abcam

ab256535

Anti-KIR2DL3 + KIR2DL1 + KIR2DL2 antibody [EPR22541-47] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR22541-47
Purification technique	Affinity purification Protein A
Concentration	1.029 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human, Recombinant fragment
Dilution info	-

Notes	-

Flow Cyt

Tested

Species	Human, Recombinant fragment
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human, Recombinant fragment
Dilution info	-
Notes	-

Target data

See full target information KIR2DL1

Function	Receptor on natural killer (NK) cells for some HLA-C alleles such as w4 and w6. Inhibits the activity of NK cells thus preventing cell lysis.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab256535 is the carrier-free version of ab224697.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


2 product images



Flow Cytometry - Anti-KIR2DL3 + KIR2DL1 + KIR2DL2 antibody [EPR22541-47] - BSA and Azide free (ab256535)

KIR2DL3 + KIR2DL1 + KIR2DL2 Flow Cytometry staining using rabbit Anti-KIR2DL3 + KIR2DL1 + KIR2DL2 antibody

Flow cytometric analysis of human peripheral blood mononuclear cell (PBMC) cells labeling KIR2DL2+KIR2DL1+KIR2DL3 with ab224697 at 1/600 compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control. Goat anti rabbit IgG (Alexa Fluor® 488, ab150097) at 1/5000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG (Left) or ab224697 (Right). Then stained with anti-CD56 conjugated to BV421. Gated on viable cells.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab224697).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KIR2DL3 + KIR2DL1 + KIR2DL2 antibody [EPR22541-47] - BSA and Azide free (ab256535)

KIR2DL3 + KIR2DL1 + KIR2DL2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KIR2DL3 + KIR2DL1 + KIR2DL2 antibody

Immunohistochemical analysis of paraffin-embedded human endometrium tissue labeling KIR2DL2+KIR2DL1+KIR2DL3 with ab224697 at 1/500 dilution (1.13µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in natural killer (NK) cells in human endometrium (PMID: 7749980). Counterstained with hematoxylin. Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab224697).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.