abcam

ab259833

Anti-Bcl 7A antibody [EPR24162-226]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24162-226
Purification technique	Affinity purification Protein A
Concentration	0.463 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Species	Human

Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000

Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Flow Cyt (Intra)

Expected

Species	Mouse
Dilution info	1/500
Notes	-

Species	Rat
Dilution info	1/500
Notes	-

Species	Human
Dilution info	1/500
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/20000
Notes	-

Species	Rat
Dilution info	1/20000
Notes	-

Species	Human
Dilution info	1/20000
Notes	-

Target data

See full target information BCL7A

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Bcl 7A also known as B-cell CLL/lymphoma 7 protein family member A plays a role in cell regulation. It is a protein with a mass of approximately 23 kDa. Bcl 7A is expressed in several tissues notably the immune system including B cells. This expression pattern suggests its function is significant in cellular processes related to this system.

Biological function summary	Bcl 7A plays a role in apoptosis regulation affecting cell survival and death. It is part of a multiprotein complex and collaborates with other Bcl family members to maintain balance between cell survival and programmed cell death. The protein is necessary for proper lymphocyte development contributing to the immune response.
Pathways	Bcl 7A functions in the Wnt signaling pathway which is important for various developmental processes and cancer pathways. Bcl 7A interacts with other proteins like β-catenin facilitating the modulation of transcription of Wnt target genes. The protein's involvement in apoptosis influences pathways important for maintaining cellular homeostasis.
Associated diseases and disorders	Altered Bcl 7A function is associated with non-Hodgkin's lymphoma and chronic lymphocytic leukemia. Its irregular expression or mutation can lead to uncontrolled cell growth representing a step in lymphomagenesis. Bcl 7A interacts with Bcl-xL another member of the Bcl family in these contexts impacting the progression of these disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


12 product images



Immunocytochemistry/ Immunofluorescence - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Bcl 7A antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HEK293T cells labelling Bcl 7A with ab259833 at 1/50 (9.26 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing mostly nuclear staining in HEK-293T cell line is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Bcl 7A antibody

Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labelling Bcl 7A with ab259833 at 1/20000 (0.023 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Nuclear staining on human tonsil (PMID:23063359). The section was incubated with ab259833 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Bcl 7A antibody

Immunohistochemical analysis of paraffin-embedded Human B cell lymphoma tissue labelling Bcl 7A with ab259833 at 1/20000 (0.023 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Nuclear staining on human B cell lymphoma (PMID:22856870). The section was incubated with ab259833 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Flow Cytometry (Intracellular) staining using rabbit Anti-Bcl 7A antibody

Flow cytometric analysis of 4% paraformaldehyde-fixed. 90% methanol-permeabilized HEK-293 (Human embryonic kidney epithelial cell) cells labelling Bcl 7A with ab259833 at 1/500 dilution (0.1ug)(Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A was immunoprecipitated from 0.35 mg HEK-293 (human embryonic kidney epithelial cell) whole cell lysate with ab259833 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259833 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HEK-293 (human embryonic kidney epithelial cell) whole cell lysate 10 ug

Lane 2: ab259833 IP in HEK-293 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259833 in HEK-293 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds

All lanes:
Immunoprecipitation - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Predicted band size: 22 kDa


Observed band size: 37 kDa, 42 kDa



Flow Cytometry (Intracellular) - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Flow Cytometry (Intracellular) staining using rabbit Anti-Bcl 7A antibody

Flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized Neuro-2a (Mouse neuroblastoma neuroblast) cells labelling Bcl 7A with ab259833 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Bcl 7A antibody

Immunohistochemical analysis of paraffin-embedded Rat cerebrum tissue labelling Bcl 7A with ab259833 at 1/20000 (0.023 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Nuclear staining on rat cerebrum. The section was incubated with ab259833 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Bcl 7A antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Neuro-2a cells labelling Bcl 7A with ab259833 at 1/50 (9.26 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing nuclear staining in Neuro-2a cell line is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Bcl 7A antibody

Immunohistochemical analysis of paraffin-embedded Mouse cerebrum tissue labelling Bcl 7A with ab259833 at 1/20000 (0.023 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Nuclear staining on mouse cerebrum. The section was incubated with ab259833 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A was immunoprecipitated from 0.35 mg Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate with ab259833 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259833 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate 10 ug

Lane 2: ab259833 IP in Neuro-2a whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259833 in Neuro-2a whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds

All lanes:
Immunoprecipitation - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Predicted band size: 22 kDa

Observed band size: 37 kDa



Western blot - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Western blot staining using rabbit Anti-Bcl 7A antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 29213114)

Negative control：HDLM-2 (HPA database: https://www.proteinatlas.org/ENSG00000110987-BCL7A/celltype)

Exposure time: Lanes 1-2:3 seconds；Lanes 3-:5：15 seconds.

All lanes:
Western blot - Anti-Bcl 7A antibody [EPR24162-226] (ab259833) at 1/1000 dilution

Lane 1:
Daudi (human burkitt's lymphoma lymphoblast) whole cell lysate at 20 µg

Lane 2:
HEK-293 (human embryonic kidney epithelial cell) whole cell lysate at 20 µg

Lane 3:
HDLM-2 (human hodgkin lymphoma) whole cell lysate at 20 µg

Lane 4:
HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg


Lane 5:
Neuro-2a (mouse neuroblastoma neuroblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 22 kDa

Observed band size: 37 kDa, 42 kDa



Western blot - Anti-Bcl 7A antibody [EPR24162-226] (ab259833)

Bcl 7A Western blot staining using rabbit Anti-Bcl 7A antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID: 29213114)

Exposure time: 15 seconds.

All lanes:
Western blot - Anti-Bcl 7A antibody [EPR24162-226] (ab259833) at 1/1000 dilution

Lane 1:
Mouse E17 brain tissue lysate at 20 µg

Lane 2:
Mouse cerebral cortex tissue lysate at 20 µg

Lane 3:
Rat hippocampus tissue lysate at 20 µg

Lane 4:
C6 (rat glial tumor glial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 22 kDa

Observed band size: 37 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.