abcam

ab259855

Anti-CENPB antibody [EPR24047-64]

Anti-CENPB antibody [EPR24047-64] (ab259855) is a rabbit monoclonal antibody detecting CENPB in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24047-64
Purification technique	Affinity purification Protein A
Concentration	0.541 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information CENPB

Function	Interacts with centromeric heterochromatin in chromosomes and binds to a specific 17 bp subset of alphoid satellite DNA, called the CENP-B box (PubMed:11726497). May organize arrays of centromere satellite DNA into a higher-order structure which then directs centromere formation and kinetochore assembly in mammalian chromosomes (Probable).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-CENPB antibody [EPR24047-64] (ab259855) is a rabbit recombinant monoclonal antibody and is validated for use

in Western Blot (WB), Flow Cytometry (Intra), Flow Cytometry (Flow Cyt), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of CENPB?
Anti-CENPB [EPR24047-64] (ab259855) specifically detects a band for CENPB (UniProt: P07199) at a molecular weight of 65kDa.

Specificity confirmed
The specificity of Anti-CENPB antibody [EPR24047-64] (ab259855) has been confirmed by Western blot testing in CENPB Knockout A431 cells.

Other related products
We have a range of other formats of antibody clone [EPR24047-64] also available for your convenience: ab259855, Carrier free - ab284394, Alexa Fluor® 488 - ab300674, Alexa Fluor® 647 - ab300733

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CENP-B also known as Centromere Protein B is a protein with a molecular weight of approximately 80 kDa. It binds to the centromeric DNA and plays a critical role in centromere formation and function. CENP-B is heavily expressed in dividing cells particularly in tissues where cell proliferation is high such as bone marrow and epithelial tissues. Researchers often use tools like anti-CENP-B antibodies and CENP-B ELISA kits to study its expression in various experimental settings.
Biological function summary	CENP-B participates in the assembly of the centromere protein complex contributing to the stability and maintenance of the centromere structure during cell division. It acts as a foundation for kinetochore nucleation ensuring proper chromosome segregation. CENP-B is involved in the formation of the CENP-A nucleosome complex important for accurate chromosomal alignment and separation.

Pathways	CENP-B is essential in the mitotic spindle assembly and chromosome segregation pathways. This protein interacts with CENP-A a variant histone which integrates into centromeric nucleosomes to initiate the formation of functional kinetochores. Additionally CENP-B interacts with CENP-C facilitating the recruitment of other centromere-associated proteins needed for chromosome dynamics during mitosis.
Associated diseases and disorders	CENP-B is primarily associated with autoimmune disorders such as systemic sclerosis (scleroderma). Patients with these conditions often produce anti-CENP-B antibodies which serve as diagnostic markers. Moreover CENP-B's dysregulation links to chromosomal instability a contributor to cancer. Research has shown connections between CENP-B and related proteins like CENP-A in the context of chromosomal aneuploidies and malignancies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

11 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CENPB antibody


Immunohistochemical analysis of paraffin-embedded: Image A -Wild-type A431 (Human epidermoid carcinoma epithelial cell) cell pellet; Image B - CENPB knockout A431 cell pellet labeling CENPB with ab259855 at 1/100 (5.34 µg/ml) dilution, followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).
Positive staining on (A) wild-type A431 cell pellet, no staining on (B) CENPB knockout A431 cell pellet. The primary antibody was incubated for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Counterstained with Hematoxylin.

Secondary antibody only control : Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CENPB antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HeLa cells labelling CENPB with ab259855 at 1/50 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 µg/mL dilution (Green). Confocal image showing centromere staining in HeLa cell line. is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5µg/mL dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 µg/mL dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CENPB antibody

Immunohistochemical analysis of paraffin-embedded Human ovarian carcinoma tissue labelling CENPB with ab259855 at 1/100 (5.39 µg/mL) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used. Positive staining on centromere in human ovarian carcinoma (PMID: 12839935).The section was incubated with ab259855 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CENPB antibody

Immunohistochemical analysis of paraffin-embedded Human gastric cancer tissue labelling CENPB with ab259855 at 1/100 (5.39 µg/mL) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used. Positive staining on centromere in human gastric cancer (PMID: 12839935). The section was incubated with ab259855 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CENPB antibody

Immunohistochemical analysis of paraffin-embedded Human pancreas tissue labelling CENPB with ab259855 at 1/100 (5.39 µg/mL) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used. Positive staining on centromere in human pancreas (PMID: 12839935). The section was incubated with ab259855 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Flow Cytometry (Intracellular) staining using rabbit Anti-CENPB antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HeLa (human cervix adenocarcinoma epithelial cell) cells labelling CENPB with ab259855 at 1/50 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB was immunoprecipitated from 0.35 mg HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate 10 µg with ab259855 at 1/30 dilution (2 µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259855 at 1/1000 dilution. VeriBlot for IP secondary antibody (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate 10 ug

Lane 2: ab259855 IP in HeLa whole cell lysate


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259855 in HeLa whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds.

All lanes:
Immunoprecipitation - Anti-CENPB antibody [EPR24047-64] (ab259855)

Predicted band size: 65 kDa



Immunoprecipitation - Anti-CENPB antibody [EPR24047-64] (ab259855)


CENPB was immunoprecipitated from 0.35 mg NIH/3T3 (mouse embryonic fibroblast) whole cell lysate 10 µg with ab259855 at 1/30 dilution (2 µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259855 at 1/1000 dilution. VeriBlot for IP secondary antibody (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: NIH/3T3 (mouse embryonic fibroblast) whole cell lysate 10 ug
Lane 2: ab259855 IP in NIH/3T3 whole cell lysate
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259855 in NIH/3T3 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes

All lanes:
Immunoprecipitation - Anti-CENPB antibody [EPR24047-64] (ab259855)

Predicted band size: 65 kDa



Western blot - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Western blot staining using rabbit Anti-CENPB antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST. Lysates were made freshly and used in WB test immediately to minimize protein degradation. Fresh lysates are preferred in this product. Exposure time: 3 minutes
All lanes:
Western blot - Anti-CENPB antibody [EPR24047-64] (ab259855) at 1/1000 dilution

Lane 1:
HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

Lane 2:
HEK-293T (human embryonic kidney epithelial cell) whole cell lysate

Lane 2:
Western blot - Human CENPB knockout A-431 cell line (ab274919)

Lane 3:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate

Lane 4:
PC-12 (rat adrenal gland pheochromocytoma) whole cell lysate

Secondary


Lane 1:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated

Lanes 2 - 4:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Lanes 2 - 3:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/50000 dilution

Lane 4:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 65 kDa



Western blot - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Western blot staining using rabbit Anti-CENPB antibody

Lanes 1-4: Merged signal (red and green). Green - ab259855 observed at 75 kDa. Red - loading control ab8245 observed at 36 kDa.
Blocking and diluting buffer and concentration: Intercept® (TBS) Blocking Buffer diluted with an equal volume of 0.1% TBST.

ab259855 Anti-CENPB antibody [EPR24047-64] was shown to specifically react with CENPB in wild-type A431 cells. Loss of signal was observed when the knockout cell line ab274919 (knockout cell lysate ab274977) was used. Wild-type and CENPB knockout samples were subjected to SDS-PAGE. ab259855 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) were incubated at 4°C overnight at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.

Fresh lysates are preferred in this product.

All lanes:
Western blot - Anti-CENPB antibody [EPR24047-64] (ab259855) at 1/1000 dilution

Lane 1:
Wild-type A431 (human epidermoid carcinoma epithelial cell) whole cell lysate

Lane 2:
CENPB knockout A431 whole cell lysate


Lane 3:
Jurkat (human T cell leukemia T lymphocyte) whole cell lysate

Lane 4:
HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate

Secondary

All lanes:
Western blot - Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) at 1/10000 dilution

Predicted band size: 65 kDa



Western blot - Anti-CENPB antibody [EPR24047-64] (ab259855)

CENPB Western blot staining using rabbit Anti-CENPB antibody

Blocking and diluting buffer and concentration: Intercept® (TBS) Blocking Buffer diluted with an equal volume of 0.1% TBS
Lanes 1-4: Merged signal (red and green). Green - ab259855 observed at 75 kDa. Red - loading control ab8245 observed at 36 kDa.

ab259855 Anti-CENPB antibody [EPR24047-64] was shown to specifically react with CENPB in wild-type A431 cells. Loss of signal was observed when the knockout cell line ab274919 (knockout cell lysate ab274977) was used. Wild-type and CENPB knockout samples were subjected to SDS-PAGE. ab259855 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) were incubated at 4°C overnight at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.

Fresh lysates are preferred in this product.

All lanes:
Western blot - Anti-CENPB antibody [EPR24047-64] (ab259855) at 1/1000 dilution

Lane 1:
Wild-type A431 (human epidermoid carcinoma epithelial cell) whole cell lysate

Lane 2:
CENPB knockout A431 whole cell lysate

Lane 3:
Jurkat (human T cell leukemia T lymphocyte) whole cell lysate

Lane 4:
HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate

Secondary

All lanes:
Goat Anti-Rabbit IgG H&L (IRDye® 800CW) (ab216773) and Goat Anti-Mouse IgG H&L (IRDye® 680RD) (ab216776) at 1/10000 dilution

Predicted band size: 65 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.