abcam

ab259934

Anti-MICA + MICB antibody [EPR24086-121]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24086-121
Purification technique	Affinity purification Protein A
Concentration	0.457 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information MICA

Function	Widely expressed membrane-bound protein which acts as a ligand to stimulate an activating receptor KLRK1/NKG2D, expressed on the surface of essentially all human natural killer (NK), gammadelta T and CD8 alphabeta T-cells (PubMed:11491531, PubMed:11777960). Up-regulated in stressed conditions, such as viral and bacterial infections or DNA damage response, serves as signal of cellular stress, and engagement of KLRK1/NKG2D by MICA triggers NK-cells resulting in a range of immune effector functions, such as cytotoxicity and cytokine production (PubMed:10426993).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	MICA and MICB also referred to as Mhc class I polypeptide-related sequence A and B are stress-induced proteins and are known to have a molecular weight of approximately 62 kDa and 70 kDa respectively. They are expressed on the surface of many cell types especially epithelial cells fibroblasts and upon certain cancers. These proteins are encoded within the MHC class I region yet unlike classic MHC class I molecules they do not associate with beta-2-microglobulin and they do not present antigens to T cells. Instead MICA and MICB can activate immune responses by engaging the NKG2D receptor on natural killer (NK) cells CD8+ T cells and gamma-delta T cells.
Biological function summary	MICA and MICB contribute to the immune surveillance system by marking stressed or transformed cells for destruction. These proteins are not part of a complex but are known ligands for the NKG2D receptor. Their interaction with NKG2D triggers the cytotoxic activity of NK cells and CD8+ T cells enhancing the destruction of cancerous or infected cells. Especially interesting is their role in tumor immunology where they serve as a danger signal to the immune system pointing NK cells toward tumor cells that would otherwise evade detection.
Pathways	The role of MICA and MICB centers around the immune response and cell stress pathways. One important pathway is the stress-induced signal transduction which upregulates these proteins in response to cellular stress such as heat shock. They fit into the immune surveillance pathway interacting with proteins like the NKG2D receptor and other signaling molecules that modulate immune cell activation. The pathway emphasizes the removal of damaged or abnormal cells and this interaction orchestrates timely immune responses.
Associated diseases and disorders	MICA and MICB have associations with certain cancers like liver and colorectal cancers where they become over-expressed and shed from tumor cells to escape immune detection. Additionally their presence links to autoimmune diseases such as rheumatoid arthritis where the immune system attacks the body's own cells unjustly. The shedding of MICA/MICB from tumor cells involves enzymes like

metalloproteinases affecting the engagement of NKG2D and complicating the immune response to cancerous cells.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images




Flow Cytometry - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

MICA + MICB Flow Cytometry staining using rabbit Anti-MICA + MICB antibody

Flow cytometric analysis of THP-1 (Human monocytic leukemia monocyte, left)/ HeLa (human cervix adenocarcinoma epithelial cell, right) cells labelling MICA with ab259934 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Negative control: THP-1 (PMID: 28154561).

Gated on viable cells.



Immunoprecipitation - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

MICA was immunoprecipitated from 0.35 mg HUVEC (human umbilical vein endothelial cell) whole cell lysate with ab259934 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259934 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: HUVEC (human umbilical vein endothelial cell) whole cell lysate 10 ug

Lane 2: ab259934 IP in HUVEC whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259934 in HUVEC whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 seconds.

This blot was developed using a higher sensitivity ECL substrate.

All lanes:
Immunoprecipitation - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

Observed band size: 60 kDa



Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

Blocking and diluting buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934) at 1/1000 dilution

Lane 1:
His-tagged human recombinant protein MHC class I polypeptide-related sequence A (24-297aa) at 0.05 µg


Lane 2:
His-tagged human recombinant protein MHC class I polypeptide-related sequence B (24-297aa) at 0.1 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 43 kDa

Observed band size: 60 kDa



Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

MICA + MICB Western blot staining using rabbit Anti-MICA + MICB antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Negative control: THP-1 (PMID: 28154561).

MICA is a glycoprotein and can be de-glycosylated by PNGase F. The molecular mass observed is consistent with the literature (PMID: 10359807, 9396860).

Exposure time: 3 minutes.
All lanes:
Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934) at 1/1000 dilution


Lane 1:
HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
THP-1 (Human monocytic leukemia monocyte) whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Observed band size: 60 kDa



Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
MICA is a glycoprotein and can be de-glycosylated by PNGase F. The molecular mass observed is consistent with the literature (PMID: 10359807, 9396860).

Exposure time: 3 minutes.

All lanes:
Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934) at 1/1000 dilution

Lane 1:
HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 2:
HeLa whole cell lysate treated with PNGase F at 20 µg


Lane 3:
HUVEC (human umbilical vein endothelial cell) whole cell lysate at 20 µg

Lane 4:
HUVEC whole cell lysate treated with PNGase F at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Observed band size: 32 kDa, 50-60 kDa



Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
MICA is a glycoprotein and can be de-glycosylated by PNGase F. The molecular mass observed is consistent with the literature (PMID: 10359807, 9396860).

Exposure time: 3 minutes.

All lanes:
Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934) at 1/1000 dilution


Lane 1:
Human breast cancer tissue lysate at 20 µg

Lane 2:
Human colon cancer tissue lysate at 20 µg

Secondary

All lanes:
VeriBlot for IP secondary antibody(HRP)(ab131366) at 1/1000 dilution

Observed band size: 50-60 kDa



Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-MICA + MICB antibody [EPR24086-121] (ab259934) at 1/1000 dilution

Lane 1:
His-tagged human recombinant protein MHC class I polypeptide-related sequence A, 10 ng

Lane 2:
His-tagged human recombinant protein MHC class I polypeptide-related sequence B, 10 ng

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Observed band size: 50 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.