abcam

ab259967

Anti-SIRT5 antibody [EPR23787-116]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23787-116
Purification technique	Affinity purification Protein A
Concentration	0.496 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Species	Human
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information SIRT5

Function	NAD-dependent lysine demalonylase, desuccinylase and deglutarylase that specifically removes malonyl, succinyl and glutaryl groups on target proteins (PubMed:21908771, PubMed:22076378, PubMed:24703693, PubMed:29180469). Activates CPS1 and contributes to the regulation of blood ammonia levels during prolonged fasting: acts by mediating desuccinylation and deglutarylation of CPS1, thereby increasing CPS1 activity in response to elevated NAD levels during fasting (PubMed:22076378, PubMed:24703693). Activates SOD1 by mediating its desuccinylation, leading to reduced reactive oxygen species (PubMed:24140062). Activates SHMT2 by mediating its desuccinylation (PubMed:29180469). Modulates ketogenesis through the desuccinylation and activation of HMGCS2 (By similarity). Has weak NAD-dependent protein deacetylase activity; however this activity may not be physiologically relevant in vivo. Can deacetylate cytochrome c (CYCS) and a number of other proteins in vitro such as UOX.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	SIRT5 also known as Sirtuin 5 is a mitochondrial protein with enzymatic activity as a lysine desuccinylase demalonylase and deglutarylase. It has a molecular mass of approximately 34 kDa. This protein is expressed highly in tissues with high metabolic activity like the liver heart and brain where it participates in regulating energy metabolism. The activity of SIRT5 affects cellular processes by removing post-translational modifications from lysine residues modulating proteins' functions and interactions in these tissues.
Biological function summary	SIRT5 functions as an important regulator of mitochondrial metabolism and influences cellular energy homeostasis. It is not part of larger complexes but it interacts with various metabolic enzymes to control their function. Through its deacylase activity SIRT5 affects the urea cycle and fatty acid oxidation by modifying key enzymes involved in these processes. Its role in maintaining a balance in energy production and consumption highlights its importance in cellular metabolism.
Pathways	SIRT5 is involved in both the tricarboxylic acid (TCA) cycle and the regulation of reactive oxygen species (ROS). Within the TCA cycle SIRT5 desuccinylates and activates enzymes like citrate synthase enhancing the cycle's efficiency. In regulating ROS SIRT5 reduces oxidative stress by affecting enzymes such as superoxide dismutase 2 (SOD2). Its interaction with these pathways and proteins reflects its role in maintaining metabolic health and cellular integrity.
Associated diseases and disorders	Alterations in SIRT5 levels have been connected to metabolic diseases such as obesity and neurodegenerative disorders like Alzheimer's disease. In obesity SIRT5 dysregulation can lead to inefficient metabolic processes and energy imbalance. In Alzheimer’s decreased SIRT5 activity may contribute to increased mitochondrial oxidative stress implicated in disease progression. These connections highlight SIRT5’s significance in both metabolic and neurodegenerative contexts linking with proteins like AMPK in obesity and tau proteins in Alzheimer's disease.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SIRT5 antibody

Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling SIRT5 with ab259967 at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human liver (PMID 29491006). Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Flow Cytometry (Intracellular) - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Flow Cytometry (Intracellular) staining using rabbit Anti-SIRT5 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HepG2 (Human hepatocellular carcinoma epithelial cell) cells labelling SIRT5 with ab259967 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SIRT5 antibody

Immunohistochemical analysis of paraffin-embedded Human ovarian adenocarcinoma tissue labeling SIRT5 with ab259967 at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human ovarian adenocarcinoma (PMID 31456942). Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Immunoprecipitation - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 was immunoprecipitated from 0.35 mg HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate with ab259967 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259967 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate 10 ug

Lane 2: ab259967 IP in HepG2 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259967 in HepG2 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

All lanes:
Immunoprecipitation - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

Predicted band size: 33 kDa

Observed band size: 30 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SIRT5 antibody

Immunohistochemical analysis of paraffin-embedded Mouse liver tissue labeling SIRT5 with ab259967 at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse liver (PMID 29491006). Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Flow Cytometry (Intracellular) - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Flow Cytometry (Intracellular) staining using rabbit Anti-SIRT5 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized Hepa1-6 (Mouse hepatoma epithelial cell) cells labelling SIRT5 with ab259967 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SIRT5 antibody

Immunohistochemical analysis of paraffin-embedded Rat liver tissue labeling SIRT5 with ab259967 at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on rat liver (PMID 29491006). Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Immunoprecipitation - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 was immunoprecipitated from 0.35 mg Mouse liver tissue lysate with ab259967 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab259967 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Mouse liver tissue lysate 10 ug

Lane 2: ab259967 IP in Mouse liver tissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab259967 in mouse liver tissue lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

All lanes:
Immunoprecipitation - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

Predicted band size: 33 kDa

Observed band size: 30 kDa



Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Western blot staining using rabbit Anti-SIRT5 antibody

Western blot: Anti-SIRT5 antibody [EPR23787-116] (ab259967) staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (ab7291) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab259967 was shown to bind specifically to SIRT5. A band was observed at 33 kDa in wild-type A549 cell lysates with no signal observed at this size in SIRT5 knockout cell line. To generate this image, wild-type and SIRT5 knockout A549 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in fluorescent western blot (TBS-based) blocking solution before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967) at 1/1000 dilution

Lane 1:
Wild-type A549 cell lysate at 20 µg

Lane 2:
SIRT5 knockout A549 cell lysate at 20 µg

Lane 3:
PC-3 cell lysate at 20 µg

Lane 4:
MCF7 Membrane Prep cell lysate at 20 µg


Secondary

Lanes 1 - 4:
Goat anti-Rabbit IgG H&L 800CW at 1/20000 dilution

Lanes 1 - 4:
Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 33 kDa



Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

SIRT5 Western blot staining using rabbit Anti-SIRT5 antibody

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967) at 1/1000 dilution

Lane 1:
Human liver tissue lysate at 10 µg


Lane 2:
Human heart tissue lysate at 10 µg

Secondary

All lanes:
VeriBlot for IP secondary antibody(HRP)(ab131366) at 1/1000 dilution

Predicted band size: 33 kDa

Observed band size: 30 kDa



Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967) at 1/1000 dilution

Lane 1:
HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Huh7 (human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 4:
HEK-293T (human embryonic kidney epithelial cell) whole cell lysate at 20 µg

Lane 5:
Hepa1-6 (mouse hepatoma epithelial cell) whole cell lysate at 20 µg

Lane 6:
RAW 264.7 (mouse abelson murine leukemia virus-induced tumor macrophage) whole cell lysate at 20 µg


Lane 7:
C6 (rat glial tumor glial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Predicted band size: 33 kDa

Observed band size: 30 kDa



Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967)

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 48 seconds.

All lanes:
Western blot - Anti-SIRT5 antibody [EPR23787-116] (ab259967) at 1/1000 dilution

Lane 1:
Mouse liver tissue lysate at 20 µg

Lane 2:
Mouse heart tissue lysate at 20 µg

Lane 3:
Rat liver tissue lysate at 20 µg

Lane 4:
Rat heart tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Predicted band size: 33 kDa

Observed band size: 30 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.