abcam

ab263950

Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR22599-78
Purification technique	Affinity purification Protein A
Specificity	STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13.
Concentration	1.011 - 1.039 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt (Intra)

Tested

Species	Mouse, Human
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Rat, Mouse, Synthetic peptide
Dilution info	-
Notes	-

IP

Tested

Species	Human, Mouse
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

ChIP

Not recommended

Species	Mouse, Rat, Human, Synthetic peptide
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	p-STAT6 expression is typically undetectable or present at very low levels in healthy tissue lysates (e.g., kidney, lung, and skin) under physiological conditions (PMID: 34831280; PMID: 22401596; PMID: 24955743; PMID: 36629780). However, its expression is markedly upregulated in pathological states, including inflammatory responses and oncogenic activation (PMID: 36884218).

Species	Human
Dilution info	1/1000
Notes	p-STAT6 expression is typically undetectable or present at very low levels in healthy tissue lysates (e.g., kidney, lung, and skin) under physiological conditions (PMID: 34831280; PMID: 22401596; PMID: 24955743; PMID: 36629780). However, its expression is markedly upregulated in pathological states, including inflammatory responses and oncogenic activation (PMID: 36884218).

Species	Rat
Dilution info	1/1000
Notes	p-STAT6 expression is typically undetectable or present at very low levels in healthy tissue lysates (e.g., kidney, lung, and skin) under physiological conditions (PMID:

34831280; PMID: 22401596; PMID: 24955743; PMID: 36629780). However, its expression is markedly upregulated in pathological states, including inflammatory responses and oncogenic activation (PMID: 36884218).

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

Dot

Tested

Species	Synthetic peptide
Dilution info	-
Notes	-

Expected

Species	Mouse, Human, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Not recommended

Species	Mouse, Rat, Human, Synthetic peptide
Dilution info	-
Notes	-

Target data

See full target information Stat6 pY641

Function	Carries out a dual function: signal transduction and activation of transcription. Involved in IL4/interleukin-4- and IL3/interleukin-3-mediated signaling.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab263950 is the carrier-free version of ab263947.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	STAT6 also known as Signal Transducer and Activator of Transcription 6 is a transcription factor with a mass of around 94 kDa. This protein is an important component in the signaling pathways that respond to cytokines especially those of

	the IL-4 and IL-13 receptors. STAT6 mainly expresses in immune cells including T cells B cells and macrophages. It exists in the cytoplasm until the cytokine stimulation prompts its phosphorylation and translocation to the nucleus where it influences gene expression.
Biological function summary	STAT6 plays a central role in the development and function of Th2 cells a subset of T helper cells. The protein controls genes linked to the immune response and inflammation such as IL-4 and IL-13 by binding to specific DNA sequences as part of the transcriptional complex. These processes are critical for antibody class switching in B cells and also for regulating the differentiation of T cells which are processes important in defending against parasitic infections and are linked to allergies.
Pathways	STAT6 functions mainly in the JAK-STAT signaling pathway a pathway important for transmitting information from chemical signals outside cells to the cell nucleus resulting in DNA transcription. Within this pathway STAT6 works closely with JAK1 and JAK3 kinases which phosphorylate STAT6 enabling its activation. The protein also connects with other members of the STAT family like STAT1 and STAT3 during overlapping signaling processes.
Associated diseases and disorders	STAT6 has associations with asthma and certain types of cancer such as colorectal cancer. In asthma STAT6 influences the production of IgE and eosinophilic inflammation often connected to allergic responses. Meanwhile STAT6 promotes tumorigenesis in some cancers by regulating genes responsible for cell survival and proliferation. It connects with proteins like IL-4 and IL-13 in these disorders which suggests potential therapeutic targets for treatment.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images




Flow Cytometry (Intracellular) - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

STAT6 (phospho Y641) Flow Cytometry (Intracellular) staining using rabbit Anti-STAT6 (phospho Y641) antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Daudi (Human Burkitt's lymphoma lymphoblast) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min (Red)/Untreated control (Green) cells labelling STAT6 (phospho Y641) with ab263947 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730)isotype control (black)and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).



Flow Cytometry (Intracellular) - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

STAT6 (phospho Y641) Flow Cytometry (Intracellular) staining using rabbit Anti-STAT6 (phospho Y641) antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Daudi (Human Burkitt's lymphoma lymphoblast) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min (Red)/Untreated control (Green) cells labelling STAT6 (phospho Y641) with ab263947 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730)isotype control (black)and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13. This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).



Immunoprecipitation - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

STAT6 (phospho Y641) was immunoprecipitated from 0.35 mg Daudi (Human Burkitt's lymphoma lymphoblast) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate 10ug with ab263947 at 1:30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab263947 at 1:1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1:5000 dilution.
Lane 1: Daudi (Human Burkitt's lymphoma lymphoblast) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate 10ug

Lane 2: ab263947 IP in Daudi was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab263947 in Daudi was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 1 min

STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).
All lanes:
Immunoprecipitation - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] (ab263947)

Predicted band size: 94 kDa




Flow Cytometry (Intracellular) - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

STAT6 (phospho Y641) Flow Cytometry (Intracellular) staining using rabbit Anti-STAT6 (phospho Y641) antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized RAW 264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min (Red) / Untreated control (Green) cells labelling STAT6 (phospho Y641) with ab263947 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black)and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).



Flow Cytometry (Intracellular) - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

STAT6 (phospho Y641) Flow Cytometry (Intracellular) staining using rabbit Anti-STAT6 (phospho Y641) antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized RAW 264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min (Red) / Untreated control (Green) cells labelling STAT6 (phospho Y641) with ab263947 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black)and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13. This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).



Immunoprecipitation - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

STAT6 (phospho Y641) was immunoprecipitated from 0.35 mg RAW 264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate 10ug with ab263947 at 1:30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab263947 at 1:1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1:5000 dilution.
Lane 1: RAW 264.7 (Mouse Abelson murine leukemia virus-induced tumor macrophage) was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate 10ug

Lane 2: ab263947 IP in RAW 264.7 was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab263947 in RAW 264.7 was serum starved for 24h, then treated with 100ng/ml IL-4 for 15min whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 2 min

STAT6 is activated via phosphorylation at Tyr641 and is required for responsiveness to IL-4 and IL-13.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).

All lanes:
Immunoprecipitation - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] (ab263947)

Predicted band size: 94 kDa



Dot Blot - Anti-STAT6 (phospho Y641) antibody [EPR22599-78] - BSA and Azide free (ab263950)

Dot blot analysis of STAT6 (phospho Y641) peptide (Lane 1 STAT6 non-phospho peptide peptide (Lane 2), labelling STAT 6 (pY641) with purified ab263947 at a dilution of 1/1000. ab97051 (Peroxidase conjugated goat anti-rabbit IgG (H+L)) was used as the secondary antibody at a dilution of 1/100000.
Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 26 seconds.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab263947).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.