abcam

ab264098

Anti-GAS 6 antibody [EPR23931-27]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23931-27
Purification technique	Affinity purification Protein A
Concentration	0.547 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information GAS6

Function	Ligand for tyrosine-protein kinase receptors AXL, TYRO3 and MER whose signaling is implicated in cell growth and survival, cell adhesion and cell migration. GAS6/AXL signaling plays a role in various processes such as endothelial cell survival during acidification by preventing apoptosis, optimal cytokine signaling during human natural killer cell development, hepatic regeneration, gonadotropin-releasing hormone neuron survival and migration, platelet activation, or regulation of thrombotic responses. (Microbial infection) Can bridge virus envelope phosphatidylserine to the TAM receptor tyrosine kinase Axl to mediate viral entry by apoptotic mimicry (PubMed:21501828). Plays a role in Dengue cell entry by apoptotic mimicry (PubMed:23084921). Plays a role in Vaccinia virus cell entry by apoptotic mimicry (PubMed:21501828). Plays a role in ebolavirus and marburgvirus cell entry by apoptotic mimicry (PubMed:17005688).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The protein Gas6 also known as growth arrest-specific 6 weighs approximately 75 kDa and functions mechanically as a ligand for the TAM (Tyro3 Axl and Mer) receptor tyrosine kinases. Gas6 protein facilitates cell signaling by binding especially to AXL receptor initiating various downstream cellular responses. It is widely expressed in different tissues including the nervous system cardiovascular system and kidneys. Additionally its expression increases in response to cellular stress and inflammatory stimuli.
Biological function summary	Gas6 influences several cellular processes such as cell survival proliferation and migration. Gas6 exerts its effects in a complex with AXL receptor which is part of a larger network involving other TAM receptors. By activating AXL Gas6 modulates immune responses inflammation and tissue repair. The protein also plays a significant role in phagocytosis helping remove apoptotic cells thereby maintaining tissue homeostasis.
Pathways	The Gas6 protein is significantly engaged in the PI3K/AKT signaling pathway important for promoting cell survival and inhibiting apoptosis. Additionally Gas6 interacts with the NF-kB pathway which regulates inflammation and immune responses. Both pathways highlight the synergy between Gas6 and AXL emphasizing their contribution to cellular integrity and response to injury.
Associated diseases and disorders	Gas6's dysregulation associates with cancer and atherosclerosis. In oncogenesis altered Gas6-AXL signaling promotes tumor progression and resistance to therapy implicating it in diverse cancers. Meanwhile in cardiovascular diseases elevated Gas6 levels correlate with atherosclerosis interacting with inflammatory proteins such as interleukins. Understanding Gas6's role in these conditions offers potential for therapeutic targeting and biomarker development.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


6 product images



Flow Cytometry (Intracellular) - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

GAS 6 Flow Cytometry (Intracellular) staining using rabbit Anti-GAS 6 antibody


Flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized THP-1 (Human monocytic leukemia monocyte) cells labelling GAS 6 with ab264098 at 1/500 dilution (Right) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Left). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab264098.



Flow Cytometry (Intracellular) - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

GAS 6 Flow Cytometry (Intracellular) staining using rabbit Anti-GAS 6 antibody


Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized THP-1 (Human monocytic leukemia monocyte) cells labelling GAS 6 with ab264098 at 1/500 dilution (0.1ug)/ Right (Red) compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab264098. Cells were co-stained with DRAQ/ Ki-67 to differentiate cell cycle phase.



Flow Cytometry (Intracellular) - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

GAS 6 Flow Cytometry (Intracellular) staining using rabbit Anti-GAS 6 antibody

Flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized Jurkat (Human T cell leukemia T lymphocyte (Left) / THP-1 (Human monocytic leukemia monocyte, Right) cells labelling GAS 6 with ab264098 at 1/500 dilution (Right). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. **Negative control**: Jurkat (PMID:10400186)



Immunoprecipitation - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

GAS 6 was immunoprecipitated from 0.35 mg THP-1 (human monocytic leukemia monocyte) whole cell lysate with ab264098 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab264098 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: THP-1 (human monocytic leukemia monocyte) whole cell lysate 10 ug

Lane 2: ab264098 IP in THP-1 whole cell lysate

Lane 3: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated: Rabbit monoclonal IgG (ab172730) instead of ab264098 in THP-1 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds

All lanes:
Immunoprecipitation - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

Predicted band size: 80 kDa

Observed band size: 78 kDa



Western blot - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

GAS 6 Western blot staining using rabbit Anti-GAS 6 antibody

False colour image of Western blot: Anti-GAS 6 antibody [EPR23931-27] staining at 1/1000 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (ab7291) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab264098 was shown to bind specifically to GAS 6. A band was observed at 75 kDa in wild-type HEK-293 cell lysates with no signal observed at this size in GAS6 knockout cell line ab275837 (knockout cell lysate ab275811). To generate this image, wild-type and GAS6 knockout HEK-293 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.
All lanes:
Western blot - Anti-GAS 6 antibody [EPR23931-27] (ab264098) at 1/1000 dilution

Lane 1:
Wild-type HEK-293 cell lysate at 20 µg

Lane 2:
GAS6 knockout HEK-293 cell lysate at 20 µg


Lane 2:
Western blot - Human GAS6 knockout HEK-293 cell line (ab275837)

Lane 3:
THP-1 cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 80 kDa

Observed band size: 75 kDa



Western blot - Anti-GAS 6 antibody [EPR23931-27] (ab264098)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Negative control: Jurkat (PMID:10400186)

Samples are non-boiled as boiling may cause protein aggregates.

Exposure time: Lanes 1-2: 48 seconds;Lanes 3-4: 3 minutes.

All lanes:
Western blot - Anti-GAS 6 antibody [EPR23931-27] (ab264098) at 1/1000 dilution

Lane 1:
THP-1 (human monocytic leukemia monocyte) whole cell lysate at 20 µg

Lane 2:
Jurkat (human t cell leukemia T lymphocyte) whole cell lysate at 20 µg

Lane 3:
SH-SY5Y (human neuroblastoma epithelial cell) whole cell lysate at 20 µg

Lane 4:
RPMI-8226 (human plasmacytoma, myeloma B lymphocyte) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 80 kDa

Observed band size: 78 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.