abcam

ab264563

Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR23059-124
Purification technique	Affinity purification Protein A
Concentration	1.041 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	-

Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information OPTN

Function	Plays an important role in the maintenance of the Golgi complex, in membrane trafficking, in exocytosis, through its interaction with myosin VI and Rab8 (PubMed:27534431). Links myosin VI to the Golgi complex and plays an important role in Golgi ribbon formation (PubMed:27534431). Plays a role in the activation of innate immune response during viral infection. Mechanistically, recruits TBK1 at the Golgi apparatus, promoting its trans-phosphorylation after RLR or TLR3 stimulation (PubMed:27538435). In turn, activated TBK1 phosphorylates its downstream partner IRF3 to produce IFN-beta/IFNB1. Plays a neuroprotective role in the eye and optic nerve. May act by regulating membrane trafficking and cellular morphogenesis via a complex that contains Rab8 and huntingtin (HD). Mediates the interaction of Rab8 with the probable GTPase-activating protein TBC1D17 during Rab8-mediated endocytic trafficking, such as that of transferrin receptor (TFRC/TfR); regulates Rab8 recruitment to tubules emanating from the endocytic recycling compartment (PubMed:22854040). Autophagy receptor that interacts directly with both the cargo to become degraded and an autophagy modifier of the MAP1 LC3 family; targets ubiquitin-coated bacteria (xenophagy), such as cytoplasmic Salmonella enterica, and appears to function in the same pathway as SQSTM1 and CALCOCO2/NDP52. (Microbial infection) May constitute a cellular target for various viruses, such as adenovirus E3 14.7 or Bluetongue virus, to inhibit innate immune response (PubMed:27538435, PubMed:9488477). During RNA virus infection, such as that of Sendai virus, negatively regulates the induction of IFNB1 (PubMed:20174559).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab264563 is the carrier-free version of ab242146.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Optineurin also known as OPTN is a multifunctional protein of approximately 67 kDa. It is expressed in various tissues including the retina where it plays a role in maintaining cellular functions. Optineurin interacts with several binding partners to execute its roles within the cell. It is involved in intracellular trafficking signaling and inflammation processes. As an essential component of the cellular machinery optineurin aids in proper cellular communication and responses.
Biological function summary	Optineurin influences different cellular processes notably autophagy and regulation of inflammation. It participates in forming a complex with other proteins to facilitate cargo transport and degradation via the autophagy-lysosome pathway. This function is important for removing damaged proteins and organelles ensuring cellular homeostasis. Inflammation regulation occurs through optineurin's ability to inhibit certain signaling pathways therefore modulating immune responses and preventing excessive inflammation.
Pathways	Optineurin plays a role in the NF-kB signaling pathway and the autophagic process. In NF-kB signaling it acts as a negative regulator by degrading specific signaling molecules. This regulation is necessary to prevent chronic inflammation and maintain immune balance. Within autophagy optineurin collaborates with other autophagy-related proteins like LC3 to facilitate the sequestration and clearance of cellular components preventing accumulation of damaged cellular materials.
Associated diseases and disorders	Optineurin connects to conditions like glaucoma and amyotrophic lateral sclerosis (ALS). Mutations in the optineurin gene link to glaucoma as they can impair autophagy and lead to optic nerve damage. In the context of ALS optineurin mutations disrupt normal cellular functions contributing to motor neuron degeneration. The protein TDP-43 is often associated with ALS and interacts with optineurin in pathological processes affecting disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

6 product images




Immunocytochemistry/ Immunofluorescence - Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free (ab264563)

Optineurin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Optineurin antibody

This data was developed using the same antibody clone in a different buffer formulation (ab242146) ab242146 was shown to react with OPTN in wild-type U20S cells in immunocytochemistry with loss of signal observed in a OPTN knockout cell line. Wild-type and knockout cells were mixed and pelleted at a 1:1 ratio on coverslips. The cells were fixed with 4% paraformaldehyde (15 min) then permeabilized with 0.1% Triton X-100 (10min) and then blocked with 1x PBS, 0.01% Triton X-100, 5% BSA, 5% NGS. The cells were then incubated with ab242146 at dilution overnight at 4°C followed by a further incubation at room temperature for 1h with a goat anti-rabbit secondary antibody to (Alexa Fluor® 555) at 0.5 µg/ml. Acquisition of the green (wild-type), red (antibody staining) and far-red (knockout) channels was performed. Representative grayscale images of the red channel are shown. Wild-type and knockout cells are outlined with yellow and magenta dashed line, respectively. Schematic representation of the mosaic strategy used is shown on the bottom-right panel. Image was acquired with a Zeiss(LSM-880).

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free (ab264563)

Optineurin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Optineurin antibody

Immunohistochemical analysis of paraffin-embedded Human cerebrum tissue labeling Optineurin with ab242146 at 1/500 dilution (0.98 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Cytoplasmic staining on human cerebrum tissue. The section was incubated with ab242146 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND®RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20mins.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab242146).



Flow Cytometry (Intracellular) - Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free (ab264563)

Optineurin Flow Cytometry (Intracellular) staining using rabbit Anti-Optineurin antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HeLa (human cervix adenocarcinoma) cells labelling Optineurin with ab242146 at 1/50 (Red) compared with a Rabbit monoclonal IgG (ab172730, Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor®488, ab150077) at 1/2000 dilution was used as the secondary antibody. This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab242146).



Immunoprecipitation - Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free (ab264563)

This data was developed using the same antibody clone in a different buffer formulation (ab242146)

Immunoprecipitation of OPTN in U20S cells. Lysates were prepared and immunoprecipitation was performed using 1.0 µg of ab242146 pre-coupled to Protein A beads. Samples were washed and processed for western blot with OPTN Monoclonal Antibody at 1/10000


This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.
Lane 1:
Immunoprecipitation at 1/10000 dilution

Lane 3:
Immunoprecipitation - Anti-Optineurin antibody [EPR23059-124] (ab242146) at 1 µg

All lanes:

U20S cells

Observed band size: 66 kDa



Western blot - Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free (ab264563)

Optineurin Western blot staining using rabbit Anti-Optineurin antibody

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab242146).

Western blot: Anti-Optineurin antibody [EPR23059-124] ab242146 staining at 1/1000 dilution, shown in green; Mouse anti GAPDH ab8245 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 73 kDa in Wild-type MCF7 cell lysates with no signal observed at this size in OPTN knockout MCF7 cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3pc Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-Optineurin antibody [EPR23059-124] (ab242146) at 1/1000 dilution

Lane 1:
Wild-type MCF7 whole cell lysate at 20 µg

Lane 2:
OPTN knockout MCF7 whole cell lysate at 20 µg

Lane 3:
U-2 OS whole cell lysate at 20 µg

Lane 4:
Raji whole cell lysate at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 66 kDa

Observed band size: 73 kDa



Western blot - Anti-Optineurin antibody [EPR23059-124] - BSA and Azide free (ab264563)

Optineurin Western blot staining using rabbit Anti-Optineurin antibody

This data was developed using the same antibody clone in a different buffer formulation (ab242146)

ab242146 was shown to react with OPTN in wild-type U-2 OS cells in Western blot with loss of signal observed in OPTN knockout cell line ab261906. Wild-type U-2 OS and OPTN knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab242146 overnight at 4 °C at a 1/1000 dilution. Blots were incubated with secondary antibodies at 0.2ug/mL before imaging.

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.
All lanes:
Western blot - Anti-Optineurin antibody [EPR23059-124] (ab242146) at 1/1000 dilution

Lane 1:
Wild-type U-2 OS cell lysate at 20 µg

Lane 2:
OPTN knockout U-2 OS cell lysate at 20 µg

Observed band size: 66 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.