abcam

ab267351

Anti-TFEB antibody [EPR22941-6]

Anti-TFEB antibody [EPR22941-6] (ab267351) is a rabbit monoclonal antibody detecting TFEB in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**, **ChIC/CUT&RUN sequencing**. Suitable for **Human**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | Advanced Validation | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR22941-6
Purification technique	Affinity purification Protein A
Concentration	0.524 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human

Dilution info	1/30
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Mouse
Dilution info	-
Notes	-

ChIC/CUT&RUN-seq

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information TFEB

Function

Transcription factor that acts as a master regulator of lysosomal biogenesis, autophagy, lysosomal exocytosis, lipid catabolism, energy metabolism and immune response (PubMed:21617040, PubMed:22343943, PubMed:22576015, PubMed:22692423, PubMed:25720963, PubMed:30120233, PubMed:31672913, PubMed:32612235, PubMed:32753672, PubMed:35662396, PubMed:36697823, PubMed:36749723, PubMed:37079666). Specifically recognizes and binds E-box sequences (5'-CANNTG-3'); efficient DNA-binding requires dimerization with itself or with another MiT/TFE family member such as TFE3 or MITF (PubMed:1748288, PubMed:19556463, PubMed:29146937). Involved in the cellular response to amino acid availability by acting downstream of MTOR: in the presence of nutrients, TFEB phosphorylation by MTOR promotes its cytosolic retention and subsequent inactivation (PubMed:21617040, PubMed:22343943, PubMed:22576015, PubMed:22692423, PubMed:25720963, PubMed:32612235, PubMed:32753672, PubMed:35662396, PubMed:36697823). Upon starvation or lysosomal stress, inhibition of MTOR induces TFEB dephosphorylation, resulting in nuclear localization and transcription factor activity (PubMed:22343943, PubMed:22576015, PubMed:22692423, PubMed:25720963, PubMed:32612235, PubMed:32753672, PubMed:35662396, PubMed:36697823). Specifically recognizes and binds the CLEAR-box sequence (5'-GTCACGTGAC-3') present in the regulatory region of many lysosomal genes, leading to activate their expression, thereby playing a central role in expression of lysosomal genes (PubMed:19556463, PubMed:22692423). Regulates lysosomal positioning in response to nutrient deprivation by promoting the expression of PIP4P1 (PubMed:29146937). Acts as a positive regulator of autophagy by promoting expression of genes involved in autophagy (PubMed:21617040, PubMed:22576015, PubMed:23434374, PubMed:27278822). In association with TFE3, activates the expression of CD40L in T-cells, thereby playing a role in T-cell-dependent antibody responses in activated CD4(+) T-cells and thymus-dependent humoral immunity (By similarity). Specifically recognizes the gamma-E3 box, a subset of E-boxes, present in the heavy-chain immunoglobulin enhancer (PubMed:2115126). Plays a role in the signal transduction processes required for normal vascularization of the placenta (By similarity). Involved in the immune response to infection by the bacteria S.aureus, S.typhimurium or S.enterica: infection promotes itaconate production, leading to alkylation, resulting in nuclear localization and transcription factor activity (PubMed:35662396). Itaconate-mediated alkylation activates TFEB-dependent lysosomal biogenesis, facilitating the bacteria clearance during the antibacterial innate immune response (PubMed:35662396). In association with ACSS2, promotes the expression of genes involved in lysosome biogenesis and both autophagy upon glucose deprivation (PubMed:28552616).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-TFEB antibody [EPR22941-6] (ab267351) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) and ChIC/CUT&RUN sequencing in Human samples.

What is the molecular weight of TFEB?
Anti-TFEB [EPR22941-6] (ab267351) specifically detects a band for TFEB (UniProt: P19484) at a molecular weight of 53kDa.

Specificity confirmed
The specificity of Anti-TFEB antibody [EPR22941-6] (ab267351) has been confirmed by Western blot testing in TFEB Knockout HepG2 cell line, ab277871.

Other related products
We have a range of other formats of antibody clone [EPR22941-6] also available for your convenience: ab270604, Carrier free - ab270614, Alexa Fluor® 488 - ab282162, PE - ab282787, Alexa Fluor® 647 - ab283274, APC - ab283612, Alexa Fluor® 555 - ab320288, Alexa Fluor® 594 - ab320289, Alexa Fluor® 750 - ab321534, Carrier free - ab267342

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


15 product images



Flow Cytometry (Intracellular) - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Flow Cytometry (Intracellular) staining using rabbit Anti-TFEB antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized IM-9 (human multiple myeloma B Lymphoblast) cells labelling TFEB with ab267351 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TFEB antibody

Immunohistochemical analysis of paraffin-embedded Human lung cancer tissue labeling TFEB with ab267351 at 1/500 dilution (1.042ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining in human lung cancer (PMID: 26264650) The section was incubated with ab256492 for 15 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-TFEB antibody


Immunofluorescent analysis of 100% Methanol-fixed IM-9 (human multiple myeloma B Lymphoblast) cells labelling TFEB with ab267351 at 1/50 dilution, followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in IM-9 cells. 100% methanol fixation is recommended. ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TFEB antibody [EPR22941-6] (ab267351)

Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling TFEB with ab267351 at 1/500 dilution (1.042ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining human colon (PMID:30519051). The section was incubated with ab256492 for 15 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-TFEB antibody

Immunofluorescent analysis of 100% Methanol-fixed Raji (human Burkitt's lymphoma B lymphocyte) cells labelling TFEB with ab267351 at 1/50 dilution, followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in Raji cells. 100% methanol fixation is recommended. ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TFEB antibody

Immunohistochemical analysis of paraffin-embedded Human breast cancer tissue labeling TFEB with ab267351 at 1/500 dilution (1.042ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining in human breast cancer (PMID:28017540). The section was incubated with ab256492 for 15 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND®RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.



Immunoprecipitation - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB was immunoprecipitated from 0.35 mg Raji (human Burkitt's lymphoma B lymphocyte) whole cell lysate with ab267351 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab267351 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: Raji (human Burkitt's lymphoma B lymphocyte) whole cell lysate 10ug

Lane 2: ab267351 IP in Raji whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab267351 in Raji whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 15 seconds.

The weak band around 50kDa is one isoform of TFEB(PMID:28993453).

All lanes:
Immunoprecipitation - Anti-TFEB antibody [EPR22941-6] (ab267351)

Predicted band size: 53 kDa

Observed band size: 66 kDa



Immunoprecipitation - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB was immunoprecipitated from 0.35 mg IM-9 (human multiple myeloma B Lymphoblast) whole cell lysate with ab267351 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab267351 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: IM-9 (human multiple myeloma B Lymphoblast) whole cell lysate 10ug

Lane 2: ab267351 IP in IM-9 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab267351 in IM-9 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 2 min.

The weak band around 50kDa is one isoform of TFEB(PMID:28993453).

All lanes:
Immunoprecipitation - Anti-TFEB antibody [EPR22941-6] (ab267351)

Predicted band size: 53 kDa

Observed band size: 66 kDa



Western blot - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Western blot staining using rabbit Anti-TFEB antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression of TFEB (phospho S211) is downregulated in response to Torin 1 treatment (PMID: 28055300).

In Western blot, anti--TFEB antibody (ab267351) staining at 1/1000 dilution. Anti-GAPDH antibody (ab181602) loading control staining at 1/200000 dilution.

All lanes:
Western blot - Anti-TFEB (phospho S211) antibody [EPR27960-64] (ab310331) at 1/1000 dilution

Lane 1:
Untreated Raji (human Burkitts lymphoma B lymphocyte) whole cell lysate at 20 µg


Lane 2:
Raji treated with 250 nM Torin 1 for 5 hours whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 70 kDa

Exposure time: 81s



Western blot - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Western blot staining using rabbit Anti-TFEB antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Low expression: MCF7.

The lysates were freshly made and used for Western blotting immediately to minimize protein degradation.

In Western blot, anti-TFEB antibody (ab267351) staining at 1/1000 dilution. Anti-GAPDH antibody (ab181602) loading control staining at 1/200000 dilution.

All lanes:
Western blot - Anti-TFEB (phospho S211) antibody [EPR27960-64] (ab310331) at 1/1000 dilution

Lane 1:
Raji (human Burkitts lymphoma B lymphocyte) whole cell lysate (untreated membrane) at 20 µg

Lane 2:
MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate (untreated membrane) at 20 µg

Lane 3:
Raji (human Burkitts lymphoma B lymphocyte) whole cell lysate (phosphatase treated membrane) at 20 µg


Lane 4:
MCF7 (human breast adenocarcinoma epithelial cell) whole cell lysate (phosphatase treated membrane) at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 70 kDa

Exposure time: 81s



Western blot - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Western blot staining using rabbit Anti-TFEB antibody

Western blot: Anti-TFEB antibody [EPR22941-6] ab267351 staining at 1/1000 dilution, shown in green; Mouse anti GAPDH ab8245 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 60 kDa in Wild-type Raji cell lysates with no signal observed at this size in TFEB knockout Raji cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3% Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-TFEB antibody [EPR22941-6] (ab267351) at 1/1000 dilution

Lane 1:
Wild-type Raji cell lysates at 20 µg

Lane 2:

Western blot - Human TFEB knockout Raji cell line (ab275832) at 20 µg

Lanes 3 and 5:
EMPTY

Lane 4:
Human wild-type Hep G2 cell line at 40 µg

Lane 6:
Western blot - Human TFEB knockout Hep G2 cell line (ab277871) at 40 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 60 kDa

Observed band size: 60 kDa



Western blot - Anti-TFEB antibody [EPR22941-6] (ab267351)

TFEB Western blot staining using rabbit Anti-TFEB antibody

False colour image of Western blot: Anti-TFEB antibody [EPR22941-6] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab267351 was shown to bind specifically to TFEB. A band was observed at 60 kDa in wild-type HepG2 cell lysates with no signal observed at this size in TFEB knockout cell line ab277871 (knockout cell lysate ab284220). To generate this image, wild-type and TFEB knockout HepG2 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.
All lanes:
Western blot - Anti-TFEB antibody [EPR22941-6] (ab267351) at 1/1000 dilution

Lane 1:
Wild-type HepG2 cell lysate at 20 µg

Lane 2:
TFEB knockout HepG2 cell lysate at 20 µg

Lane 2:
Western blot - Human TFEB knockout Hep G2 cell line (ab277871)


Lane 3:
Raji cell lysate at 20 µg

Lane 4:
MCF7 cell lysate at 20 µg

Performed under reducing conditions.


Predicted band size: 53 kDa

Observed band size: 60 kDa




ChIC/CUT&RUN sequencing - Anti-TFEB antibody [EPR22941-6] (ab267351)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 Raji (human Burkitt's lymphoma B lymphocyte) cells and 5 µg of ab267351 [EPR22941-6]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.
The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-TFEB antibody [EPR22941-6] (ab267351)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 Raji (human Burkitt's lymphoma B lymphocyte) cells and 5 µg of ab267351 [EPR22941-6]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.
The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-TFEB antibody [EPR22941-6] (ab267351)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2.5 x 10^5 Raji (human Burkitt's lymphoma B lymphocyte) cells and 5 µg of ab267351 [EPR22941-6]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control ab172730 is also shown.
The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.