abcam

ab268021

Anti-Fibronectin antibody [EPR23110-25]

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23110-25
Purification technique	Affinity purification Protein A
Concentration	0.469 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Expected

Species	Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

ICC/IF

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Species	Rat
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information FN1

Function	Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin (PubMed:3024962, PubMed:3593230, PubMed:3900070, PubMed:7989369). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape (PubMed:3024962, PubMed:3593230, PubMed:3900070, PubMed:7989369). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4 receptor, inhibiting FCGR1A/CD64-mediated monocyte activation (PubMed:34089617). Anastellin. Binds fibronectin and induces fibril formation. This fibronectin polymer, named superfibronectin, exhibits enhanced adhesive properties. Both anastellin and superfibronectin inhibit tumor growth, angiogenesis and metastasis. Anastellin activates p38 MAPK and inhibits lysophospholipid signaling. Secreted by contracting muscle, induces liver autophagy, a degradative pathway for nutrient mobilization and damage removal, and systemic insulin sensitization via hepatic ITGA5:ITGB1 integrin receptor signaling.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.


Full details and terms and conditions can be found here:
Terms & Conditions.

10 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Immunohistochemical analysis of paraffin-embedded human pancreatic carcinoma labeling Fibronectin with ab268021 at 1/2000 dilution (0.228 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP).The section was incubated with ab268021 at 4°C overnight. Positive staining in the human pancreatic carcinoma noted. Counterstained with hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP). Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Fibronectin antibody

Immunohistochemical analysis of paraffin-embedded human mammary gland tissue labeling Fibronectin with ab268021 at 1/2000 dilution (0.23 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the stroma of human mammary gland (PMID:650101). Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Fibronectin antibody

Immunohistochemical analysis of paraffin-embedded human esophageal squamous cell carcinoma tissue labeling Fibronectin with ab268021 at 1/2000 dilution (0.23 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the stroma of human esophageal squamous cell carcinoma (PMID:30314454). Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Fibronectin antibody

Immunohistochemical analysis of paraffin-embedded rat liver tissue labeling Fibronectin with ab268021 at 1/2000 dilution (0.23 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the perivascular tissue of sinusoids of rat liver (PMID:650101). Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Frozen sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Immunohistochemistry (Frozen sections) staining using rabbit Anti-Fibronectin antibody

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized rat colon tissue labeling Fibronectin with ab268021 at 1/100 dilution followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Positive staining mainly on smooth muscle of rat colon (PMID: 17881565). The nuclear counterstain is DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Frozen sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Immunohistochemistry (Frozen sections) staining using rabbit Anti-Fibronectin antibody

Immunohistochemical analysis of frozen section of 4% PFA-fixed, 0.2% Triton X-100 permeabilized mouse colon tissue labeling Fibronectin with ab268021 at 1/100 dilution followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Positive staining mainly on smooth muscle of mouse colon (PMID: 17881565). The nuclear counterstain is DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Fibronectin antibody

Immunohistochemical analysis of paraffin-embedded mouse stomach tissue labeling Fibronectin with ab268021 at 1/2000 dilution (0.23 µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Positive staining in the smooth muscle of mouse stomach (PMID:650101). Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).


Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Flow Cytometry - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Flow Cytometry staining using rabbit Anti-Fibronectin antibody

Flow cytometric analysis of NIH/3T3 (mouse embryonic fibroblast) cells labeling Fibronectin with ab268021 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Gated on viable cells.



Western blot - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Western blot staining using rabbit Anti-Fibronectin antibody

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 44 seconds.

The expression molecular weight observed is consistent with what has been described in the literature (PMID:28367057).

ab268021 was shown to specifically react with Fibronectin in wild-type HAP1 cells as signal was lost in Fibronectin knockout cells. Wild-type and Fibronectin knockout samples were subjected to SDS-PAGE. ab268021 and ab179475 (Rabbit anti-Integrin alpha V loading control) were incubated 1 hour at room temperature at 1/1000 dilution and 1/5000 dilution respectively. Blots were developed with Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) secondary antibody at 1/100,000 dilution for 1 hour at room temperature before imaging. The blot was developed on a BIO-RAD® ChemiDoc™ MP instrument using the ECL technique.

All lanes:
Western blot - Anti-Fibronectin antibody [EPR23110-25] (ab268021) at 1/1000 dilution

Lane 1:
Human colon lysate at 20 µg

Lane 2:
Human stomach lysate at 20 µg

Lane 3:
Wild-type HAP1 whole cell lysate at 40 µg

Lane 4:
Fibronectin knockout HAP1 whole cell lysate at 40 µg


Secondary

Lanes 1 - 2:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/1000 dilution

Lanes 3 - 4:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 262 kDa

Observed band size: 285 kDa



Western blot - Anti-Fibronectin antibody [EPR23110-25] (ab268021)

Fibronectin Western blot staining using rabbit Anti-Fibronectin antibody

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 3.25 seconds.

The expression molecular weight observed is consistent with what has been described in the literature (PMID:28367057).

All lanes:
Western blot - Anti-Fibronectin antibody [EPR23110-25] (ab268021) at 1/1000 dilution

Lane 1:
Human serum at 20 µg

Lane 2:
Human plasma at 20 µg

Lane 3:
Mouse plasma at 20 µg

Lane 4:
Rat plasma at 20 µg

Secondary

Lanes 1 - 2:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/1000 dilution

Lanes 3 - 4:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 262 kDa

Observed band size: 285 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.