abcam

ab268270

Anti-TNFSF5 antibody [CD40LG/2763] - BSA and Azide free

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Recombinant Fragment Protein within Human CD40LG aa 100 to C-terminus. The exact immunogen used to generate this antibody is proprietary information. Database link P29965
Carrier free	Yes
Clone number	CD40LG/2763
Purification technique	Affinity purification Protein A/G
Light chain type	kappa
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from bioreactor concentrate.

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	1-2 µg for 10^6 Cells
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Protein Array

Tested

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1-2 µg/mL
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-

Notes	-

IHC-P

Tested

Species	Human
Dilution info	1-2 µg/mL
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Target data

See full target information CD40LG

Function	Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:1280226, PubMed:31331973). Costimulates T-cell proliferation and cytokine production (PubMed:8617933). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:8617933). Induces the activation of NF-kappa-B (PubMed:15067037, PubMed:31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity). CD40 ligand, soluble form. Acts as a ligand for integrins, specifically ITGA5:ITGB1 and ITGAV:ITGB3; both integrins and the CD40 receptor are required for activation of CD40-CD40LG signaling, which have cell-type dependent effects, such as B-cell activation, NF-kappa-B signaling and anti-apoptotic signaling.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

5 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TNFSF5 antibody [CD40LG/2763] - BSA and Azide free (ab268270)

TNFSF5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using mouse Anti-TNFSF5 antibody

Formalin-fixed, paraffin-embedded human spleen tissue stained for TNFSF5 using ab268146 at 2 µg/ml in immunohistochemical analysis.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and Sodium azide (ab268146)



Flow Cytometry - Anti-TNFSF5 antibody [CD40LG/2763] - BSA and Azide free (ab268270)

TNFSF5 Flow Cytometry staining using mouse Anti-TNFSF5 antibody


Flow cytometric analysis of Jurkat (Human T cell leukemia cell line from peripheral blood) cells labeling TNFSF5 using ab268146 at 2 µg/million cells followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and Sodium azide (ab268146)



Immunocytochemistry/ Immunofluorescence - Anti-TNFSF5 antibody [CD40LG/2763] - BSA and Azide free (ab268270)

TNFSF5 Immunocytochemistry/ Immunofluorescence staining using mouse Anti-TNFSF5 antibody


Immunofluorescence staining of Jurkat (Human T cell leukemia cell line from peripheral blood) cells stained for TNFSF5 with ab268146 at 2 µg/ml followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and Sodium azide (ab268146)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TNFSF5 antibody [CD40LG/2763] - BSA and Azide free (ab268270)

TNFSF5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using mouse Anti-TNFSF5 antibody

Formalin-fixed, paraffin-embedded human spleen tissue stained for TNFSF5 using ab268146 at 2 µg/ml in immunohistochemical analysis.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and Sodium azide (ab268146)



Protein Array - Anti-TNFSF5 antibody [CD40LG/2763] - BSA and Azide free (ab268270)

Protein Array containing more than 19,000 full-length human proteins using ab268146.
Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an

S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and Sodium azide (ab268146)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.