abcam

ab269948

Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23244-12
Purification technique	Affinity purification Protein A
Concentration	0.555 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

IHC-P

Not recommended

Species	Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Target data

See full target information F11R

Function Seems to play a role in epithelial tight junction formation. Appears early in primordial forms of cell junctions and recruits PARD3 (PubMed:11489913). The association of the PARD6-PARD3 complex may prevent the interaction of PARD3

with JAM1, thereby preventing tight junction assembly (By similarity). Plays a role in regulating monocyte transmigration involved in integrity of epithelial barrier (By similarity). Ligand for integrin alpha-L/beta-2 involved in memory T-cell and neutrophil transmigration (PubMed:11812992). Involved in platelet activation (PubMed:10753840). (Microbial infection) Acts as a receptor for Mammalian reovirus sigma-1. (Microbial infection) Acts as a receptor for Human Rotavirus strain Wa.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Junctional Adhesion Molecule 1 also known as JAM-A or JAM-A protein is a member of the immunoglobulin superfamily. It has a mass of roughly 32 kDa. This molecule functions in tight junctions of epithelial and endothelial cells playing an important

	role in regulating paracellular permeability. JAM-A is expressed in tissues where tight junctions are present including the skin gastrointestinal tract and endothelia throughout the body.
Biological function summary	JAM-A functions as a cell adhesion molecule influencing processes like leukocyte transmigration and epithelial barrier formation. It does not act alone but rather forms complexes with other proteins at cell junctions. JAM-A interacts with other junctional proteins like occludin and claudins contributing to the structure and function of tight junctions.
Pathways	JAM-A participates in cellular processes involving inflammation and cell migration. It is an integral part of the signal transduction pathways that regulate leukocyte migration an important aspect of immune response. JAM-A is also associated with G-protein coupled receptors in these pathways highlighting its role in regulating intracellular signaling that affects cell adhesion and migration.
Associated diseases and disorders	JAM-A has connections with cancer progression and cardiovascular diseases. Its overexpression has been linked to several cancers including breast cancer due to its role in enhancing cell migration and invasion. In cardiovascular diseases JAM-A interacts with platelet-endothelial cell adhesion molecule-1 (PECAM-1) influencing the inflammatory response within vascular tissues. Understanding these interactions can aid in developing therapeutic strategies targeting JAM-A in related diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Flow Cytometry - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Junctional Adhesion Molecule 1/JAM-A Flow Cytometry staining using rabbit Anti-Junctional Adhesion Molecule 1/JAM-A antibody

Flow cytometric analysis of HT-29 (Human colorectal adenocarcinoma epithelial cell) cells labeling Junctional Adhesion Molecule 1/JAM-A with ab269948 at 1/500 compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 was used as the secondary antibody.
Gated on viable cells.




Flow Cytometry - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Junctional Adhesion Molecule 1/JAM-A Flow Cytometry staining using rabbit Anti-Junctional Adhesion Molecule 1/JAM-A antibody

Flow cytometric analysis of human B lymphocytes (Left) / human monocytes (Right) cells labeling Junctional Adhesion Molecule 1/JAM-A with ab269948 at 1/500 compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 was used as the secondary antibody.
Human peripheral blood mononuclear cell (PBMC) co-stained with anti-CD19 conjugated to PE-Cy7 and anti-CD14 conjugated to BV510. JAM-A expression on B lymphocytes (CD19+) and monocytes (CD14+) population are shown respectively.


Gated on viable cells.



Immunocytochemistry/ Immunofluorescence - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Junctional Adhesion Molecule 1/JAM-A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Junctional Adhesion Molecule 1/JAM-A antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HT-29 (human colorectal adenocarcinoma epithelial cell) cells labeling Junctional Adhesion Molecule 1/JAM-A with ab269948 at 1/50 dilution, followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing membranous staining in HT-29 cell line is observed. ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).
Secondary antibody only control: ab150077 AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.



Immunoprecipitation - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Junctional Adhesion Molecule 1/JAM-A was immunoprecipitated from 0.35 mg HT-29 (Human colorectal adenocarcinoma epithelial cell) whole cell lysate 10µg with ab269948 at 1/30 dilution (2µg in 0.35 mg lysates). Western blot was performed on the immunoprecipitate using ab269948 at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/1000 dilution.
Lane 1: HT-29 whole cell lysate 10µg.

Lane 2: ab269948 IP in HT-29 whole cell lysate.


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab269948 in HT-29 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds.

All lanes:
Immunoprecipitation - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Predicted band size: 32 kDa



Immunoprecipitation - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Junctional Adhesion Molecule 1/JAM-A was immunoprecipitated from 0.35 mg human lung lysate 10µg with ab269948 at 1/30 dilution (2µg in 0.35 mg lysates). Western blot was performed on the immunoprecipitate using ab269948 at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/1000 dilution.
Lane 1: Human lung lysate 10µg.

Lane 2: ab269948 IP in human lung lysate.


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab269948 in human lung lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds.

All lanes:
Immunoprecipitation - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Predicted band size: 32 kDa



Western blot - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948)

Junctional Adhesion Molecule 1/JAM-A Western blot staining using rabbit Anti-Junctional Adhesion Molecule 1/JAM-A antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression profile/ molecular weight observed is consistent with what has been described in the literature (PMID:10698320, 10753840).

Low expression control: THP-1 and Molt-4 (PMID:10698320).

Exposure time: Lanes 1-5: 15 seconds; Lane 6:70 seconds.
All lanes:
Western blot - Anti-Junctional Adhesion Molecule 1/JAM-A antibody [EPR23244-12] (ab269948) at 1/1000 dilution

Lane 1:
Human lung lysate at 20 µg

Lane 2:
HT-29 (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
THP-1 (human monocytic leukemia monocyte) whole cell lysate at 20 µg

Lane 4:
HUVEC (human umbilical vein endothelial cell) whole cell lysate at 20 µg

Lane 5:
MOLT-4 (human lymphoblastic leukemia t lymphoblast) whole cell lysate at 20 µg

Lane 6:
TF-1 (human erythroleukemia erythroblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 32 kDa

Observed band size: 35 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.