abcam

ab271057

Anti-MAGEA10 antibody [EPR24366-89]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24366-89
Purification technique	Affinity purification Protein A
Concentration	0.519 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/2000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information MAGEA10

Function	Not known, though may play a role in embryonal development and tumor transformation or aspects of tumor progression.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	MAGEA10 also known as Melanoma Antigen Family A 10 is a member of the MAGE (Melanoma Antigen Gene) family. This protein has a molecular mass of about 36 kDa. MAGEA10 is expressed in various tumor cells and some male germline cells in the testis but not in normal somatic tissues. The expression pattern makes it an interesting target for cancer research and potential therapeutic interventions.
Biological function summary	The function of MAGEA10 remains not fully understood but it likely plays a role in cell proliferation and survival. It is not a component of any known protein complex. Research suggests that MAGEA10 might modulate cellular processes by interacting with other proteins impacting pathways related to oncogenesis and cell cycle regulation.
Pathways	MAGEA10 appears to be involved in critical pathways associated with cancer progression and possibly immune evasion. It interacts with proteins involved in cell signaling pathways such as the p53 pathway which is pivotal in regulating the cell cycle and apoptosis. MAGEA10 may influence these pathways by affecting gene transcription and modulating protein-protein interactions.
Associated diseases and disorders	MAGEA10 has a strong connection to cancer particularly in melanoma and lung cancer. Overexpression of MAGEA10 in these cancers suggests a possible role in

tumor growth and metastasis. The protein is part of cancer-testis antigens similar to MAGEA1 another member of the same family implicated in various malignancies suggesting that targeting MAGEA10 could aid in developing specific cancer immunotherapies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MAGEA10 antibody

Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue labelling MAGEA10 with ab271057 at 1/2000 dilution (0.26 ug/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). Nuclear staining in human lung carcinoma. The section was incubated with ab271057 at 4°C overnight. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MAGEA10 antibody

Immunohistochemical analysis of paraffin-embedded human placenta tissue labelling MAGEA10 with ab271057 at 1/2000 dilution (0.26 ug/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). Nuclear staining in human placenta. The section was incubated with ab271057 at 4°C overnight. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer).


Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0)



Flow Cytometry (Intracellular) - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Flow Cytometry (Intracellular) staining using rabbit Anti-MAGEA10 antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized SK-MEL-2 (human skin malignant melanoma) cells labelling MAGEA10 with ab271057 at 1/500 dilution (0.1ug)/ (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunocytochemistry/ Immunofluorescence - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MAGEA10 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized A375 (human malignant melanoma epithelial cell) cells labelling MAGEA10 with ab271057 at 1/500 dilution (1.038 ug/ml), followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)

preadsorbed antibody at 1/1000 dilution (Green). Confocal image showing nuclear staining in A375 cell line. **Negative control**: 293T (PMID: 10446460) is observed.
ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MAGEA10 antibody

Immunohistochemical analysis of paraffin-embedded human liver tissue labelling MAGEA10 with ab271057 at 1/2000 dilution (0.26 ug/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). **Negative control**: no staining in human liver. The section was incubated with ab271057 at 4°C overnight. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MAGEA10 antibody

Immunohistochemical analysis of paraffin-embedded human kidney tissue labelling MAGEA10 with ab271057 at 1/2000 dilution (0.26 ug/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). **Negative control**: no staining in human kidney. The section was incubated with ab271057 at 4°C overnight. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP polymer).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0)




Immunocytochemistry/ Immunofluorescence - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MAGEA10 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized SK-MEL-2 (human skin malignant melanoma) cells labelling MAGEA10 with ab271057 at 1/500 dilution (1.038 ug/ml), followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green). Confocal image showing nuclear staining in SK-MEL-2 cell line. **Negative control**:293T (PMID: 10446460) is observed.
ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Flow Cytometry (Intracellular) staining using rabbit Anti-MAGEA10 antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized 293T (Human embryonic kidney epithelial cell, Left)/ A375 (Human malignant melanoma epithelial cell, Right) cells labelling MAGEA10 with ab271057 at 1/500 dilution (0.1ug)/ (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. **Negative control**: 293T.(PMID:10446460)



Immunoprecipitation - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 was immunoprecipitated from 0.35 mg A375 (human malignant melanoma epithelial cell) whole cell lysate with ab271057 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab271057 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: A375 (human malignant melanoma epithelial cell) whole cell lysate 10 ug

Lane 2: ab271057 IP in A375 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab271057 in A375 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 7.75 seconds

All lanes:
Immunoprecipitation - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

Predicted band size: 41 kDa

Observed band size: 65 kDa



Immunoprecipitation - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 was immunoprecipitated from 0.35 mg SK-MEL-2 (human skin malignant melanoma) whole cell lysate with ab271057 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab271057 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: SK-MEL-2 (human skin malignant melanoma) whole cell lysate 10 ug

Lane 2: ab271057 IP in SK-MEL-2 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab271057 in SK-MEL-2 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 67 seconds

All lanes:
Immunoprecipitation - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

Predicted band size: 41 kDa

Observed band size: 65 kDa



Western blot - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Western blot staining using rabbit Anti-MAGEA10 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Negative control: 293T (PMID:10446460)

Exposure time: Lane 1: 3.75 seconds Lane 2-4: 36 seconds

All lanes:
Western blot - Anti-MAGEA10 antibody [EPR24366-89] (ab271057) at 1/1000 dilution

Lane 1:
SK-MEL-2 (human skin malignant melanoma), whole cell lysate at 20 µg

Lane 2:
A375 (human malignant melanoma epithelial cell), whole cell lysate at 20 µg

Lane 3:
SH-SY5Y (human neuroblastoma epithelial cell), whole cell lysate at 20 µg


Lane 4:
293T (human embryonic kidney epithelial cell), whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 41 kDa

Observed band size: 65 kDa



Western blot - Anti-MAGEA10 antibody [EPR24366-89] (ab271057)

MAGEA10 Western blot staining using rabbit Anti-MAGEA10 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Negative control: Colon, liver (PMID: 10446460).

This blot was developed using a higher sensitivity ECL substrate.

Exposure time: 3 minutes

All lanes:
Western blot - Anti-MAGEA10 antibody [EPR24366-89] (ab271057) at 1/1000 dilution

Lane 1:
Human placenta tissue lysate at 20 µg

Lane 2:
Human colon tissue lysate at 20 µg

Lane 3:
Human liver tissue lysate at 20 µg

Secondary

All lanes:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/1000 dilution

Predicted band size: 41 kDa

Observed band size: 65 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.