abcam

ab271092

Anti-CD133 antibody [EPR24348-38]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24348-38
Purification technique	Affinity purification Protein A
Concentration	0.473 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Rat, Human, Mouse
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Not recommended

Species	Rat, Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Not recommended

Species	Rat, Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Not recommended

Species	Rat, Human
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	1/100
Notes	-

Not recommended

Species	Rat, Human
Dilution info	-
Notes	-

Target data

See full target information Prom1

Function	May play a role in cell differentiation, proliferation and apoptosis. Binds cholesterol in cholesterol-containing plasma membrane microdomains and may play a role in the organization of the apical plasma membrane in epithelial cells. During early retinal development acts as a key regulator of disk morphogenesis (PubMed:19228982). Involved in regulation of MAPK and Akt signaling pathways. In neuroblastoma cells suppresses cell differentiation such as neurite outgrowth in a RET-dependent manner.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD133 also known as Prominin-1 is a transmembrane glycoprotein with a molecular weight of approximately 120 kDa. This protein stands out due to its expression in various stem and progenitor cells including those of the neural

	hematopoietic and epithelial lineages. CD133 is recognized for its distinct five transmembrane domains and two large extracellular loops. Researchers commonly use CD133 as a cell surface marker to identify stem and progenitor cells. The significance of CD133 extends to immunohistochemistry (IHC) and Western blot applications where it is a popular target for detection and analysis.
Biological function summary	CD133 is involved in maintaining the cellular architecture and signaling processes. It plays a role in cellular differentiation and proliferation while acting as a part of larger protein complexes on the cell surface. This protein is notable as it contributes significantly to the regulation of stem cell properties supporting the growth and repair in various tissues. Anti-CD133 antibodies are valuable tools in studying these functions due to their specificity in targeting this marker.
Pathways	CD133 participates in cellular signaling and transport pathways vital for intercellular communication and homeostasis. It interacts within pathways related to cell polarity and asymmetric division both essential for tissue development and maintenance. CD133 is associated with proteins like CD133/1 and CD133/2 which further highlights its involvement in these complex biological networks and on pathways such as the Wnt signaling pathway which regulates gene expression associated with cell proliferation and differentiation.
Associated diseases and disorders	Several studies indicate a strong relationship between CD133 expression and certain cancers such as glioblastoma and colorectal cancer. In these cases CD133 serves not only as a marker but also as an indicator of the presence of cancer stem cells which are capable of tumor initiation and maintenance. Furthermore researchers have identified connections between CD133 and other proteins such as c-Met (cMab) within these pathological contexts suggesting interactions that could contribute to disease progression and offering potential targets for therapeutic intervention.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

13 product images




Immunohistochemistry (Frozen sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Frozen sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse E14.5 embryonic kidney tissue labeling CD133 with ab271092 at 1/100 (4.59 ug/ml) dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Positive staining on the apical membrane of mouse E14.5 embryonic kidney is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)at 1/1000 dilution.


Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Flow Cytometry (Intracellular) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Flow Cytometry (Intracellular) staining using rabbit Anti-CD133 antibody

Flow Cytometry analysis of Mouse bone marrow cells labelling CD133 with ab271092 at 1/50 (right). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 was used as the secondary antibody. Rabbit monoclonal IgG (ab172730) was used as the isotype control (left). Cells were co-stained with anti-CD117 conjugated to BV421. Gated on viable cells.




Immunohistochemistry (Frozen sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Frozen sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse E14.5 embryonic lung tissue labeling CD133 with ab271092 at 1/100 (4.59 ug/ml) dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Positive staining on the apical membrane of lung buds is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labelling CD133 with ab271092 at 1/2000 (0.23 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse kidney. The section was incubated with ab271092 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Flow Cytometry (Intracellular) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Flow Cytometry (Intracellular) staining using rabbit Anti-CD133 antibody


Flow Cytometry analysis of Mouse bone marrow cells labelling CD133 with ab271092 at 1/50 (right). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat Anti-Rabbit IgG (Alexa Fluor® 647) (ab150083) at 1/2000 was used as the secondary antibody. Rabbit monoclonal IgG (ab172730) was used as the isotype control (left). Cells were stained with rabbit IgG or ab271092. Then stained with Lineage cocktail conjugated to FITC, anti-Sca-1 conjugated to PE/Cy7, anti-CD117 conjugated to APC/Cy7 and anti-CD16/32 conjugated to BV510. Gated on viable MP cells. Positive staining on CD16/32+ MP cells is consistent with literature. (PMID: 23509298).



Flow Cytometry (Intracellular) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Flow Cytometry (Intracellular) staining using rabbit Anti-CD133 antibody

Flow Cytometry analysis of Mouse bone marrow cells labelling CD133 with ab271092 at 1/50 (right). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 was used as the secondary antibody. Rabbit monoclonal IgG (ab172730) was used as the isotype control (left). Gated on viable cells.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of paraffin-embedded Mouse liver tissue labelling CD133 with ab271092 at 1/2000 (0.23 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on the biliary ducts of mouse liver. The section was incubated with ab271092 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of paraffin-embedded Mouse pancreas tissue labelling CD133 with ab271092 at 1/2000 (0.23 ug/ml) followed by a ready to use LeicaDS9800(Bond™ Polymer Refine Detection). Positive staining on mouse pancreatic ducts. The section was incubated with ab271092 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of paraffin-embedded Mouse retina tissue labelling CD133 with ab271092 at 1/2000 (0.23 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on the cone and rod cells of mouse retina. The section was incubated with ab271092 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Frozen sections) - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Immunohistochemistry (Frozen sections) staining using rabbit Anti-CD133 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse E14.5 embryonic cortex tissue labeling CD133 with ab271092 at 1/100 (4.59 ug/ml) dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Positive staining on the apical membrane of mouse E14.5 embryonic cortex is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Immunoprecipitation - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 was immunoprecipitated from 0.35 mg Mouse E14.5 brain tissue lysate 10 ug with ab271092 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab271092 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Mouse E14.5 brain tissue lysate 10 ug

Lane 2: ab271092 IP in Mouse E14.5 brain tissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab271092 in Mouse E14.5 brain tissue lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds

All lanes:
Immunoprecipitation - Anti-CD133 antibody [EPR24348-38] (ab271092)

Predicted band size: 97 kDa

Observed band size: 120 kDa



Western blot - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Western blot staining using rabbit Anti-CD133 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Negative control: mouse heart (PMID:17575139).

Exposure time: Lanes 1-2: 10 seconds; Lane 3: 70 seconds.

All lanes:
Western blot - Anti-CD133 antibody [EPR24348-38] (ab271092) at 1/1000 dilution

Lane 1:
Mouse kidney tissue lysate at 20 µg

Lane 2:
Mouse heart tissue lysate at 20 µg


Lane 3:
Mouse E14.5 brain tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 97 kDa

Observed band size: 120 kDa



Western blot - Anti-CD133 antibody [EPR24348-38] (ab271092)

CD133 Western blot staining using rabbit Anti-CD133 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-CD133 antibody [EPR24348-38] (ab271092) at 1/1000 dilution

Lane 1:
Mouse kidney tissue lysate

Lane 2:
Mouse kidney tissue lysate treated with PNGase F

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 97 kDa

Observed band size: 120 kDa, 97 kDa

Exposure time: 10s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.