abcam

ab271123

Anti-TM9SF2 antibody [EPR24416-17]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24416-17
Purification technique	Affinity purification Protein A
Concentration	0.593 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)

Species	Rat
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat
Dilution info	1/2000
Notes	-

Species	Human
Dilution info	1/2000
Notes	-

Target data

See full target information TM9SF2

Function	In the intracellular compartments, may function as a channel or small molecule transporter.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	TM9SF2 also known as Transmembrane 9 Superfamily Member 2 is a membrane protein with a mass of approximately 80 kDa. It functions as a transmembrane transporter that may influence ion homeostasis. TM9SF2 is mainly expressed in the endoplasmic reticulum and Golgi apparatus of cells where it likely impacts protein processing and trafficking. Due to its nature as a membrane protein TM9SF2 may also have roles in cellular signaling. Alternate names for TM9SF2 include TMS9 and Nonaspanin.
Biological function summary	TM9SF2 influences cellular processes by contributing to membrane integrity and protein sorting. As a possible member of larger protein complexes within the cell TM9SF2 interacts with other transmembrane proteins to facilitate intracellular trafficking pathways. This protein may modify the sorting and secretion of enzymes or hormones through its action on vesicular transport mechanisms therefore affecting various metabolic and signaling processes within the cell.
Pathways	TM9SF2 is involved in membrane trafficking and protein secretion important for maintaining cellular homeostasis. It associates with important biological pathways such as vesicular transport pathways and intracellular sorting pathways. TM9SF2 acts together with proteins like SNAREs and Rab GTPases contributing to vesicle budding fusion and transport. This association helps coordinate how proteins are delivered to different cellular destinations integrating TM9SF2's function within the broader picture of cellular operations.
Associated diseases and disorders	TM9SF2 has connections with metabolic diseases and neurological disorders. Alterations in TM9SF2 expression have been linked with abnormal lipid metabolism and could play a role in the pathogenesis of obesity and type 2 diabetes. Additionally irregularities in its function might influence neurodegenerative conditions by affecting protein trafficking and degradation pathways. TM9SF2 interacts with proteins such as APP (Amyloid precursor protein) suggesting potential involvement in Alzheimer's disease where protein misprocessing leads to amyloid plaque formation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


9 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labelling TM9SF2 with ab271123 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human kidney. The section was incubated with ab271123 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Flow Cytometry (Intracellular) - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

TM9SF2 Flow Cytometry (Intracellular) staining using rabbit Anti-TM9SF2 antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized U-87 MG (Human glioblastoma-astrocytoma epithelial cell) cells labelling with ab271123 at 1/50 (Red) compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

TM9SF2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TM9SF2 antibody

Immunohistochemical analysis of paraffin-embedded Rat pancreas tissue labelling TM9SF2 with ab271123 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on rat pancreas. The section was incubated with ab271123for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

TM9SF2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-TM9SF2 antibody

Immunohistochemical analysis of paraffin-embedded Mouse pancreas tissue labelling TM9SF2 with ab271123 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse pancreas. The section was incubated with ab271123 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Frozen sections) - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

TM9SF2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-TM9SF2 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse kidney tissue labelling TM9SF2 with ab271123 at 1100 (5.93 µg/ml) dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Cytoplasmic staining on mouse kidney is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Frozen sections) - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

TM9SF2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-TM9SF2 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat kidney tissue labelling TM9SF2 with ab271123 at 1100 (5.93 µg/ml) dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Cytoplasmic staining on rat kidney is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Western blot - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

TM9SF2 Western blot staining using rabbit Anti-TM9SF2 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 3 minutes

All lanes:
Western blot - Anti-TM9SF2 antibody [EPR24416-17] (ab271123) at 1/1000 dilution

Lane 1:
HeLa (human epithelial cell line from cervix adenocarcinoma), whole cell lysate at 40 µg

Lane 2:
HT-29 (human colorectal adenocarcinoma epithelial cell), whole cell lysate at 40 µg

Lane 3:
PANC-1 (human pancreatic epithelioid carcinoma epithelial cell), whole cell lysate at 40 µg

Lane 4:
U-87 MG (human glioblastoma-astrocytoma epithelial cell), whole cell lysate at 40 µg


Lane 5:
Human kidney tissue lysate at 40 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/2000 dilution

Predicted band size: 76 kDa

Observed band size: 55 kDa



Western blot - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Samples are non-boiled as boiling may cause protein aggregates.

Exposure time 3 minutes

All lanes:
Western blot - Anti-TM9SF2 antibody [EPR24416-17] (ab271123) at 1/1000 dilution

Lane 1:
C6 (rat glial tumor glial cell), whole cell lysate at 20 µg

Lane 2:
PC-12 (rat adrenal gland pheochromocytoma), whole cell lysate at 20 µg

Lane 3:
Mouse kidney tissue lysate at 20 µg

Lane 4:
Mouse colon tissue lysate at 20 µg


Lane 5:
Rat colon tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 76 kDa

Observed band size: 55 kDa



Western blot - Anti-TM9SF2 antibody [EPR24416-17] (ab271123)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Band higher than 250 kDa should be TM9SF2 aggregates.

Exposure time: 10 seconds

All lanes:
Western blot - Anti-TM9SF2 antibody [EPR24416-17] (ab271123) at 1/1000 dilution

Lane 1:
HEK-293T (human embryonic kidney) transfected with an empty vector, containing a myc-His-tag®, whole cell lysate at 10 µg

Lane 2:
HEK-293T transfected with human TM9SF2 expression vector containing a myc-His-tag®, whole cell lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 76 kDa

Observed band size: 55 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.