abcam

ab271180

Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118]

Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180) is a rabbit monoclonal antibody detecting SARS-CoV-2 nucleocapsid protein in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF, ELISA**. Suitable for **SARS-CoV-2**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24334-118
Purification technique	Affinity purification Protein A
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Recombinant fragment - SARS-CoV-2
Dilution info	1/1000
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

IP

Tested

Species	Recombinant fragment - SARS-CoV-2
Dilution info	1/30
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

ELISA

Tested

Species	Recombinant fragment - SARS-CoV-2
Dilution info	-
Notes	Use at 31.25 ng /ml

Predicted

Species	SARS-CoV-2
Dilution info	-

Notes	-

WB

Tested

Species	Recombinant fragment - SARS-CoV-2
Dilution info	1/1000
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Recombinant fragment - SARS-CoV-2
Dilution info	1/500
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

IHC-P

Tested

Species	Recombinant fragment - SARS-CoV-2
Dilution info	1/4000
Notes	

Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Target data

See full target information SARS-CoV-2 nucleocapsid protein

Function

Packages the positive strand viral genome RNA into a helical ribonucleocapsid (RNP) and plays a fundamental role during virion assembly through its interactions with the viral genome and membrane protein M (PubMed:33264373). Plays an important role in enhancing the efficiency of subgenomic viral RNA transcription as well as viral replication. Attenuates the stress granules formation by reducing host G3BP1 access to host mRNAs under stress conditions (PubMed:34901782, PubMed:36534661). May block host chemokine function in vivo, facilitating viral replication and transmission (PubMed:35921414). Acts by being secreted into the extracellular space where it competes to host chemokines for binding to host glycosaminoglycans (GAG) (PubMed:35921414). May induce inflammasome responses in cultured cells and mice. Acts by interacting with host NLRP3 to facilitate inflammasome assembly, which induces cytokine release that may play a role in COVID lung injury. Disrupts the association betwen NLRP3 and ABHD8, enhancing NLRP3 stability and leading to increased inflammasome activation (PubMed:39225180).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF), ELISA in SARS-CoV-2 samples.

Other related products
We have a range of other formats of antibody clone [EPR24334-118] also available for your convenience: ab271180, Carrier free - ab280201, Alexa Fluor® 488 - ab283243, PE - ab283244, Alexa Fluor® 647 - ab302552

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The SARS-CoV-2 nucleocapsid protein also known as N protein plays a central role in the packaging and replication of the viral RNA genome. It functions mechanically by binding to the viral RNA to facilitate its replication and encapsidation. The nucleocapsid protein is expressed within the virus-infected cells particularly in the cytoplasm. It is a highly conserved structural protein of the virus and has a molecular weight of approximately 50 kDa. Additionally the nucleocapsid protein aids in RNA binding and is essential for the virus's structural stability.
Biological function summary	The nucleocapsid protein of SARS-CoV-2 is involved in maintaining the integrity of the viral genome. It forms a ribonucleoprotein complex with the viral RNA which is necessary for viral replication. The nucleocapsid protein is integral in the formation of the virus's structure and assists in the assembly and release of new virus particles. It interacts with other viral proteins contributing to the stability and functioning of the virus.
Pathways	The nucleocapsid protein is involved in the viral life cycle and replication pathways. The protein participates in the RNA replication and transcription complex working

	alongside proteins such as the RNA-dependent RNA polymerase and other non-structural proteins. These interactions within the pathways are essential for the efficient replication of the viral genome and production of new viral particles which are key steps in the virus's lifecycle.
Associated diseases and disorders	The nucleocapsid protein is directly linked to COVID-19 the disease caused by the SARS-CoV-2 virus. It is a target for various diagnostic tests such as the COVID ELISA test and corona IgA test due to its immunogenic properties. The presence of nucleocapsid antibodies in patients is often measured through tests like the ELISA test which helps in diagnosing COVID-19. The nucleocapsid protein's involvement in COVID-19 progression connects it with other viral proteins highlighting its significance in disease mechanisms and potential therapeutic targeting.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


15 product images



Flow Cytometry (Intracellular) - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Flow Cytometry (Intracellular) staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HEK-293T (human embryonic kidney) transfected with myc-tagged SARS-CoV-1 Nucleocapsid protein expression construct (Left)/ HEK-293T (human embryonic kidney) transfected with myc-tagged MERS Nucleocapsid protein expression construct (Right).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077 at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Immunohistochemical analysis of paraffin-embedded Human lung labeling SARS-CoV-2 with ab271180 at 1/4000 dilution followed by ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). **Negative control**: No staining on human lung. The section was incubated with ab271180 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Goat Anti-Rabbit IgG H&L (HRP polymer).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100-permeabilized 293T cells transfected with myc-tagged SARS-CoV-1 and MERS Nucleocapsid protein labelling SARS-CoV-2 with ab271180 at 1/1000 dilution, followed by ab150080 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) antibody at 1/1000 dilution. ab202008 Anti-Myc tag mouse monoclonal antibody (Alexa Fluor® 488) was used to counterstain Myc tag at 1/200 dilution (Green). The nuclear counterstain was DAPI (Blue). Confocal image showing no staining in 293T cells transfected with myc-tagged SARS-CoV-1 and MERS Nucleocapsid protein, which are **negative controls** for SARS-CoV-2.



Immunocytochemistry/ Immunofluorescence - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100-permeabilized 293T cells transfected with myc-tagged SARS-CoV-2 Nucleocapsid protein expression vector labelling SARS-CoV-2 with ab271180 at 1/1000 dilution, followed by ab150080 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) antibody at 1/1000 dilution (Red). ab202008 Anti-Myc tag mouse monoclonal antibody (Alexa Fluor® 488) was used to counterstain Myc tag at 1/200 dilution (Green). The nuclear counterstain was DAPI (Blue). Confocal image showing cytoplasmic staining in 293T cells transfected with myc-tagged SARS-CoV-2 Nucleocapsid protein expression vector.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody


Immunohistochemical analysis of paraffin-embedded HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) transfected with SARS-CoV-2 Nucleocapsid protein expression vector containing a myc-His-tag® (Panel A)
HEK-293T transfected with SARS-CoV-1 Nucleocapsid protein expression vector containing a myc-His-tag® (Panel B) HEK-293T transfected with MERS Nucleocapsid protein expression vector containing a myc-His-tag® (Panel C) HEK-293T transfected with control vector containing a myc-His-tag® (Panel D) with 271180 at 1/4000 dilution followed by ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). Positive staining on (A) HEK-293T transfected with a SARS-CoV-2 Nucleocapsid protein expression vector is observed. No staining on (B) HEK-293T transfected with a SARS-CoV-1 Nucleocapsid protein expression vector, (C) HEK-293T transfected with MERS Nucleocapsid protein expression vector and (D) HEK-293T transfected with empty vector. The section was incubated with ab271180 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Flow Cytometry (Intracellular) - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Flow Cytometry (Intracellular) staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HEK-293T (human embryonic kidney) transfected with myc-tagged SARS-CoV-2 Nucleocapsid protein expression construct labelling SARS-CoV-2 with ab271180 at 1/500 dilution (Right) compared with a Rabbit monoclonal IgG (ab172730) (Left) isotype control.
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077 at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 was immunoprecipitated from 0.35 mg HEK-293T (human embryonic kidney) transfected with SARS-CoV-2 Nucleocapsid protein expression vector containing a myc-His-tag®, whole cell lysate 5 µg with ab271180 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab271180 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HEK-293T (human embryonic kidney) transfected with SARS-CoV-2 Nucleocapsid protein expression vector containing a myc-His-tag® whole cell lysate 5 µg


Lane 2: ab271180 IP in HEK-293T transfected with SARS-CoV-2 Nucleocapsid protein expression vector containing a myc-His-tag® whole cell lysate.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab271180 in HEK-293T transfected with SARS-CoV-2 Nucleocapsid protein expression vector containing a myc-His-tag® whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3.25 seconds.
All lanes:
Immunoprecipitation - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)


Observed band size: 51 kDa



ELISA - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

Plates were coated with Recombinant Human coronavirus SARS-CoV-2 nucleocapsid protein (His tag, ab273530) and Recombinant Human coronavirus SARS Nucleocapsid Protein (ab270829) at 500 ng/ml.
Primary antibody (ab271180) concentration range 0.045-100 ng /ml.

Pre-adsorbed secondary antibody, goat anti-rabbit IgG H&L (HRP, ab97080) used at 1/2000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Immunohistochemical analysis of paraffin-embedded Rat lung labeling SARS-CoV-2 with ab271180 at 1/4000 dilution followed by ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). **Negative control**: No staining on Rat lung. The section was incubated with ab271180 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Goat Anti-Rabbit IgG H&L (HRP polymer).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Immunohistochemical analysis of paraffin-embedded Mouse lung labeling SARS-CoV-2 with ab271180 at 1/4000 dilution followed by ready to use Goat Anti-Rabbit IgG H&L (HRP polymer). **Negative control**: No staining on Mouse lung. The section was incubated with ab271180 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Goat Anti-Rabbit IgG H&L (HRP polymer).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Western blot staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Samples
Lanes 1 & 3: Recombinant Human coronavirus SARS-CoV-2 Nucleocapsid Protein (His tag) (ab273530) (0.2 µg)

Lanes 2 & 4: Recombinant Human coronavirus SARS-CoV-2 Nucleocapsid Protein (His tag) (ab272107) (0.2 µg)

Green: Rabbit monoclonal [EPR24334-118] to SARS-CoV-2 nucleocapsid protein (ab271180)

Red: Mouse monoclonal [HIS.H8] to 6X His tag® (ab18184)

Lanes 1 & 2: Green - ab271180 observed at 50 kDa.


Lanes 3 & 4: Red - loading control Mouse anti-6x HisTag observed at 50 kDa.

ab271180 was shown to react with SARS-CoV-2 nucleocapsid protein in Western blot. Membranes were blocked in 3% milk before incubation with ab271180 and Mouse anti-6x HisTag (ab18184) overnight at 4 °C at a 1/1000 dilution. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1/20000 dilution for 1 h at room temperature before imaging.

All lanes:
Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

All lanes:
Western blot - Recombinant Human coronavirus SARS-CoV-2 nucleocapsid protein (His tag) (ab273530)


Observed band size: 50 kDa



Immunocytochemistry/ Immunofluorescence - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100-permeabilized Vero cells labelling SARS-CoV-2 with ab271180 at 1/1000 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/500 dilution. Fluorescent image showing positve staining in Vero cells after infecting with SARS-CoV-2 virus for 48 h. Nuclear counter stain DAPI. Two indenpendent optical fields were captured for imaging at each given antibody dilution.
This piece of data was kindly provided by Dr. Hangping Yao, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases in Zhejiang University, School of Medicine



Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Western blot staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Exposure time 7.75 seconds

All lanes:
Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180) at 1/1000 dilution

Lane 1:
HEK-293T (human embryonic kidney) transfected with SARS-CoV-2 Nucleocapsid protein expression vector containing a myc-His-tag®, whole cell lysate at 10 µg


Lane 2:
HEK-293T (human embryonic kidney) transfected with SARS-CoV-1 Nucleocapsid protein expression vector containing a myc-His-tag®, whole cell lysate at 10 µg

Lane 3:
HEK-293T (human embryonic kidney) transfected with MERS Nucleocapsid protein expression vector containing a myc-His-tag®, whole cell lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 51 kDa



Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Western blot staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Data was kindly provided by Dr. Hangping Yao, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases in Zhejiang University, School of Medicine.


All lanes:
Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180) at 1/1000 dilution

Lane 1:
SARS-CoV-2 Nucleocapsid Recombinant Protein at 0.1 µg

Lane 2:
SARS-CoV-2 infected Vero for 48 hours

Lane 3:
Uninfected Vero

Observed band size: 51 kDa



Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180)

SARS-CoV-2 nucleocapsid protein Western blot staining using rabbit Anti-SARS-CoV-2 nucleocapsid protein antibody

Lanes 1 - 3: Green - ab271180 observed at 51 kDa. **Lanes 4 - 6:** Red - loading control ab18184 (Mouse monoclonal [HIS.H8] to 6X His tag®) observed at 51 kDa.
ab271180 was shown to react with SARS-CoV-2 nucleocapsid protein in Western blot. Membranes were blocked in 3 % milk in TBS-T (0.1 % Tween®) before incubation with ab271180 and ab18184 (Mouse monoclonal [HIS.H8] to 6X His tag®) overnight at 4 °C at a 1 in 1000 dilution of both antibodies. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.

All lanes:
Western blot - Anti-SARS-CoV-2 nucleocapsid protein antibody [EPR24334-118] (ab271180) at 1/1000 dilution

Lanes 1 and 4:
Western blot - Recombinant Human coronavirus SARS-CoV-2 nucleocapsid protein (His tag) (ab273530) at 0.1 µg

Lanes 2 and 5:
Recombinant Human coronavirus SARS Nucleocapsid Protein (ab270829) at 0.1 µg

Lanes 3 and 6:
Western blot - Recombinant Human coronavirus SARS-CoV-2 Nucleocapsid Protein (His tag) (ab272107) at 0.1 µg

Performed under reducing conditions.

Observed band size: 51 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.