abcam

ab272016

Anti-GPNMB antibody [SP299] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	SP299
Purification technique	Affinity purification Protein A/G
Concentration	0.997 - 1.013 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	Primary incubation for 30 minutes at 4°;C.

WB

Tested

Species	Human
Dilution info	-
Notes	Primary incubation for ;1 hour at room temperature.

IHC-P

Tested

Species	Human
Dilution info	-
Notes	Perform heat mediated antigen retrieval with EDTA buffer pH 8.0 before commencing with IHC staining protocol.Primary incubation for ;10 minutes at room temperature.

Target data

See full target information GPNMB

Function	Could be a melanogenic enzyme.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab272016 is the carrier-free version of ab227695.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	GPNMB also known as osteoactivin is a type I transmembrane glycoprotein with a molecular mass of approximately 115 kDa. This protein is highly expressed in various tissues including the bone skin and brain. It also shows notable presence in certain immune cells such as dendritic cells and macrophages. GPNMB localizes mainly on the cell membrane and within lysosomal compartments indicating its involvement in cellular and signaling processes.
Biological function summary	GPNMB plays versatile roles in promoting cell growth differentiation and repair. It is involved in processes like tissue regeneration possibly due to its role in cell adhesion and migration. The protein also partakes in osteoclast differentiation and bone resorption therefore linking to skeletal homeostasis. GPNMB forms part of a

	broader protein complex at the cellular junctions interacting with integrins to mediate its functions.
Pathways	GPNMB features prominently in osteoclastogenesis and melanoma metastasis pathways. It cooperates with RANKL an important protein in bone metabolism by influencing osteoclast differentiation and activation. In the context of melanoma GPNMB's action in pathways may enhance tumor proliferation and invasion possibly interacting with signaling molecules like AKT or ERK although the specifics remain actively researched.
Associated diseases and disorders	GPNMB has significant links to both osteoarthritis and certain types of cancer particularly breast cancer and melanoma. In osteoarthritis GPNMB expression often increases reflecting involvement in cartilage degradation processes. Its overexpression correlates with tumor progression in melanoma potentially driven by its interaction with integrin β1 which assists in cellular adhesion and migration within the tumor microenvironment.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

9 product images




Immunocytochemistry/ Immunofluorescence - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunocytochemistry/ Immunofluorescence staining of U-87 MG cells using rabbit Anti-GPNMB antibody

This data was developed using ab227695, the same antibody clone in a different buffer formulation.

ab227695 was shown to react with GPNMB in wild-type U-87 MG cells in immunocytochemistry with loss of signal observed in a GPNMB siRNA knockdown cell line. Wild-type and siRNA knockdown cells were mixed and pelleted at a 1:1 ratio on coverslips. The cells were fixed with 4% paraformaldehyde (15 min) then permeabilized with 0.1% Triton X-100 (10min) and then blocked with 1x PBS, 0.01% Triton X-100, 5% BSA, 5% NGS. The cells were then incubated with ab227695 at 1/500 dilution overnight at 4°C followed by a further incubation at room temperature for 1h with a goat anti-rabbit secondary antibody to (Alexa Fluor® 555) at 0.5 µg/ml. Acquisition of the green (wild-type), red (antibody staining) and far-red (siRNA knockdown) channels was performed. Representative grayscale images of the red channel are shown. Wild-type and siRNA knockdown are outlined with yellow and magenta dashed line, respectively. Schematic representation of the mosaic strategy used is shown on the bottom-right panel. Image was acquired with a Zeiss(LSM-880).

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GPNMB antibody

Formalin-fixed, paraffin-embedded human colon tissue stained for GPNMB using ab227695 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GPNMB antibody

Formalin-fixed, paraffin-embedded human reactive lymph node tissue stained for GPNMB using ab227695 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GPNMB antibody

Formalin-fixed, paraffin-embedded human tonsil tissue stained for GPNMB using ab227695 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GPNMB antibody

Formalin-fixed, paraffin-embedded human lung squamous cell carcinoma tissue stained for GPNMB using ab227695 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).




Flow Cytometry - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Flow Cytometry staining using rabbit Anti-GPNMB antibody

Flow Cytometry analysis of A-375 (human malignant melanoma cell line) cells, labeling GPNMB with ab227695 at 1/100 dilution (green) compared to a Rabbit IgG negative control (blue).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GPNMB antibody

Formalin-fixed, paraffin-embedded human cervical squamous cell carcinoma tissue stained for GPNMB using ab227695 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

GPNMB Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-GPNMB antibody

Formalin-fixed, paraffin-embedded human lung tissue stained for GPNMB using ab227695 at 1/100 dilution in immunohistochemical analysis.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab227695).



Western blot - Anti-GPNMB antibody [SP299] - BSA and Azide free (ab272016)

This data was developed using ab227695, the same antibody clone in a different buffer formulation.

ab227695 was shown to react with GPNMB in wild-type U-87 MG cells in Western blot with loss of signal observed in a GPNMB siRNA knockdown cell line. Cell lysates from wild-type U-87 MG transfected with either scrambled siRNA or GPNMB siRNA were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab227695 overnight at 4 °C at a 1/500 dilution. Blots were incubated with secondary antibodies at 0.2 µg/mL before imaging.

This data was provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.

All lanes:
Western blot - Anti-GPNMB antibody [SP299] (ab227695) at 1/500 dilution

Lane 1:
Wild-type U-87 MG transfected with scrambled siRNA control lysate at 60 µg

Lane 2:
U-87 MG transfected with siRNA specifically targeting GPNMB cell lysate at 60 µg

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.