abcam

ab273038

Anti-IL-10RB/IL-10R2 antibody [EPR24691-11]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24691-11
Purification technique	Affinity purification Protein A
Concentration	0.48 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information IL10RB

Function Shared cell surface receptor required for the activation of five class 2 cytokines: IL10, IL22, IL26, IL28, and IFNL1. The IFNLR1/IL10RB dimer is a receptor for the cytokine ligands IFNL2 and IFNL3 and mediates their antiviral activity. The ligand/receptor complex stimulate the activation of the JAK/STAT signaling pathway leading to the expression of IFN-stimulated genes (ISG), which contribute to the antiviral state.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Flow Cytometry - Anti-IL-10RB/IL-10R2 antibody [EPR24691-11] (ab273038)

IL-10RB/IL-10R2 Flow Cytometry staining using rabbit Anti-IL-10RB/IL-10R2 antibody


Flow cytometric analysis of Human peripheral blood mononuclear cell (PBMC) cells labelling IL-10RB/IL-10R2 with ab273038 at 1/500 dilution (0.1ug) (Right) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Left). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab273038. Then stained with anti-CD14 conjugated to BV510. Gated on viable cells.



Flow Cytometry - Anti-IL-10RB/IL-10R2 antibody [EPR24691-11] (ab273038)

IL-10RB/IL-10R2 Flow Cytometry staining using rabbit Anti-IL-10RB/IL-10R2 antibody


Flow cytometric analysis of Human peripheral blood mononuclear cell (PBMC) cells labelling IL-10RB/IL-10R2 with ab273038 at 1/500 dilution (0.1ug) (Right) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Left) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab273038. Then stained with anti-CD3 conjugated to Alexa Fluor® 647. Gated on viable cells.



Flow Cytometry - Anti-IL-10RB/IL-10R2 antibody [EPR24691-11] (ab273038)

IL-10RB/IL-10R2 Flow Cytometry staining using rabbit Anti-IL-10RB/IL-10R2 antibody

Flow cytometric analysis of Human peripheral blood mononuclear cell (PBMC) cells labelling IL-10RB/IL-10R2 with ab273038 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab273038. Then stained with anti-CD56 conjugated to PE. Gated on viable NK cells.



Flow Cytometry - Anti-IL-10RB/IL-10R2 antibody [EPR24691-11] (ab273038)

IL-10RB/IL-10R2 Flow Cytometry staining using rabbit Anti-IL-10RB/IL-10R2 antibody

Flow cytometric analysis of Human peripheral blood mononuclear cell (PBMC) cells labelling IL-10RB/IL-10R2 with ab273038 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab273038. Then stained with anti-CD3 conjugated to Alexa Fluor® 647 and anti-CD14 conjugated to BV510. Gated on viable T cells and monocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.