abcam

ab273506

Anti-HAI-1 antibody [EPR23294-49] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR23294-49
Purification technique	Affinity purification Protein A
Specificity	Unsuitable for human and rat IHC-P.
Concentration	1.067 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Expected

Species	Mouse

Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

WB

Expected

Species	Mouse, Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

IHC-Fr

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	-
Notes	This antibody is not suitable for human and rat species in IHC-P application.

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Target data

See full target information SPINT1

Function	Inhibitor of HGFAC (PubMed:9045658). Inhibits serine protease activity of ST14/matriptase in vitro (PubMed:28710277). Inhibits serine protease activity of TMPRSS13, via the BPTI/Kunitz inhibitor 1 domain (PubMed:20977675).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab273506 is the carrier-free version of ab272721.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	HAI-1 known also as Hepatocyte Growth Factor Activator Inhibitor 1 or SPINT1 is a type 1 transmembrane protein with a molecular mass of around 55 kDa. This protein commonly expresses in epithelial cells particularly in tissues like the stomach and liver. It serves as a serine protease inhibitor playing a role in regulating the activity of certain proteases. Mouse HAI protein comparable in function and structure is often used in research as a model for understanding HAI-1 in humans. Researchers frequently use HAI detectors and HAI-1 ELISA methods for precise quantification and analysis.
Biological function summary	HAI-1 performs a regulatory role by inhibiting proteases such as matriptase and hepsin which are critical for many cellular functions. As part of a complex HAI-1 binds to these proteases controlling their activity to prevent excessive proteolysis which can damage tissues. This regulatory function is key in maintaining epithelial integrity and homeostasis impacting processes like cell growth and migration. Anti-HAI studies show that disrupting this balance can lead to a cascade of biological alterations.
Pathways	Several cellular processes heavily depend on the balance maintained by HAI-1 in protease activity. One significant pathway involves the HGF/c-Met signaling cascade which impacts cell motility and proliferation. By inhibiting matriptase HAI-1 indirectly affects this pathway regulating cellular responses. Additionally HAI-1 also relates to epithelial-mesenchymal transition (EMT) through controlling proteolytic activity which can influence how cells migrate and differentiate in development or healing.
Associated diseases and disorders	Deregulation of HAI-1 often links to cancer progression specifically in cases like breast or prostate cancer. In these contexts improper regulation of proteases can lead to enhanced tumor invasiveness and metastasis. HAI-1's interaction with the c-Met receptor pathway facilitated by HGFA illustrates its significant role in cancer biology. Disorders of the liver such as fibrosis also reveal connections where HAI-1 modulation of protease activity affects extracellular matrix turnover implicating HAI-1 in the fibrogenic response alongside proteins like matriptase.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

5 product images




Flow Cytometry (Intracellular) - Anti-HAI-1 antibody [EPR23294-49] - BSA and Azide free (ab273506)

HAI-1 Flow Cytometry (Intracellular) staining using rabbit Anti-HAI-1 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Daudi (Human Burkitt's lymphoma lymphoblast) (Left panel) / T-47D (Human ductal breast epithelial tumor epithelial cell) (Right panel) cells labelling HAI-1 with ab272721 at 1/600 dilution (0.1µg) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor®488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Negative control: Daudi (PMID: 24070417).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab272721).



Immunocytochemistry/ Immunofluorescence - Anti-HAI-1 antibody [EPR23294-49] - BSA and Azide free (ab273506)


HAI-1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-HAI-1 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized T-47D cells labelling HAI-1 with ab272721 at 1/50 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 (2µg/ml) dilution (Green). Confocal image showing membranous staining in T-47D cell line.
Negative control: Daudi (PMID: 24070417).

ab195889 anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5µg/ml) dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2µg/ml) dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab272721).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HAI-1 antibody [EPR23294-49] - BSA and Azide free (ab273506)

HAI-1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HAI-1 antibody

Immunohistochemical analysis of paraffin-embedded mouse stomach tissue labeling HAI-1 with ab272721 at 1/2000 followed by ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse stomach. The section was incubated with ab272721 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab272721).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HAI-1 antibody [EPR23294-49] - BSA and Azide free (ab273506)

HAI-1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-HAI-1 antibody

Immunohistochemical analysis of paraffin-embedded mouse testis tissue labeling HAI-1 with ab272721 at 1/2000 followed by ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on germ cells of mouse testis (PMID: 16740973). The section was incubated with ab272721 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab272721).



Immunoprecipitation - Anti-HAI-1 antibody [EPR23294-49] - BSA and Azide free (ab273506)

HAI-1 was immunoprecipitated from mouse placenta tissue lysate with ab272721 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab272721 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Mouse placenta tissue lysate 10µg.

Lane 2: ab272721 IP in mouse placenta tissue lysate.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab272721 in mouse placenta tissue lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

This blot was developed using a higher sensitivity ECL substrate.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab272721).

All lanes:
Immunoprecipitation - Anti-HAI-1 antibody [EPR23294-49] (ab272721)

Predicted band size: 58 kDa

Observed band size: 56 kDa, 66 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.