abcam

ab273630

Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free

Recombinant | RabMAb | Advanced Validation

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR23695-115
Purification technique	Affinity purification Protein A
Concentration	1.001 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Mouse
Dilution info	-

Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

ChIP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human, Mouse
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse, Rat
Dilution info	-
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

mIHC

Tested

Species	Mouse
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Expected

Species	Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information CDK7

Function	Serine/threonine kinase involved in cell cycle control and in RNA polymerase II-mediated RNA transcription (PubMed:9852112, PubMed:19136461, PubMed:26257281, PubMed:28768201). Cyclin-dependent kinases (CDKs) are activated by the binding to a cyclin and mediate the progression through the cell cycle. Each different complex controls a specific transition between 2 subsequent phases in the cell cycle. Required for both activation and complex formation of CDK1/cyclin-B during G2-M transition, and for activation of CDK2/cyclins during G1-S transition (but not complex formation). CDK7 is the catalytic subunit of the CDK-activating kinase (CAK) complex. Phosphorylates SPT5/SUPT5H, SF1/NR5A1, POLR2A, p53/TP53, CDK1, CDK2, CDK4, CDK6 and CDK11B/CDK11 (PubMed:9372954, PubMed:9840937, PubMed:19136461, PubMed:26257281, PubMed:28768201). Initiates transcription by RNA polymerase II by mediating phosphorylation of POLR2A at 'Ser-5' of the repetitive C-terminal domain (CTD) when POLR2A is in complex with DNA, promoting dissociation from DNA and initiation (PubMed:19136461, PubMed:26257281, PubMed:28768201). CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation, thus regulating cell cycle progression. CAK complexed to the core-TFIIH basal transcription factor activates RNA polymerase II by serine phosphorylation of the CTD of POLR2A, allowing its escape from the promoter and elongation of the transcripts (PubMed:9852112). Its expression and activity are constant throughout the cell cycle. Upon DNA damage, triggers p53/TP53 activation by phosphorylation, but is inactivated in turn by p53/TP53; this feedback loop may lead to an arrest of the cell cycle and of the transcription, helping in cell recovery, or to apoptosis. Required for DNA-bound peptides-mediated transcription and cellular growth inhibition.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab273630 is the carrier-free version of ab256787.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Cyclin-dependent kinase 7 (Cdk7) also known as CAK is a protein that functions as a serine/threonine kinase. It has a mass of approximately 37 kDa and is expressed in various tissues including the liver heart and testis. Cdk7 forms part of the CDK-activating complex required for the cell cycle's progression and transcription processes. By phosphorylating important sites on target proteins Cdk7 contributes to the regulation of the cell cycle and RNA polymerase II-mediated transcription.
Biological function summary	Cdk7 interacts with other molecules playing a fundamental role in the regulation of cell division and transcription. Cdk7 is a component of the Cdk-activating kinase (CAK) complex alongside cyclin H and MAT1. This complex enables activation of other cyclin-dependent kinases through phosphorylation which is essential for DNA replication and repair. As part of the transcription complex TFIIH Cdk7 enhances RNA polymerase II's transcriptional activity influencing gene expression and cellular responses to external signals.
Pathways	Cdk7 participates in significant regulatory pathways controlling cell cycle and transcription. It is integral to the p53 Signaling and Cell Cycle pathways where it

	facilitates the activation of CDK1 CDK2 and CDK4/6 key players in cell cycle progression from the G1 to S phase. Through these pathways Cdk7 maintains cellular control upon environmental stress responses and DNA damage providing connections to tumor suppressor proteins such as p53.
Associated diseases and disorders	Cdk7 has associations with cancer and neurodegenerative diseases. Its dysregulation can lead to unchecked cell proliferation contributing to the development of various cancers like breast and lung cancer. Cdk7's role in transcription implicates it in disorders like Alzheimer's disease where altered transcriptional regulation is observed. Furthermore its interaction with proteins such as p53 links it to cancer as p53 mutations can result in the loss of control over the cell cycle exacerbating tumorigenic processes.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images




Flow Cytometry (Intracellular) - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Cdk7 Flow Cytometry (Intracellular) staining using rabbit Anti-Cdk7 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized HeLa (Human cervix adenocarcinoma epithelial cell) cells labelling Cdk7 with ab256787 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor®488, ab150077) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab256787).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Cdk7 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cdk7 antibody

Immunohistochemical analysis of paraffin-embedded Mouse testis tissue labeling Cdk7 with ab256787 at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Nuclear staining in mouse testis (PMID: 11319144, PMID: 8521393). The section was incubated with ab256787 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab256787).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Immunohistochemical analysis of paraffin-embedded Mouse breast cancer tissue labeling Cdk7 with ab256787 at 1/100 followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Nuclear staining in mouse breast cancer (PMID: 27301701). The section was incubated with ab256787 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab256787).



Multiplex immunohistochemistry - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Cdk7 Multiplex immunohistochemistry staining of Mouse skin using rabbit Anti-Cdk7 antibody

This data was developed using ab256787, the same antibody clone in a different buffer formulation.
Multiplex immunohistochemistry analysis of formalin/PFA-fixed paraffin-embedded mouse skin staining Collagen VII with ab309143 at a 1/100 dilution, ab256787 anti-Cdk7 used at 1/100 dilution and ab315806 anti-Loricrin used at a 1/500 dilution.

Panel A: merged staining of anti-Collagen VII (green; Opal™520), anti-Cdk7 (magenta; Opal™690) and anti-Loricrin (gray; Opal™570) on mouse skin.

Panel B: anti-Collagen VII staining basement membrane in mouse skin.

Panel C: anti-Cdk7 staining nucleus in mouse skin.


Panel D: anti-Loricrin staining terminally differentiated epidermal cells in mouse skin.

Nuclear DNA was labeled with DAPI (shown in blue).

The section was incubated in three rounds of staining: in the order of ab309143, ab256787 and ab315806 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Cdk7 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cdk7 antibody

Immunohistochemical analysis of paraffin-embedded Rat testis tissue labeling Cdk7 with ab256787 at 1/100 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Nuclear staining in rat testis (PMID: 11319144, PMID: 8521393). The section was incubated with ab256787 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab256787).




Flow Cytometry (Intracellular) - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Cdk7 Flow Cytometry (Intracellular) staining using rabbit Anti-Cdk7 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized NIH/3T3 (Mouse embryonic fibroblast) cells labelling Cdk7 with ab256787 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor®488, ab150077) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab256787).



Immunoprecipitation - Anti-Cdk7 antibody [EPR23695-115] - BSA and Azide free (ab273630)

Cdk7 was immunoprecipitated from 0.35 mg NIH/3T3 (mouse embryonic fibroblast) whole cell lysate with ab256787 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab256787 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: NIH/3T3 (mouse embryonic fibroblast) whole cell lysate 10 ug

Lane 2: ab256787 IP in NIH/3T3 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab256787 in NIH/3T3 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab256787).
All lanes:
Immunoprecipitation - Anti-Cdk7 antibody [EPR23695-115] (ab256787)

Predicted band size: 39 kDa

Observed band size: 39 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.