abcam

ab273877

Anti-Thioredoxin / TRX antibody [EPR23843-48]

Lab Essentials | Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23843-48
Purification technique	Affinity purification Protein A
Concentration	0.513 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/250
Notes	-

Species	Rat

Dilution info	1/250
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Species	Rat
Dilution info	1/500
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat
Dilution info	1/2000
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

Target data

See full target information Txn

Function	Participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions (By similarity). Plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity. Induces the FOS/JUN AP-1 DNA binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity (By similarity). ADF augments the expression of the interleukin-2 receptor TAC (IL2R/P55).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Thioredoxin (TRX) also known as TXN or Trx is a small redox protein with a molecular mass of approximately 12 kDa. It acts as an oxidoreductase enzyme facilitating the reduction of other proteins by cysteine thiol-disulfide exchange. Thioredoxin is ubiquitously expressed across various tissues indicating its function in a broad range of cellular activities. The protein comprises an active site with a conserved Cys-Gly-Pro-Cys sequence critical for its reducing activity.
Biological function summary	Thioredoxin influences cellular redox homeostasis and plays direct roles in regulating cell growth and apoptosis. It participates as a major reductant in cells influencing transcription factors and enzymatic activities that depend on thiol-disulfide exchanges. Thioredoxin can interact with other components like thioredoxin reductase and NADPH to form the thioredoxin system a powerful antioxidant defense mechanism. Diseases typically arise from abnormal regulation of this system highlighting its influence on cellular survival and proliferation.
Pathways	Thioredoxin integrates into significant signaling and metabolic processes. It is an important player in the cellular response to oxidative stress and participates in signaling pathways such as the MAPK and apoptosis pathways. In these pathways thioredoxin reduces the oxidative stress transcription factor AP-1 influencing cell fate decisions. Its interaction with thioredoxin-interacting protein (TXNIP) further modulates cellular responses to oxidative environments.
Associated diseases and disorders	Altered thioredoxin activity associates with cancer and neurodegenerative diseases. Its overexpression links to tumor progression where it promotes survival and resistance to apoptosis. Additionally in the context of neurodegenerative disorders thioredoxin's interaction with amyloid-beta peptides implicates it in Alzheimer's disease by mitigating oxidative stress damage. Through these conditions its role as a protector against oxidative stress remains pivotal.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


10 product images



Immunocytochemistry/ Immunofluorescence - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Thioredoxin / TRX antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 cells labelling Thioredoxin / TRX with ab273877 at 1/250 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing cytoplasmic and nuclear staining in NIH/3T3 cells. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2 ug/ml) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Thioredoxin / TRX antibody

Immunohistochemical analysis of paraffin-embedded Mouse testis tissue labeling Thioredoxin / TRX with ab273877 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on sertoli and leydig cells of mouse testis. The section was incubated with ab273877 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Thioredoxin / TRX antibody

Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labeling Thioredoxin / TRX with ab273877 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse kidney (PMID: 20682242). The

section was incubated with ab273877 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Thioredoxin / TRX antibody

Immunohistochemical analysis of paraffin-embedded Rat kidney tissue labeling Thioredoxin / TRX with ab273877 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on rat kidney. The section was incubated with ab273877 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Flow Cytometry (Intracellular) staining using rabbit Anti-Thioredoxin / TRX antibody


Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized C6 (Rat glial tumor glial cell) cells labelling Thioredoxin / TRX with ab273877 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Flow Cytometry (Intracellular) - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Flow Cytometry (Intracellular) staining using rabbit Anti-Thioredoxin / TRX antibody


Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized NIH/3T3 (Mouse embryonic fibroblast) cells labelling Thioredoxin / TRX with ab273877 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Thioredoxin / TRX antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized C6 cells labelling Thioredoxin / TRX with ab273877 at 1/250 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing cytoplasmic and nuclear staining in C6 cells. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2 ug/ml) dilution.



Western blot - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877) at 1/1000 dilution


All lanes:
Rat kidney tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 12 kDa

Observed band size: 11 kDa



Western blot - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Exposure time: Lane 1, 2: 7.75 seconds Lane 3, 4: 3 minutes.

All lanes:
Western blot - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877) at 1/1000 dilution

Lane 1:
NIH/3T3 (mouse embryonic fibroblast), whole cell lysate at 20 µg

Lane 2:
Neuro-2a (mouse neuroblastoma neuroblast), whole cell lysate at 20 µg

Lane 3:
C2C12 (mouse myoblasts myoblast), whole cell lysate at 20 µg


Lane 4:
PC-12 (rat adrenal gland pheochromocytoma), whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 12 kDa

Observed band size: 11 kDa



Western blot - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877)

Thioredoxin / TRX Western blot staining using rabbit Anti-Thioredoxin / TRX antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-Thioredoxin / TRX antibody [EPR23843-48] (ab273877) at 1/1000 dilution

Lane 1:
Mouse kidney tissue lysate at 20 µg

Lane 2:
Mouse liver tissue lysate at 20 µg

Lane 3:
Mouse testis tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 12 kDa

Observed band size: 11 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.