abcam

ab275022

Anti-CEACAM6 antibody [EPR23956-80]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23956-80
Purification technique	Affinity purification Protein A
Concentration	0.498 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Mouse
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information CEACAM6

Function Cell surface glycoprotein that plays a role in cell adhesion and tumor progression (PubMed:10910050, PubMed:11590190, PubMed:1378450, PubMed:16204051, PubMed:2022629, PubMed:2803308, PubMed:8776764). Intercellular adhesion

occurs in a calcium- and fibronectin-independent manner (PubMed:16204051, PubMed:2022629). Mediates homophilic and heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM5 and CEACAM8 (PubMed:11590190, PubMed:16204051, PubMed:2022629, PubMed:2803308, PubMed:8776764). Heterophilic interaction with CEACAM8 occurs in activated neutrophils (PubMed:8776764). Plays a role in neutrophil adhesion to cytokine-activated endothelial cells (PubMed:1378450). Plays a role in cell migration and cell adhesion to endothelial cells (PubMed:16204051).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CEACAM6 also known as 9a6 CEACAM-6 and 1h7 is a cell surface glycoprotein weighing approximately 35-40 kDa. The protein plays a mechanical role as an

	intercellular adhesion molecule. It displays broad expression across multiple tissues with high levels found in epithelial cells of the gastrointestinal tract respiratory tract and in leukocytes.
Biological function summary	CEACAM6 acts in cell signaling and adhesion processes. It often forms complexes with other members of the CEACAM family such as CEACAM5. This protein is involved in the regulation of cell-cell adhesion which impacts cell migration and organization. It also influences immune reactions by modulating the interactions between pathogens and host cells.
Pathways	CEACAM6 integrates into pathways associated with cell adhesion and immune regulation. It engages in the PI3K-Akt signaling pathway tightly interacting with proteins such as CEACAM1 and CEACAM5. Through these interactions CEACAM6 plays an essential role in cell survival proliferation and immune responses.
Associated diseases and disorders	CEACAM6 shows significant links with cancer especially colorectal and pancreatic cancers. Overexpression of this protein correlates with tumor progression and metastasis. Additionally in inflammatory bowel disease CEACAM6 association with CEACAM7 often results in altered epithelial barrier function contributing to disease pathogenesis.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


9 product images



Immunocytochemistry/ Immunofluorescence - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CEACAM6 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized BxPC-3 BxPC-3 (human pancreas adenocarcinoma cell) cells labelling CEACAM6 with ab275022 at 1/500 (1.048 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing membranous and cytoplamic staining in BxPC-3 cell line. **Low expression control:** LNCaP: PMID: 16204051. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CEACAM6 antibody

Immunohistochemical analysis of paraffin-embedded human spleen tissue labeling CEACAM6 with ab275022 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human spleen. The section was incubated with ab275022 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CEACAM6 antibody

Immunohistochemical analysis of paraffin-embedded human colon cancer tissue labeling CEACAM6 with ab275022 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human colon cancer (PMID: 27284373). The section was incubated with ab275022 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Flow Cytometry - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 Flow Cytometry staining using rabbit Anti-CEACAM6 antibody

Flow cytometric analysis of LNCaP (human prostate carcinoma epithelial cell, Left) / BxPC-3 (human pancreas adenocarcinoma epithelial cell, Right) cells labelling CEACAM6 with ab275022 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.

Negative control: LNCaP.

Gated on viable cells.



Immunoprecipitation - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 was immunoprecipitated from 0.35 mg BxPC-3 (human pancreas adenocarcinoma epithelial cell) whole cell lysate with ab275022 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab275022 at 1/2000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: BxPC-3 (human pancreas adenocarcinoma epithelial cell) whole cell lysate 10 ug

Lane 2: ab275022 IP in BxPC-3 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab275022 in BxPC-3 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 seconds.

All lanes:
Immunoprecipitation - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

Predicted band size: 37 kDa

Observed band size: 40-100 kDa



Immunoprecipitation - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 was immunoprecipitated from 0.35 mg A549 (human lung carcinoma epithelial cell) whole cell lysate with ab275022 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab275022 at 1/2000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: A549 (human lung carcinoma epithelial cell) whole cell lysate 10 ug

Lane 2: ab275022 IP in A549 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab275022 in A549 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 seconds.

All lanes:
Immunoprecipitation - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

Predicted band size: 37 kDa

Observed band size: 40-100 kDa



Western blot - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure times: Lanes 1-2: 15 seconds; Lane3: 3 seconds; Lane 4: 26 seconds.

The molecular weight observed is consistent with what has been described in the literature (PMID: 19244123, 28892050, 16204051).

Low expression control: LNCaP (PMID: 16204051).

All lanes:
Western blot - Anti-CEACAM6 antibody [EPR23956-80] (ab275022) at 1/1000 dilution

Lane 1:
A549 (human lung carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
LNCaP (human prostate carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
BxPC-3 (human pancreas adenocarcinoma epithelial cell) whole cell lysate at 20 µg


Lane 4:
LoVo (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 37 kDa

Observed band size: 40-100 kDa



Western blot - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

CEACAM6 Western blot staining of Human lung tissue lysate using rabbit Anti-CEACAM6 antibody

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 26 seconds.

The molecular weight observed is consistent with what has been described in the literature (PMID: 19244123, 28892050, 16204051).

All lanes:
Western blot - Anti-CEACAM6 antibody [EPR23956-80] (ab275022) at 1/1000 dilution


All lanes:
Human lung tissue lysate at 20 µg

Secondary

All lanes:
VeriBlot for IP secondary antibody(HRP)(ab131366) at 1/1000 dilution

Predicted band size: 37 kDa

Observed band size: 40-100 kDa



Western blot - Anti-CEACAM6 antibody [EPR23956-80] (ab275022)

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 15 seconds.

The other family members have lower homology than the tested proteins.

All lanes:
Western blot - Anti-CEACAM6 antibody [EPR23956-80] (ab275022) at 1/1000 dilution

Lane 1:
His-tagged human CEACAM6 recombinant protein, 20ng

Lane 2:
His-tagged human CEACAM1 recombinant protein (aa35-428), 20ng

Lane 3:
His-tagged human CEACAM5 recombinant protein (aa35-685), 20ng

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 37 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.