abcam

ab275025

Anti-C10 antibody [EPR23475-105]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23475-105
Purification technique	Affinity purification Protein A
Concentration	0.55 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Rat, Mouse, Human
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	1/30
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Expected

Species	Rat
Dilution info	1/500
Notes	-

Not recommended

Species	Human

Dilution info	-
Notes	-

WB

Not recommended

Species	Mouse
Dilution info	-
Notes	Does not give a signal with endogenous expression samples.

Species	Human
Dilution info	-
Notes	Does not give a signal with endogenous expression samples.

Species	Rat
Dilution info	-
Notes	Does not give a signal with endogenous expression samples.

IHC-Fr

Tested

Species	Mouse
Dilution info	1/500
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Not recommended

Species	Rat
Dilution info	-
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Species	Human
Dilution info	-

Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Human
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information Ccl6

Function	Chemotactic factor that attracts mostly macrophage, but it can also attract B cells, CD4(+) lymphocytes and eosinophils.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	C10 also known as complement component C10 plays an important role in the immune system by facilitating cell lysis through the membrane attack complex (MAC) assembly. It is part of the complement cascade traditionally involved in destroying pathogens and infected host cells. The protein has a molecular mass of approximately 30 kilodaltons. C10 expression occurs mainly in liver hepatocytes but its product circulates in the bloodstream and can also be found in various body fluids.
Biological function summary	This complement component carries out an essential function in innate immunity by contributing to the execution of cellular apoptosis and pathogen elimination. C10 becomes a part of the larger MAC complex interacting with other complement proteins such as C5b C6 C7 C8 and C9 to form pores on target cell membranes. These pore formations help in creating a passage for ions and molecules leading to osmotic lysis and subsequent cell death integral for immune defense.
Pathways	C10 integrates into the complement activation pathway particularly within the terminal pathway of the complement system. This pathway involves a cascade of protein interactions initiated by different triggers such as antigen-antibody complexes or pathogen surfaces. Related proteins like C3 and C5 facilitate the formation and activation of MAC focusing on pathogen neutralization. C10's mechanistic role ensures the effective completion of this terminal pathway contributing significantly to both classical and alternative complement pathways.

Associated diseases and disorders	Abnormalities in C10 activity can relate to conditions such as paroxysmal nocturnal hemoglobinuria (PNH) and systemic lupus erythematosus (SLE). In PNH faulty complement regulation due to mutations in complement regulatory proteins like CD55 and CD59 leads to red blood cell destruction where C10's role in MAC formation aggravates hemolysis. In SLE autoantibody production and complement activation create immune complexes that accumulate in organs exacerbating inflammation and tissue damage. Understanding C10's interaction with these proteins provides insight into the therapeutic potential for managing such autoimmune-related conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images




Immunohistochemistry (Frozen sections) - Anti-C10 antibody [EPR23475-105] (ab275025)

C10 Immunohistochemistry (Frozen sections) staining using rabbit Anti-C10 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse colon tissue labeling C10 with ab275025 at 1/500 dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2 ug/ml) dilution (Green). Positive staining on mouse colon (PMID: 7579352). The nuclear counterstain was DAPI (Blue).

Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)at 1/1000 (2 ug/ml) dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-C10 antibody [EPR23475-105] (ab275025)

C10 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-C10 antibody

Immunohistochemical analysis of paraffin-embedded Mouse lung tissue labeling C10 with ab275025 at 1/500 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Postive staining on mouse lung (PMID: 14734772). The section was incubated with ab275025 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Frozen sections) - Anti-C10 antibody [EPR23475-105] (ab275025)

C10 Immunohistochemistry (Frozen sections) staining using rabbit Anti-C10 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse small intestine tissue labeling C10 with ab275025 at 1/500 dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2 ug/ml) dilution (Green). Positive staining on mouse small intestine (PMID: 7579352). The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2 ug/ml) dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-C10 antibody [EPR23475-105] (ab275025)

C10 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-C10 antibody

Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling C10 with ab275025 at 1/500 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Postive staining on mouse spleen (PMID: 15770051). The section was incubated with ab275025 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-C10 antibody [EPR23475-105] (ab275025)

C10 Flow Cytometry (Intracellular) staining using rabbit Anti-C10 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed, 90% methanol permeabilized P388D1 (Mouse lymphoma monocyte; macrophage) (treated with 1ug/ml lipopolysaccharide (LPS) for 4h then add 300ng/ml Brefeldin A for 16h) (Red) / Untreated control (Green) cells labelling C10 with ab275025 at 1/500 dilution (0.1ug) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-C10 antibody [EPR23475-105] (ab275025)

C10 was immunoprecipitated from 0.35 mg Mouse small intestine tissue lysate with ab275025 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab275025 at 1/500 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/1000 dilution.
Lane 1: Mouse small intestine tissue lysate 10 ug

Lane 2: ab275025 IP in Mouse small intestine tissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab275025 in mouse small intestine tissue lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 3 minutes.

The expression profile observed is consistent with what has been described in the literature (PMID: 19812544).

All lanes:
Immunoprecipitation - Anti-C10 antibody [EPR23475-105] (ab275025)

Predicted band size: 13 kDa

Observed band size: 13 kDa



Western blot - Anti-C10 antibody [EPR23475-105] (ab275025)

Blocking and dilution buffer: 5% NFDM/TBST.
Exposure time: 8 seconds.

This product does not recognize CCL9.

All lanes:
Western blot - Anti-C10 antibody [EPR23475-105] (ab275025) at 1/1000 dilution

Lane 1:

His-tagged mouse C10 recombinant protein, 20ng

Lane 2:
His-tagged mouse MIP-1 gamma (CCL9) recombinant protein (aa 22-122), 50ng

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 13 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.