abcam

ab275393

Anti-Semaphorin 4D/CD100 antibody [EPR23866-69]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23866-69
Purification technique	Affinity purification Protein A
Concentration	0.51 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information SEMA4D

Function	Cell surface receptor for PLXNB1 and PLXNB2 that plays an important role in cell-cell signaling (PubMed:20877282). Regulates GABAergic synapse development (By similarity). Promotes the development of inhibitory synapses in a PLXNB1-dependent manner (By similarity). Modulates the complexity and arborization of developing neurites in hippocampal neurons by activating PLXNB1 and interaction with PLXNB1 mediates activation of RHOA (PubMed:19788569). Promotes the migration of cerebellar granule cells (PubMed:16055703). Plays a role in the immune system; induces B-cells to aggregate and improves their viability (in vitro) (PubMed:8876214). Induces endothelial cell migration through the activation of PTK2B/PYK2, SRC, and the phosphatidylinositol 3-kinase-AKT pathway (PubMed:16055703).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Semaphorin 4D also known as CD100 is a transmembrane protein that plays important roles in cellular signaling. It belongs to the semaphorin protein family known for their role in axonal guidance. CD100 has a molecular mass of approximately 150 kDa. This protein expresses mainly in the immune system including T cells and B cells but can also be found in certain neurons and endothelial cells. It can exist in both a membrane-bound form and a soluble form which arise from proteolytic cleavage.
Biological function summary	CD100 influences various cellular functions by interacting with multiple receptors such as Plexin-B1 and CD72 as part of a larger signaling complex. It contributes to immune response regulation by promoting T-cell activation and B-cell aggregation which are essential for adaptive immunity. In the nervous system it guides neural development including axon growth and synaptic plasticity. CD100's interaction with Plexin-B1 plays a significant role in these developmental processes often affecting cell morphology and movement.
Pathways	Semaphorin 4D is involved in both immune and neural signaling pathways. In the immune system it influences the PI3K-Akt signaling pathway which impacts cell survival and immune cell activation. It associates with pathways involving Plexin-B1 and RhoA to regulate cytoskeletal dynamics affecting cell motility and growth. Additionally CD100's role in neural pathways intersects with proteins like neuropilin-1 influencing semaphorin-mediated signaling essential for neural connectivity.
Associated diseases and disorders	Semaphorin 4D has links to cancer and autoimmune diseases. High expression levels of CD100 are observed in several tumor types where it might promote tumor growth and metastasis through its signaling pathways. In autoimmune conditions like multiple sclerosis CD100 affects immune cell interactions and can exacerbate inflammatory responses. Its involvement with proteins such as CD72 and Plexin-B1 in these conditions highlights its contribution to pathogenesis and potential as a therapeutic target.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

2 product images




Flow Cytometry - Anti-Semaphorin 4D/CD100 antibody [EPR23866-69] (ab275393)

Semaphorin 4D/CD100 Flow Cytometry staining using rabbit Anti-Semaphorin 4D/CD100 antibody

Flow cytometric analysis of PC-3 (human prostate adenocarcinoma epithelial cell, Left) / HDLM-2(Human Hodgkin lymphoma, Right) cells labelling Semaphorin 4D/CD100 with ab275393 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.

Negative control: PC-3 (PMID: 23775445).

Gated on viable cells.



Western blot - Anti-Semaphorin 4D/CD100 antibody [EPR23866-69] (ab275393)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression profile observed is consistent with what has been described in the literature (PMID: 23775445).

Negative control: PC-3 (PMID: 23775445)

Exposure time: 136 seconds.

All lanes:
Western blot - Anti-Semaphorin 4D/CD100 antibody [EPR23866-69] (ab275393) at 1/1000 dilution

Lane 1:
HDLM-2 (Human Hodgkin lymphoma) whole cell lysate at 20 µg

Lane 2:
PC-3 (human prostate adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Predicted band size: 96 kDa

Observed band size: 140 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.