abcam

ab275979

Anti-Integrin alpha 4/CD49D antibody [Natalizumab] - Rabbit IgG (Chimeric)

Recombinant

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.02% Proclin 300 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Cell preparation containing ITGA4 protein. The exact immunogen used to generate this antibody is proprietary information. Database link P13612
Clone number	Natalizumab
Purification technique	Affinity purification Protein A
Light chain type	kappa
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	1 µg/mL

Notes	-

Target data

See full target information ITGA4

Function	Integrins alpha-4/beta-1 (VLA-4) and alpha-4/beta-7 are receptors for fibronectin. They recognize one or more domains within the alternatively spliced CS-1 and CS-5 regions of fibronectin. They are also receptors for VCAM1. Integrin alpha-4/beta-1 recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha-4/beta-7 is also a receptor for MADCAM1. It recognizes the sequence L-D-T in MADCAM1. On activated endothelial cells integrin VLA-4 triggers homotypic aggregation for most VLA-4-positive leukocyte cell lines. It may also participate in cytolytic T-cell interactions with target cells. ITGA4:ITGB1 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1-dependent fractalkine signaling (PubMed:23125415). ITGA4:ITGB1 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:18635536, PubMed:25398877). Integrin ITGA4:ITGB1 represses PRKCA-mediated L-type voltage-gated channel Ca(2+) influx and ROCK-mediated calcium sensitivity in vascular smooth muscle cells via its interaction with SVEP1, thereby inhibiting vasocontraction (PubMed:35802072).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

This rabbit monoclonal chimeric antibody was made using the variable domain sequences of the original human IgG4 format, for improved compatibility with existing reagents, assays and techniques.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Integrin alpha 4 also known as CD49D or ITGA4 is a transmembrane receptor with a mass of approximately 150 kDa. This protein is expressed on the surface of leukocytes including T and B lymphocytes monocytes and eosinophils. Integrin alpha 4 functions by interacting mechanically with the extracellular matrix components and endothelial cell counter-receptors like VCAM-1 facilitating cell adhesion and migration. Its expression is vital for the immune cells' ability to move and locate sites of inflammation or injury.
Biological function summary	Integrin alpha 4 forms a heterodimer complex with beta subunits like beta 1 (forming VLA-4) or beta 7 allowing it to mediate adhesion and signaling pathways. In particular the alpha 4 and beta 1 dimerization gives rise to a high-affinity receptor for fibronectin and VCAM-1. This interaction is essential for regulating immune responses and maintaining cell positioning within tissues. The presence of this complex on leukocytes drives their ability to navigate through the vasculature to inflamed areas.
Pathways	Integrin alpha 4 participates in significant signal transduction pathways including the MAPK and PI3K-Akt pathways. These pathways modulate cell survival proliferation and migration. In the MAPK pathway integrin alpha 4 influences cell signaling cascades by interacting with other proteins like TALIN or PAXILLIN which are important for integrin-mediated signal relay. In connection with PI3K-Akt pathway alpha 4 integrins work closely with proteins in regulating cellular adhesion and motility showing a dynamic interplay in immune regulation.
Associated diseases and disorders	Integrin alpha 4 plays an important role in various immune-related diseases such as multiple sclerosis and Crohn's disease. The protein's participation in leukocyte adhesion and trafficking makes it a target for therapeutic intervention. Natalizumab a monoclonal antibody against alpha 4 integrin showcases the critical link between the protein and disease treatment by blocking leukocyte migration to reduce inflammation. Additionally integrin alpha 4's interaction with proteins like VCAM-1 and VLA-4 highlights its relevance in these autoimmune conditions highlighting its importance in disease pathophysiology.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

1 product image



Flow Cytometry - Anti-Integrin alpha 4/CD49D antibody [Natalizumab] - Rabbit IgG (Chimeric) (ab275979)

Integrin alpha 4/CD49D Flow Cytometry staining using rabbit Anti-Integrin alpha 4/CD49D antibody

Flow cytometry of human lymphocytes labeling Integrin alpha 4/CD49D with ab275979 (panel B) compared with an isotype control (panel A) at 1 µg/ml for 30 mins at RT. After washing, bound antibody was detected using Donkey Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150073)) and cells analysed on a FACSCanto flow-cytometer.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.