abcam

ab276132

Anti-Angiotensinogen antibody [EPR24118-2]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24118-2
Purification technique	Affinity purification Protein A
Concentration	0.42 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Rat

Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-Fr

Not recommended

Species	Rat, Human, Mouse
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-P

Tested

Species	Rat
Dilution info	1/500
Notes	-

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information AGT

Function

Essential component of the renin-angiotensin system (RAS), a potent regulator of blood pressure, body fluid and electrolyte homeostasis. Angiotensin-2. Acts directly on vascular smooth muscle as a potent vasoconstrictor, affects cardiac contractility and heart rate through its action on the sympathetic nervous system, and alters renal sodium and water absorption through its ability to stimulate the zona glomerulosa cells of the adrenal cortex to synthesize and secrete aldosterone (PubMed:10619573, PubMed:17138938). Acts by binding to angiotensin receptors AGTR1 and AGTR2 (PubMed:1567413). Also binds the DEAR/FBXW7-AS1 receptor (By similarity). Angiotensin-3. Stimulates aldosterone release. Angiotensin 1-7. Is a ligand for the G-protein coupled receptor MAS1 (By similarity). Has vasodilator and antidiuretic effects. Has an antithrombotic effect that involves MAS1-mediated release of nitric oxide from platelets (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Angiotensinogen also known as AGT is a precursor to angiotensin peptides and plays a mechanical role in the renin-angiotensin system. It is a glycoprotein with a mass of approximately 60 kDa and is mainly produced in the liver. This protein circulates in the bloodstream where it waits for renin to cleave it. Angiotensinogen's presence is significant in various tissues but primarily it functions within the plasma after hepatic secretion.
Biological function summary	Angiotensinogen is critical in the regulation of blood pressure and fluid balance. It serves as the substrate for renin an enzyme that initiates a cascade leading to the production of angiotensin I which further converts into active forms like angiotensin II. Angiotensinogen does not operate as part of a larger protein complex but interacts closely with enzymes and other substrates to mediate its effects on the cardiovascular system.

Pathways	Angiotensinogen is an essential component of the renin-angiotensin-aldosterone system (RAAS) which helps regulate blood pressure and electrolyte balance. The pathway involves the conversion of angiotensinogen to angiotensin I by renin followed by the conversion to angiotensin II by angiotensin-converting enzyme (ACE). This system also has connections with proteins such as aldosterone which is involved in sodium and water retention influencing cardiovascular health.
Associated diseases and disorders	Angiotensinogen links to hypertension and cardiovascular diseases. The protein's interaction with renin and ACE results in the formation of angiotensin II a potent vasoconstrictor that increases blood pressure. Elevated levels of angiotensinogen are often observed in individuals with high 117/85 blood pressure indicating a risk for chronic conditions. Managing angiotensinogen levels through RAAS modulation is a therapeutic strategy for controlling these disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

10 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Angiotensinogen antibody

Immunohistochemical analysis of paraffin-embedded Human placenta tissue labeling Angiotensinogen with ab276132 at 1/500 (0.854 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Cytoplasmic staining on human placenta (PMID: 21215447). The section was incubated with ab276132 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 minutes.



Immunocytochemistry/ Immunofluorescence - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Angiotensinogen antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 100% Methanol permeabilized HepG2 cells labelling Angiotensinogen with ab276132 at 1/100 (4.27 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in HepG2 cell line. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Angiotensinogen antibody

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling Angiotensinogen with ab276132 at 1/500 (0.854 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Cytoplasmic staining on human kidney. The section was incubated with ab276132 for 30 minutes at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 minutes.



Flow Cytometry (Intracellular) - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Flow Cytometry (Intracellular) staining using rabbit Anti-Angiotensinogen antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized HepG2 (human hepatocellular carcinoma epithelial cell) cells labelling Angiotensinogen with ab276132 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Angiotensinogen antibody

Immunohistochemical analysis of paraffin-embedded Human liver tissue labeling Angiotensinogen with ab276132 at 1/500 (0.854 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Cytoplasmic staining on human liver. The section was incubated with ab276132 for 30 minutes at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 minutes.



Immunoprecipitation - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen was immunoprecipitated from 0.35 mg HepG2 (human hepatocellular carcinoma epithelial cell), whole cell lysate with ab276132 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab276132 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HepG2 (human hepatocellular carcinoma epithelial cell), whole cell lysate 10 ug

Lane 2: ab276132 IP in HepG2 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab276132 in HepG2 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 23 seconds.

All lanes:
Immunoprecipitation - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Predicted band size: 53 kDa

Observed band size: 50 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Angiotensinogen antibody

Immunohistochemical analysis of paraffin-embedded Rat kidney tissue labeling Angiotensinogen with ab276132 at 1/500 (0.854 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Cytoplasmic staining on rat kidney (PMID: 28716988). The section was incubated with ab276132 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 minutes.



Western blot - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Western blot staining using rabbit Anti-Angiotensinogen antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Angiotensinogen undergoes glycosylation. The molecular weight observed is consistent with what have been described in literature(PMID: 9694881).

Exposure times:

Lane 1: 15 seconds
Lane 2: 3.25 seconds

All lanes:
Western blot - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132) at 1/1000 dilution

Lane 1:
Human plasma at 20 µL


Lane 2:
Rat plasma at 20 µL

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/100000 dilution

Predicted band size: 53 kDa

Observed band size: 48-70 kDa



Western blot - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Western blot staining using rabbit Anti-Angiotensinogen antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Angiotensinogen undergoes glycosylation. The molecular weight observed is consistent with what has been described in literature(PMID: 9694881).

Lanes 1 & 2 of the blot were developed using a higher sensitivity ECL substrate.

Exposure times: Lane 1, 2: 59 seconds; Lane 3: 3.25 seconds;Lane 4, 5: 15 seconds

All lanes:
Western blot - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132) at 1/1000 dilution

Lane 1:
HepG2 (human hepatocellar carcinoma epithelial cell), whole cell lysate at 20 µg

Lane 2:
Huh7 (human hepatocellar carcinoma epithelial cell), whole cell lysate at 20 µg

Lane 3:
Human lung tissue lysate at 20 µg

Lane 4:
Human liver tissue lysate at 20 µg


Lane 5:
Human kidney tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 53 kDa

Observed band size: 48-70 kDa



Western blot - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132)

Angiotensinogen Western blot staining using rabbit Anti-Angiotensinogen antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Angiotensinogen undergoes glycosylation. The molecular weight observed is consistent with what have been described in literature(PMID: 9694881).

This blot was developed using a higher sensitivity ECL substrate.

Exposure time: 127 seconds.

All lanes:
Western blot - Anti-Angiotensinogen antibody [EPR24118-2] (ab276132) at 1/1000 dilution

Lane 1:
Rat lung tissue lysate at 20 µg

Lane 2:
Rat liver tissue lysate at 20 µg

Lane 3:
Rat kidney tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 53 kDa

Observed band size: 48-70 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.