abcam

ab277523

Anti-NPTX2 antibody [EPR24020-38]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24020-38
Purification technique	Affinity purification Protein A
Concentration	0.504 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

IHC-Fr

Tested

Species	Human
Dilution info	0.5 µg/mL
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Human, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information NPTX2

Function	Likely to play role in the modification of cellular properties that underlie long-term plasticity. Binds to agar matrix in a calcium-dependent manner (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	NPTX2 protein also known as neuronal pentraxin II or NARP weighs approximately 47 kDa. NPTX2 is expressed mostly in the brain especially in neurons. This protein belongs to the neuronal pentraxin family and plays a mechanical role in processes like synaptic plasticity. NPTX2 operates by binding to AMPA receptors influencing their surface expression.
Biological function summary	NPTX2 affects synaptic stabilization by participating in the formation and maintenance of synaptic connections. It forms complexes with AMPA receptor subunits promoting synapses that are more active and facilitating synaptic communication. NPTX2's involvement is essential for normal synaptic function reflecting its significance in neural signaling.
Pathways	NPTX2 plays a role in the glutamatergic signaling pathway an important pathway for synaptic transmission involving excitatory neurotransmitters. NPTX2 works alongside other proteins like GluR1 subunit in this pathway mediating synaptic responses. Through its actions in these pathways NPTX2 contributes to long-term potentiation a process related to learning and memory.
Associated diseases and disorders	NPTX2 has links to neurological conditions such as Alzheimer’s disease and schizophrenia. In Alzheimer's NPTX2 dysregulation associates with synaptic dysfunction possibly involving interactions with beta-amyloid. In schizophrenia altered NPTX2 expression might contribute to glutamatergic signaling imbalances affecting cognitive function. These associations highlight NPTX2's importance in maintaining normal neural processes.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

14 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NPTX2 antibody

Immunohistochemical analysis of paraffin-embedded Human pancreas tissue labeling NPTX2 with ab277523 at 1/2000 (0.229 ug/ml) followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection). Positive staining on islet of human pancreas. The section was incubated with ab277523 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NPTX2 antibody

Immunohistochemical analysis of paraffin-embedded Human cerebrum tissue labeling NPTX2 with ab277523 at 1/2000 (0.229 ug/ml) followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection) was used. Positive staining on human cerebrum. The section was incubated with

ab277523 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunohistochemistry (Frozen sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-NPTX2 antibody


IHC image of NPTX2 staining in a section of frozen normal human pancreas performed on a Leica Biosystems BOND® RX instrument. The section was fixed in 10% paraformaldehyde (10 min) prior to staining. The section was incubated with ab277523, 0.5 ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody. For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times. *Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.



Immunohistochemistry (Frozen sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-NPTX2 antibody

Negative tissue image: IHC image of NPTX2 staining in a section of frozen normal human heart performed on a Leica Biosystems BOND® RX instrument. The section was fixed in 10% paraformaldehyde (10 min) prior to staining. The section was incubated with ab277523, 0.5 ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody. For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times. *Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.




Immunohistochemistry (Frozen sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Frozen sections) staining using rabbit Anti-NPTX2 antibody

IHC image of NPTX2 staining in a section of frozen normal human cerebral cortex performed on a Leica Biosystems BOND® RX instrument. The section was fixed in 10% paraformaldehyde (10 min) prior to staining. The section was incubated with ab277523, 0.5 ug/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody. For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and

antibody incubation times. *Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.



Immunoprecipitation - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 was immunoprecipitated from 0.35 mg Human brain tissue lysate with ab277523 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab277523 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Human brain tissue lysate 10 ug

Lane 2: ab277523 IP in Human brain tissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab277523 in human brain tissue lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds.

All lanes:
Immunoprecipitation - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

Predicted band size: 47 kDa


Observed band size: 50 kDa



Immunocytochemistry/ Immunofluorescence - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-NPTX2 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 100% Methanol permeabilized Beta-TC-6 cells labelling NPTX2 with ab277523 at 1/50 (9.16 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor®488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in Beta-TC-6 cells. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NPTX2 antibody

Immunohistochemical analysis of paraffin-embedded Mouse cerebrum tissue labeling NPTX2 with ab277523 at 1/2000 (0.229 ug/ml) followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection). Positive staining on mouse cerebrum (PMID:31772567). The section was incubated with ab277523 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection) was used.


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Flow Cytometry (Intracellular) staining using rabbit Anti-NPTX2 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized Beta-TC-6 (Mouse pancreas insulinoma beta cell) cells labelling NPTX2 with ab277523 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730)isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-NPTX2 antibody

Immunohistochemical analysis of paraffin-embedded Rat cerebrum tissue labeling NPTX2 with ab277523 at 1/2000 (0.229 ug/ml) followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection). Positive staining on rat cerebrum. The section was incubated with ab277523 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND®RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

NPTX2 was immunoprecipitated from 0.35 mg Mouse brain tissue lysate with ab277523 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab277523 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Mouse brain tissue lysate 10 ug

Lane 2: ab277523 IP in Mouse brain tissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab277523 in mouse brain tissue lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 10 seconds.

All lanes:
Immunoprecipitation - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

Predicted band size: 47 kDa

Observed band size: 50 kDa

Western blot - Anti-NPTX2 antibody [EPR24020-38] (ab277523)



Blocking and diluting buffer and concentration: 5% NFDM/TBST.
NPTX2 is a glycoprotein. The molecular weight observed is consistent with what has been described in the literature (PMID: 8786423, 28440221).

Exposure time: 15 seconds.

All lanes:
Western blot - Anti-NPTX2 antibody [EPR24020-38] (ab277523) at 1/1000 dilution

All lanes:
SH-SY5Y (human neuroblastoma epithelial cell) whole cell lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 47 kDa

Observed band size: 50-60 kDa

Western blot - Anti-NPTX2 antibody [EPR24020-38] (ab277523)



Blocking and diluting buffer and concentration: 5% NFDM/TBST.
NPTX2 is a glycoprotein. The molecular weight observed is consistent with what has been described in the literature (PMID: 8786423, 28440221).

Negative: human kidney, human lung (PMID: 8530029, 8786423).

Exposure times: Lane 1: 15 secondsLane 2-5: 3 minutes.

All lanes:
Western blot - Anti-NPTX2 antibody [EPR24020-38] (ab277523) at 1/1000 dilution

Lane 1:
Human brain tissue lysate at 20 µg

Lane 2:
Human pancreas tissue lysate at 20 µg

Lane 3:
Human testis tissue lysate at 20 µg

Lane 4:
Human kidney tissue lysate at 20 µg


Lane 5:
Human lung tissue lysate at 20 µg

Secondary

All lanes:
VeriBlot for IP secondary antibody(HRP)(ab131366) at 1/1000 dilution

Predicted band size: 47 kDa

Observed band size: 50-60 kDa



Western blot - Anti-NPTX2 antibody [EPR24020-38] (ab277523)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
NPTX2 is a glycoprotein. The molecular weight observed is consistent with what has been described in the literature (PMID: 8786423, 28440221).

Negative: mouse kidney, mouse lung (PMID: 8530029, 8786423).

Exposure time: 15 seconds.

All lanes:
Western blot - Anti-NPTX2 antibody [EPR24020-38] (ab277523) at 1/1000 dilution

Lane 1:
Beta-TC-6 (mouse pancreas insulinoma beta cell) whole cell lysate at 20 µg

Lane 2:
Mouse brain tissue lysate at 20 µg

Lane 3:
Mouse testis tissue lysate at 20 µg

Lane 4:
Mouse kidney tissue lysate at 20 µg

Lane 5:
Mouse lung tissue lysate at 20 µg

Lane 6:
Rat brain tissue lysate at 20 µg

Lane 7:
Rat testis tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 47 kDa

Observed band size: 50-60 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.