abcam

ab279647

Anti-Lin28A antibody [EPR24286-100]

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24286-100
Purification technique	Affinity purification Protein A
Concentration	0.584 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Species	Human

Dilution info	1/500
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000

Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

mIHC

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information LIN28A

Function

RNA-binding protein that inhibits processing of pre-let-7 miRNAs and regulates translation of mRNAs that control developmental timing, pluripotency and metabolism (PubMed:21247876). Seems to recognize a common structural G-quartet (G4) feature in its miRNA and mRNA targets (Probable). 'Translational enhancer' that drives specific mRNAs to polysomes and increases the efficiency of protein synthesis. Its association with the translational machinery and target mRNAs results in an increased number of initiation events per molecule of mRNA and, indirectly, in mRNA stabilization. Binds IGF2 mRNA, MYOD1 mRNA, ARBP/36B4 ribosomal protein mRNA and its own mRNA. Essential for skeletal muscle differentiation program through the translational up-regulation of IGF2 expression. Suppressor of microRNA (miRNA) biogenesis, including that of let-7, miR107, miR-143 and miR-200c. Specifically binds the miRNA precursors (pre-miRNAs), recognizing an 5'-GGAG-3' motif found in pre-miRNA terminal loop, and recruits TUT4 and TUT7 uridylyltransferases (PubMed:18951094, PubMed:19703396, PubMed:22118463, PubMed:22898984). This results in the terminal uridylation of target pre-miRNAs (PubMed:18951094, PubMed:19703396, PubMed:22118463, PubMed:22898984). Uridylated pre-miRNAs fail to be processed by Dicer and undergo degradation. The repression of let-7 expression is required for normal development and contributes to maintain the pluripotent state by preventing let-7-mediated differentiation of embryonic stem cells (PubMed:18951094, PubMed:19703396, PubMed:22118463, PubMed:22898984). Localized to the periendoplasmic reticulum area, binds to a large number of spliced mRNAs and inhibits the translation of mRNAs destined for the ER, reducing the synthesis of transmembrane proteins, ER or Golgi lumen proteins, and secretory proteins. Binds to and enhances the translation of mRNAs for several metabolic enzymes, such as PFKP, PDHA1 or SDHA, increasing glycolysis and oxidative phosphorylation. Which, with the let-7 repression may enhance tissue repair in adult tissue (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

17 product images




Immunocytochemistry/ Immunofluorescence - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Lin28A antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NCCIT (human pluripotent embryonic carcinoma epithelial cell line) cells labeling Lin28A with ab279647 at 1/500 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing nucleolar and cytoplasmic staining in NCCIT cell line. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (Red). The nuclear counterstain was DAPI (Blue).
Negative control: HEK-293T (PMID: 25479749).


Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Flow Cytometry (Intracellular) staining using rabbit Anti-Lin28A antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HEK-293T (human embryonic kidney epithelial cell, Left panel) / NCCIT (human pluripotent embryonic carcinoma epithelial cell, Right panel) cells labeling Lin28A with ab279647 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Negative control: HEK-293T (PMID: 25479749).



Immunoprecipitation - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A was immunoprecipitated from 0.35 mg NCCIT (human pluripotent embryonic carcinoma epithelial cell), whole cell lysate 10 µg with ab279647 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab279647 at 1/5000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/10000 dilution.
Lane 1: NCCIT whole cell lysate 10 µg

Lane 2: ab279647 IP in NCCIT whole cell lysate 10 µg


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab279647 in NCCIT whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 8 seconds
All lanes:
Immunoprecipitation - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Predicted band size: 23 kDa

Observed band size: 25 kDa



Multiplex immunohistochemistry - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Multiplex immunohistochemistry staining of Rat testis tissue using rabbit Anti-Lin28A antibody

Immunohistochemistry analysis of Formalin/PFA-fixed paraffin-embedded sections Rat testis tissue labelling Lin28A with ab279647 at 1:1000 dilution (B), Wilms Tumor Protein with ab267377 at 1:500 dilution (C) and DKKL1 with ab316308 at 1:2000 dilution (D). Opal Polymer HRP Ms + Rb was used as a secondary antibody, and DAPI was used for a nuclear counter stain. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

Panel A: merged staining of anti-Lin28A (green; Opal™520), anti-Wilms Tumor Protein (magenta; Opal™690) and anti-DKKL1 (gray; Opal™570) on rat testis.

Panel B: anti-Lin28A staining undifferentiated spermatogonia in rat testis.


Panel C: ant-Wilms Tumor Protein staining Sertoli cells in rat testis.

Panel D: ant-DKKL11 staining mature sperm cells in rat testis.

Nuclear DNA was labeled with DAPI (shown in blue).

The section was incubated in three rounds of staining: in the order of ab279647, ab267377 and ab316308 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.




Immunohistochemistry (Frozen sections) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunohistochemistry (Frozen sections) staining using rabbit Anti-Lin28A antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen mouse testis tissue labeling Lin28A with ab279647 at 1/100 dilution followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Positive staining on mouse undifferentiated spermatogonia (PMID: 32094118) is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488)at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Multiplex immunohistochemistry - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Multiplex immunohistochemistry staining of Mouse testis tissue using rabbit Anti-Lin28A antibody

Immunohistochemistry analysis of Formalin/PFA-fixed paraffin-embedded sections Mouse testis tissue labelling Lin28A with ab279647 at 1:1000 dilution (B), Wilms Tumor Protein with ab267377 at 1:500 dilution (C) and DKKL1 with ab316308 at 1:2000 dilution (D). Opal Polymer HRP Ms + Rb was used as a secondary antibody, and DAPI was used for a nuclear counter stain. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

Panel A: merged staining of anti-Lin28A (green; Opal™520), anti-Wilms Tumor Protein (magenta; Opal™690) and anti-DKKL1 (gray; Opal™570) on rat testis.

Panel B: anti-Lin28A staining undifferentiated spermatogonia in rat testis.


Panel C: ant-Wilms Tumor Protein staining Sertoli cells in rat testis.

Panel D: ant-DKKL11 staining mature sperm cells in rat testis.

Nuclear DNA was labeled with DAPI (shown in blue).

The section was incubated in three rounds of staining: in the order of ab279647, ab267377 and ab316308 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.



Multiplex immunohistochemistry - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Multiplex immunohistochemistry staining of Rat testis tissue using rabbit Anti-Lin28A antibody

Immunohistochemistry analysis of Formalin/PFA-fixed paraffin-embedded sections Rat testis labelling Lin28A with ab279647 at 1:1000 dilution (B), STAT4 with ab284408 at 1:500 dilution (C) and SCP1 with ab303520 at 1:500 dilution (D). Opal Polymer HRP Ms + Rb was used as a secondary antibody, and DAPI was used for a nuclear counter stain. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

Panel A: merged staining of anti-Lin28A (green; Opal™520), anti-STAT4 (magenta; Opal™690) and anti-SCP1 (gray; Opal™570) on rat testis.

Panel B: anti-Lin28A staining undifferentiated spermatogonia in rat testis.

Panel C: ant-STAT4 staining round spermatids (rs) and elongated spermatids (es) in rat testis.


Panel D: ant-SCP1 staining spermatocytes in rat testis.

Nuclear DNA was labeled with DAPI (shown in blue).

The section was incubated in three rounds of staining: in the order of ab279647, ab284408 and ab303520 for 30 mins at room temperature. Each round was followed by a separate fluorescent

tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.



Multiplex immunohistochemistry - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Multiplex immunohistochemistry staining of Mouse testis tissue using rabbit Anti-Lin28A antibody

Immunohistochemistry analysis of Formalin/PFA-fixed paraffin-embedded sections Mouse testis tissue labelling Lin28A with ab279647 at 1:1000 dilution (B), STAT4 with ab284408 at 1:500 dilution (C) and SCP1 with [ab303520 at 1:500 dilution (D). Opal Polymer HRP Ms + Rb was used as a secondary antibody, and DAPI was used for a nuclear counter stain. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.

Panel A: merged staining of anti-Lin28A (green; Opal™520), anti-STAT4 (magenta; Opal™690) and anti-SCP1 (gray; Opal™570) on mouse testis.

Panel B: anti-Lin28A staining undifferentiated spermatogonia in mouse testis.


Panel C: ant-STAT4 staining round spermatids (rs) and elongated spermatids (es) in mouse testis.

Panel D: ant-SCP1 staining spermatocytes in mouse testis.

Nuclear DNA was labeled with DAPI (shown in blue).

The section was incubated in three rounds of staining: in the order of ab279647, ab284408 and ab303520 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Lin28A antibody

Immunohistochemical analysis of paraffin-embedded mouse testis tissue labeling Lin28A with ab279647 at 1/1000 dilution followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse undifferentiated spermatogonia (PMID: 32094118). The section was incubated with ab279647 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BON$^{D®}$ RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Lin28A antibody

Immunohistochemical analysis of paraffin-embedded rat testis tissue labeling Lin28A with ab279647 at 1/1000 dilution followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat undifferentiated spermatogonia. The section was incubated with ab279647 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Flow Cytometry (Intracellular) staining using rabbit Anti-Lin28A antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NIH/3T3 (mouse embryonic fibroblast, Left panel) / F9 (mouse embryonal carcinoma epithelial cell, Right panel) cells labeling Lin28A with ab279647 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Negative control: NIH/3T3 (PMID: 21962509).




Immunocytochemistry/ Immunofluorescence - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Lin28A antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized F9 (mouse embryonal carcinoma epithelial cell line) cells labelling Lin28A with ab279647 at 1/500 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing nucleolar and cytoplasmic staining in F9 cell line. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).
Negative control: NIH/3T3 (PMID:21962509).

Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Lin28A antibody

Immunohistochemical analysis of paraffin-embedded mouse liver tissue labeling Lin28A with ab279647 at 1/1000 dilution followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). The section was incubated with ab279647 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Negative control: no staining on mouse liver (PMID: 11279525).

Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Lin28A antibody

Immunohistochemical analysis of paraffin-embedded mouse ovary tissue labeling Lin28A with ab279647 at 1/1000 dilution followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse oocytes (PMID: 23378032). The section was incubated with ab279647 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A was immunoprecipitated from 0.35 mg F9 (mouse embryonal carcinoma epithelial cell) whole cell lysate 10 µg with ab279647 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab279647 at 1/5000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/10000 dilution.
Lane 1: F9 whole cell lysate 10 µg

Lane 2: ab279647 IP in F9 whole cell lysate 10 µg

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab279647 in F9 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 5.5 seconds

All lanes:
Immunoprecipitation - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Predicted band size: 23 kDa

Observed band size: 25 kDa



Western blot - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Western blot staining using rabbit Anti-Lin28A antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
The observed MW is consistent with what has been described in the literature (PMID:25479749)

Negative control: HEK-293T (PMID: 25479749)

Exposure time: 10 seconds

All lanes:
Western blot - Anti-Lin28A antibody [EPR24286-100] (ab279647) at 1/5000 dilution

Lane 1:
NCCIT (human pluripotent embryonic carcinoma epithelial cell) whole cell lysate at 10 µg

Lane 2:
JAR (human placenta choriocarcinoma epithelial cell) whole cell lysate at 10 µg


Lane 3:
HEK-293T (human embryonic kidney epithelial cell) whole cell lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 23 kDa

Observed band size: 25 kDa



Western blot - Anti-Lin28A antibody [EPR24286-100] (ab279647)

Lin28A Western blot staining using rabbit Anti-Lin28A antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
The observed MW is consistent with what has been described in the literature (PMID:21962509)

Negative control: NIH/3T3 (PMID:21962509)

Exposure time: 3 seconds

All lanes:
Western blot - Anti-Lin28A antibody [EPR24286-100] (ab279647) at 1/1000 dilution

Lane 1:
F9 (mouse embryonal carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Secondary

All lanes:

Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 23 kDa

Observed band size: 25 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.