abcam

ab280355

Anti-IL3RA/CD123 antibody [EPR24481-20]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24481-20
Purification technique	Affinity purification Protein A
Concentration	0.521 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/2000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information IL3RA

Function	Cell surface receptor for IL3 expressed on hematopoietic progenitor cells, monocytes and B-lymphocytes that controls the production and differentiation of hematopoietic progenitor cells into lineage-restricted cells (PubMed:10527461). Ligand stimulation rapidly induces hetrodimerization with IL3RB, phosphorylation and enzyme activity of effector proteins such as JAK2 and PI3K that play a role in signaling cell proliferation and differentiation. Activation of JAK2 leads to STAT5-mediated transcriptional program (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	IL3RA also known as CD123 or interleukin-3 receptor alpha chain is a transmembrane protein with a molecular weight of approximately 70-80 kDa. This protein frequently expresses on the surface of hematopoietic cells particularly those of myeloid lineage including mast cells and basophils. Researchers use CD123 as a marker for identifying certain cell types in flow cytometry. Antibodies such as anti-CD123 are available for various applications including identifying these cells in research and clinical settings.
Biological function summary	IL3RA forms part of a heterodimeric receptor complex binding with the beta-common (βc) chain to create the high-affinity IL-3 receptor. This complex plays a role in mediating the effects of interleukin-3 a cytokine that promotes the growth and differentiation of pluripotent hematopoietic stem cells. Through this mechanism the receptor influences immune response modulation particularly in allergic reactions and inflammation.

Pathways	IL3RA interacts with the JAK/STAT pathway which is essential for signal transduction and transcription activation in response to IL-3 binding. This pathway involves proteins like Janus kinase 2 (JAK2) and Signal Transducer and Activator of Transcription 5 (STAT5) which transmit signals from IL3RA engagement to the cell nucleus promoting gene expression associated with cell proliferation and survival.
Associated diseases and disorders	IL3RA expression relates closely to certain hematological conditions such as acute myeloid leukemia (AML) and systemic mastocytosis. In AML increased expression of CD123 marks leukemic cells making it a target for therapeutic intervention. The interaction of IL3RA with FLT3 another tyrosine kinase is of particular interest in these disorders as both proteins contribute to the proliferation of malignant cells through related signaling pathways.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

9 product images




Flow Cytometry - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Flow Cytometry staining of human peripheral blood mononuclear cells (PBMCs) using rabbit Anti-IL3RA/CD123 antibody

Flow cytometry staining of human peripheral blood mononuclear cells (PBMCs) with ab280355 (right) or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (left). PBMCs were incubated for 30 mins on ice in 1x PBS containing 10 µg/ml human IgG and 10 % normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody ab280355 or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (1x 10^6 in 100 µl at 1.0 µg/ml (1/2000 dilution)) for 30 mins on ice. The cells were simultaneously stained with CD303.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 dilution for 30 mins on ice

Acquisition of >30000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. Events were gated on live cells.




Flow Cytometry - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Flow Cytometry staining using rabbit Anti-IL3RA/CD123 antibody

Flow cytometric analysis of K-562(Human chronic myelogenous leukemia lymphoblast, Left)/ HDLM-2(Human Hodgkin lymphoma, Right) cells labelling IL3RA/CD123 with ab280355 at 1/500 dilution (0.1ug)/ red compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Negative control: K-562 cell (PMID: 26531164).

Gated on viable cells.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IL3RA/CD123 antibody

Immunohistochemical analysis of paraffin-embedded human Hodgkin's lymphoma tissue labeling IL3RA/CD123 with ab280355 at 1/500 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human Hodgkin's lymphoma. The section was incubated with ab280355 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Flow Cytometry - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Flow Cytometry staining using rabbit Anti-IL3RA/CD123 antibody

Flow cytometric analysis of human peripheral blood mononuclear cell (PBMC) cells labelling IL3RA/CD123 with ab280355 at 1/500 dilution (0.1ug)/ Right compared with a Rabbit monoclonal

IgG (ab172730) / Left isotype control. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Cells were stained with rabbit IgG or ab280355. Then stained with anti-HLA-DR conjugated to PE.

Gated on viable cells.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IL3RA/CD123 antibody

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling IL3RA/CD123 with ab280355 at 1/500 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human tonsil (PMID: 27340692, PMID:15728234, PMID:28205524). The section was incubated with ab280355 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunocytochemistry/ Immunofluorescence - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IL3RA/CD123 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HDLM-2 cells labelling IL3RA/CD123 with ab280355 at 1/1000 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing membranous staining in HDLM-2 cells.
Negative control: K-562 (PMID: 26531164).

ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) at a 1/200 dilution (Red). The nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Immunoprecipitation - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 was immunoprecipitated from 0.35 mg HDLM-2 (human hodgkin lymphoma) whole cell lysate 10 µg with ab280355 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab280355 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HDLM-2 whole cell lysate 10 µg

Lane 2: ab280355 IP in HDLM-2 whole cell lysate


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab280355 in HDLM-2 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 32 seconds
All lanes:
Immunoprecipitation - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

Predicted band size: 43 kDa

Observed band size: 60 kDa



Western blot - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

IL3RA/CD123 Western blot staining of KG-1a (human bone marrow myeloblast) whole cell lysate using rabbit Anti-IL3RA/CD123 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
The expression profile/molecular weight observed is consistent with what has been described in the literature (PMID:30647406).

Exposure time: 103 seconds

All lanes:
Western blot - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355) at 1/1000 dilution


All lanes:
KG-1a (human bone marrow myeloblast) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 43 kDa

Observed band size: 60 kDa



Western blot - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
The expression profile/molecular weight observed is consistent with what has been described in the literature (PMID:30647406).

Negative control: K-562 cell (PMID: 30647406)

Exposure time: 37 seconds

All lanes:
Western blot - Anti-IL3RA/CD123 antibody [EPR24481-20] (ab280355) at 1/1000 dilution

Lane 1:
Human tonsil tissue lysate at 40 µg

Lane 2:
HDLM-2 (human hodgkin lymphoma) whole cell lysate at 20 µg

Lane 3:
K-562 (human chronic myelogenous leukemia lymphoblast) whole cell lysate at 20 µg

Secondary

Lane 1:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Lanes 2 - 3:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 43 kDa

Observed band size: 60 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.