abcam

ab280953

Anti-SARS-CoV2 orf3a antibody [EPR24848-9]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24848-9
Purification technique	Affinity purification Protein A
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Transfected cell lysate - SARS-CoV-2
Dilution info	1/1000
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell line - SARS-CoV-2, Recombinant full length protein - SARS-CoV-2
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Transfected cell line - SARS-CoV-2
Dilution info	1/500
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell lysate - SARS-CoV-2, Recombinant full length protein - SARS-CoV-2
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Transfected cell line - SARS-CoV-2

Dilution info	1/500
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell lysate - SARS-CoV-2, Recombinant full length protein - SARS-CoV-2
Dilution info	-
Notes	-

WB

Tested

Species	Transfected cell lysate - SARS-CoV-2
Dilution info	1/1000
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell line - SARS-CoV-2, Recombinant full length protein - SARS-CoV-2
Dilution info	-
Notes	-

IHC-P

Tested

Species	Transfected cell line - SARS-CoV-2
Dilution info	1/1000
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell lysate - SARS-CoV-2, Recombinant full length protein - SARS-CoV-2
Dilution info	-
Notes	-

ELISA

Tested

Species	Recombinant full length protein - SARS-CoV-2
Dilution info	-
Notes	-

Predicted

Species	SARS-CoV-2
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell lysate - SARS-CoV-2, Transfected cell line - SARS-CoV-2
Dilution info	-

Notes	-

Target data

See full target information SARS CoV-2 ORF3a Protein

Function	Plays a role in viral egress via lysosomal trafficking (PubMed:33157038, PubMed:33422265). Forms homotetrameric ion channels (viroporins) localized at endosomes and lysosomes, that may induce deacidification of lysosomes, allowing safe egress of virions via lysosomal trafficking (PubMed:33157038, PubMed:33422265, PubMed:34158638). Also blocks autolysosome formation by binding and sequestering the host component VPS39 for homotypic fusion and protein sorting (HOPS) on late endosomes (PubMed:33422265). This prevents fusion of autophagosomes with lysosomes, disrupting autophagy and facilitating virus egress (PubMed:33422265). Induces host RETREG1/FAM134B-dependent reticulophagy by interacting with host HMGB1 and enhancing the association between HMGB1 and host BECN1 (PubMed:35239449). This induces endoplasmic reticulum stress and inflammatory responses and facilitates viral infection (PubMed:35239449).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The SARS-CoV-2 ORF3a protein also known as ORF3 is an accessory protein of the SARS-CoV-2 virus weighing approximately 275 amino acids. It is expressed in infected host cells and plays an important role in the virus's lifecycle. ORF3a is known for its role as an ion channel involved in the viral replication cycle. Its functions include altering host cell processes to facilitate viral propagation and possibly contributing to viral pathogenesis.
Biological function summary	The SARS-CoV-2 ORF3a protein interferes with the host immune response promoting virus survival and replication. ORF3a is implicated in the enhancement of viral pathogenic effects by modulating innate and adaptive immunity. Though not part of a large known stable complex ORF3a interacts with other viral and host proteins impacting processes like apoptosis and inflammation.
Pathways	ORF3a affects several critical processes within the host cell. It plays a role in the apoptosis pathway influencing cell death mechanisms that help the virus evade immune defenses. Additionally it modulates the NF-kB signaling pathway which is vital for immune response regulation. Other related proteins include ORF7a and NSP6 which work alongside ORF3a to manipulate host cell environments to the virus's advantage.
Associated diseases and disorders	The SARS-CoV-2 ORF3a protein is directly connected to COVID-19 impacting the severity and progression of the disease. Its interaction with host proteins contributes to dysregulated immune responses and cytokine storms characteristic of severe COVID-19 cases. The protein's role in apoptosis links it to potential lung tissue damage exacerbating respiratory conditions associated with COVID-19 infections.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

9 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV2 orf3a antibody

Immunohistochemical analysis of paraffin-embedded HEK-293T cells labelling SARS CoV-2 ORF3a Protein with ab280953 at 1/1000 (0.53 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Panel A : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) transfected with SARS CoV-2 ORF3a Protein expression vector containing a myc-His-tag® Panel B: HEK-293T transfected with SARS-CoV-1 ORF3a Protein vector containing a myc-His-tag®. Panel C: Incubate HEK-293T transfected with SARS CoV-2 ORF3a Protein expression vector containing a myc-His-tag® with secondary antibody only, but not ab280953. Panel D: HEK-293T transfected with control vector containing a myc-His-tag®. Positive staining on (A) HEK-293T transfected with SARS-CoV-2 Protein 3a expression vector. No staining on (B) HEK-293T transfected with a SARS-CoV-1 Protein 3a expression vector, (C) HEK-293T transfected with SARS-CoV-2 Protein 3a expression vector and incubated with secondary antibody only and (D) HEK-293T transfected with empty vector. The section was incubated with ab280953 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Flow Cytometry (Intracellular) - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Flow Cytometry (Intracellular) staining using rabbit Anti-SARS-CoV2 orf3a antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HEK-293T (human embryonic kidney) transfected with myc-tagged SARS CoV-2 ORF3a Protein expression

construct (Right) with ab280953 at 1/500 dilution compared with HEK-293T (human embryonic kidney) transfected with myc-tagged SARS-COV-1 ORF3a Protein expression construct (Left).
A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077 at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV2 orf3a antibody

Immunohistochemical analysis of paraffin-embedded Human lung tissue labelling SARS CoV-2 ORF3a Protein with ab280953 at 1/1000 (0.53 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). No staining on human lung. The section was incubated with ab280953 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS CoV-2 ORF3a Protein was immunoprecipitated from 0.35 mg HEK-293T (human embryonic kidney) transfected with SARS CoV-2 ORF3a Protein expression vector containing a myc-His-tag®, whole cell lysate 20 µg with ab280953 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab280953 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HEK-293T (human embryonic kidney) transfected with SARS CoV-2 ORF3a Protein expression vector containing a myc-His-tag®, whole cell lysate 20 µg

Lane 2: ab280953 IP in HEK-293T transfected with SARS CoV-2 ORF3a Protein expression vector containing a myc-His-tag® whole cell lysate


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab280953 in HEK-293T transfected with SARS CoV-2 ORF3a Protein expression vector containing a myc-His-tag® whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 seconds.

All lanes:
Immunoprecipitation - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

Observed band size: 23 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV2 orf3a antibody

Immunohistochemical analysis of paraffin-embedded Rat lung tissue labelling SARS CoV-2 ORF3a Protein with ab280953 at 1/1000 (0.53 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). No staining on rat lung. The section was incubated with ab280953 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SARS-CoV2 orf3a antibody

Immunohistochemical analysis of paraffin-embedded Mouse lung tissue labelling SARS CoV-2 ORF3a Protein with ab280953 at 1/1000 (0.53 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). No staining on mouse lung. The section was incubated with ab280953 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection) was used.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




ELISA - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

Recombinant Anti-SARS CoV-2 ORF3a Protein antibody [EPR24848-9] (ab280953)
Plates were coated with a recombinant protein ORF3 at 1000 ng/ml.

Primary antibody (ab280953) concentration range 0.045-100 ng /ml.

Pre-adsorbed secondary antibody, goat anti-rabbit IgG H&L (HRP, ab97080) used at 1/2000 dilution.



Western blot - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Western blot staining using rabbit Anti-SARS-CoV2 orf3a antibody

Lanes 1 – 4: Green – ab280953 observed at 26 kDa.
Lanes 5 – 8: Red – loading control ab18184 (Mouse monoclonal [HIS.H8] to 6X His tag®) observed at 26 kDa.

ab280953 was shown to bind specifically to SARS CoV-2 ORF3a Protein in Western blot. Samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) at 1/20000 dilution.

Blocking buffer: 3% milk in TBS-0.1 % Tween® 20 (TBS-T)
All lanes:
Western blot - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953) at 1/1000 dilution


Lanes 1 and 5:
Expi293 transfected with SARS-CoV-2 ORF3a protein expression vector (His-tagged) cell lysate at 40 µg

Lanes 2 and 6:
Expi293 transfected with SARS-CoV-2 ORF3a protein expression vector (His-tagged) cell lysate at 20 µg

Lanes 3 and 7:
Expi293 transfected with SARS-CoV-1 ORF3a protein expression vector (His-tagged) cell lysate at 40 µg

Lanes 4 and 8:
Expi293 transfected with SARS-CoV-1 ORF3a protein expression vector (His-tagged) cell lysate at 20 µg

Observed band size: 26 kDa



Western blot - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953)

SARS-CoV2 orf3a Western blot staining using rabbit Anti-SARS-CoV2 orf3a antibody

Blocking and Diluting buffer and concentration: 5% NFDM/TBST
Exposure time 5.5 seconds

All lanes:
Western blot - Anti-SARS-CoV2 orf3a antibody [EPR24848-9] (ab280953) at 1/1000 dilution

Lane 1:
HEK-293T (human embryonic kidney) transfected with an empty vector (vector control) containing a myc-His-tag®, whole cell lysate at 20 µg

Lane 2:
HEK-293T transfected with SARS-CoV-2 ORF3a Protein expression vector containing a myc-His-tag®, whole cell lysate at 20 µg

Lane 3:
HEK-293T transfected with SARS-CoV-1 ORF3a Protein expression vector containing a myc-His-tag®, whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 23 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.