abcam

ab281845

Anti-IL-2RG antibody [EPR24561-27]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24561-27
Purification technique	Affinity purification Protein A
Concentration	0.559 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/100
Notes	-

Species	Human

Dilution info	1/100
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

WB

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

Target data

See full target information IL2RG

Function	Common subunit for the receptors for a variety of interleukins. Probably in association with IL15RA, involved in the stimulation of neutrophil phagocytosis by IL15 (PubMed:15123770).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Flow Cytometry - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

IL-2RG Flow Cytometry staining using rabbit Anti-IL-2RG antibody


Flow cytometric analysis of Human peripheral blood mononuclear cell (PBMC) cells labelling IL-2RG with ab281845 at 1/50 dilution (1ug)/ Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab281845. Then stained with anti-CD4 conjugated to Alexa Fluor® 647. Gated on viable cells.



Immunocytochemistry/ Immunofluorescence - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

IL-2RG Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IL-2RG antibody


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized U-937 cells labelling IL-2RG with ab281845 at 1/100 (5.59 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in U-937 cell line is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Flow Cytometry - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

IL-2RG Flow Cytometry staining using rabbit Anti-IL-2RG antibody

Flow cytometric analysis of HT-29 (Human colorectal adenocarcinoma epithelial cell, Left)/ U937 (human histiocytic lymphoma monocyte, Right) cells labelling IL-2RG with ab281845 at 1/50 dilution (1ug)/ (red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. **Negative control**: HT-29 (PMID: 8530527) Gated on viable cells.



Immunoprecipitation - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

IL-2RG was immunoprecipitated from 0.35 mg U-937 (human histiocytic lymphoma monocyte) whole cell lysate 10 ug with ab281845 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab281845 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: U-937 (human histiocytic lymphoma monocyte) whole cell lysate 10 ug

Lane 2: ab281845 IP in U-937 whole cell lysate 10 ug


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab281845 in U-937 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes
All lanes:
Immunoprecipitation - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

Predicted band size: 42 kDa


Observed band size: 64 kDa



Immunocytochemistry/ Immunofluorescence - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

IL-2RG Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IL-2RG antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized CTLL-2 cells labelling IL-2RG with ab281845 at 1/100 (5.59 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in CTLL-2 cell line is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Flow Cytometry - Anti-IL-2RG antibody [EPR24561-27] (ab281845)

IL-2RG Flow Cytometry staining using rabbit Anti-IL-2RG antibody

Flow cytometric analysis of Mouse blood cells cells labelling IL-2RG with ab281845 at 1/50 dilution (1ug)/ Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab281845. Then stained with anti-CD4 conjugated to Alexa Fluor® 647. Gated on viable cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.