abcam

ab281923

Anti-C5a antibody [EPR24303-10]

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24303-10
Purification technique	Affinity purification Protein A
Concentration	0.504 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information C5

Function

Precursor of the C5a anaphylatoxin and complement C5b components of the complement pathways, which consist in a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed:12878586, PubMed:18204047, PubMed:30643019, PubMed:6554279). Activated downstream of classical, alternative, lectin and GZMK complement pathways (PubMed:12878586, PubMed:18204047, PubMed:30643019, PubMed:39914456, PubMed:39814882, PubMed:6554279). Complement C5b. Component of the membrane attack complex (MAC), a multiprotein complex activated by the complement cascade, which inserts into a target cell membrane and forms a pore, leading to target cell membrane rupture and cell lysis (PubMed:26841837, PubMed:27052168, PubMed:30552328, PubMed:30643019). Complement C5b is generated following cleavage by C5 convertase and initiates formation of the MAC complex: C5b binds sequentially C6, C7, C8 and multiple copies of the pore-forming subunit C9 (PubMed:30552328, PubMed:30643019). During MAC complex assembly, the C5b6 subcomplex, composed of complement C5b and C6, associates with the outer leaflet of target cell membrane, reducing the energy for membrane bending (PubMed:30552328, PubMed:32569291). C5a anaphylatoxin. Mediator of local inflammatory process released following cleavage by C5 convertase (PubMed:8182049, PubMed:9553099). Acts by binding to its receptor (C5AR1 or C5AR2), activating G protein-coupled receptor signaling and inducing a variety of responses including

intracellular calcium release, contraction of smooth muscle, increased vascular permeability, and histamine release from mast cells and basophilic leukocytes (PubMed:36806352, PubMed:37852260, PubMed:37169960, PubMed:8182049, PubMed:9553099). C5a is also a potent chemokine which stimulates the locomotion of polymorphonuclear leukocytes and directs their migration toward sites of inflammation (PubMed:342601, PubMed:37852260, PubMed:37169960, PubMed:5765461, PubMed:8182049, PubMed:9553099).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	C5a also known as complement component 5a is a 74-amino acid peptide with a molecular mass of approximately 11 kDa. It is expressed in the blood and is a fragment that forms when the complement system part of the body's immune

	response activates. As a potent anaphylatoxin C5a plays a role in mediating inflammation and recruiting white blood cells to an infection site. It interacts with C5a receptors C5aR1 and C5aR2 located on various immune cells including neutrophils monocytes and macrophages.
Biological function summary	C5a functions as part of the complement cascade a complex series of events essential for opsonization cell lysis and inflammation. C5a binds to its receptors initiating cell responses such as chemotaxis increased vascular permeability and release of pro-inflammatory cytokines. These processes help in pathogen clearance making C5a a significant player in innate immunity. C5a exists as part of the larger complement system that includes proteins like C3 and C5 forming an important arm of the body's defense against pathogens.
Pathways	C5a integrates into the complement pathway and the coagulation cascade. It drives actions in the inflammatory response enhancing immune defense by promoting cytokine production and leukocyte migration. Important interactions occur with complement proteins such as C3 and C5b-9 key elements of the terminal complement complex. By influencing these pathways C5a directly contributes to both immune modulation and inflammatory processes.
Associated diseases and disorders	C5a is linked to conditions such as sepsis and rheumatoid arthritis. In these diseases C5a contributes to excessive inflammation and tissue damage. C5a's interaction with C5aR1 and other receptors like CR1 involved in inflammatory regulation exemplifies its role in advancing disease states. Therapeutically targeting C5a and its receptors remains an active area of research aiming to develop treatments for conditions driven by dysregulated complement activation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Flow Cytometry (Intracellular) - Anti-C5a antibody [EPR24303-10] (ab281923)

C5a Flow Cytometry (Intracellular) staining using rabbit Anti-C5a antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HEK-293 (Human embryonic kidney epithelial cell, Left)/ HepG2 (human hepatocellular carcinoma epithelial cell, Right) cells labelling C5a with ab281923 at 1/50 dilution (1ug)/ (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody. No expression was observed in cell line: HEK-293.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-C5a antibody [EPR24303-10] (ab281923)

C5a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-C5a antibody

Immunohistochemical analysis of paraffin-embedded Human colon tissue labelling C5a with ab281923 at 1/500 dilution (1.092 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on plasma of human colon. The section was incubated with ab281923 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-C5a antibody [EPR24303-10] (ab281923)

C5a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-C5a antibody

Immunohistochemical analysis of paraffin-embedded Human liver tissue labelling C5a with ab281923 at 1/500 dilution (1.092 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human liver. The section was incubated with ab281923 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-C5a antibody [EPR24303-10] (ab281923)

C5a was immunoprecipitated from 0.35 mg Human plasma with ab281923 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab281923 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: Human plasma 10 µL

Lane 2: ab281923 IP in Human plasma


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab281923 in human plasma

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 6 seconds.

All lanes:
Immunoprecipitation - Anti-C5a antibody [EPR24303-10] (ab281923)

Observed band size: 110 kDa



Western blot - Anti-C5a antibody [EPR24303-10] (ab281923)

C5a Western blot staining using rabbit Anti-C5a antibody

Western blot: Rabbit monoclonal [EPR24303-10] to C5a ab281923 staining at 1/1000 dilution, shown in green; Mouse anti-CANX (ab238078) loading control staining at 1/20,000 dilution, shown in magenta. A band was observed at 180 kDa in Wild-type A549 cell lysates with no signal observed at this size in C5 knockout A549 cell line (ab326016). To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5pc Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-C5a antibody [EPR24303-10] (ab281923) at 1/1000 dilution

Lane 1:
Wild-type A549 at 20 µg

Lane 2:
Western blot - Human C5 knockout A549 cell line (ab326016) at 20 µg

Lane 2:
C5 knockout A549 at 20 µg

Lane 3:
HepG2 at 20 µg

Lane 4:
U-2 OS at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 188 kDa

Observed band size: 180 kDa



Western blot - Anti-C5a antibody [EPR24303-10] (ab281923)

C5a Western blot staining using rabbit Anti-C5a antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
110 kDa Complement C5 alpha chain is observed. The molecular weight is consistent with what has been described in the literature (PMID: 24071736, 23401592, 25387724).

Very low or no expression of C5 alpha chain is observed in testis, skeletal muscle.

Exposure time: 6 seconds.

All lanes:
Western blot - Anti-C5a antibody [EPR24303-10] (ab281923) at 1/1000 dilution

Lane 1:
Human plasma at 10 µL

Lane 2:
Human testis tissue lysate at 20 µg

Lane 3:
Human skeletal muscle tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/1000 dilution

Observed band size: 110 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.