abcam

ab283500

Anti-Cofilin (phospho S3) antibody [EPR24753-28]

Lab Essentials | Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24753-28
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse, Synthetic peptide
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Rat, Human, Synthetic peptide
Dilution info	-
Notes	-

Dot

Tested

Species	Synthetic peptide
Dilution info	1/1000
Notes	-

Expected

Species	Human, Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/600
Notes	-

Expected

Species	Mouse
Dilution info	1/600
Notes	-

Species	Rat
Dilution info	-
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/5000
Notes	-

Species	Rat
Dilution info	1/5000
Notes	-

Species	Human
Dilution info	1/5000
Notes	-

Not recommended

Species	Synthetic peptide
Dilution info	-
Notes	-

Target data

See full target information CFL1 phospho S3

Function Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity (PubMed:11812157). In conjunction with the subcortical maternal complex (SCMC), plays an essential role for zygotes to progress beyond the first embryonic cell divisions via regulation of actin dynamics (PubMed:15580268). Required for the centralization of the mitotic spindle and symmetric division of zygotes (By similarity). Plays a role in the regulation of cell morphology and cytoskeletal organization in epithelial cells (PubMed:21834987). Required for the up-regulation

of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed:23633677). Required for neural tube morphogenesis and neural crest cell migration (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Cofilin-1 also known just as cofilin is a small actin-modulating protein with a molecular weight of approximately 19 kDa. It is part of the actin chemical structure regulation and plays an important role in actin filament dynamics by severing and depolymerizing actin filaments. Cofilin-1 expresses widely in both muscle and non-muscle tissues modulating the actin cytoskeleton which is critical for cell movement shape and various signaling pathways. The phosphorylated form of this

	protein known as phospho-cofilin or p-cofilin is another significant state to consider.
Biological function summary	Through its interaction with actin filaments cofilin-1 functions to regulate the length and turnover of the actin cytoskeleton within cells. Cofilin-1 is part of a critical complex where it binds to actin monomers and filaments to facilitate the disassembly and recycling of actin. This regulation impacts cellular activities such as migration endocytosis and division influencing how cells respond to internal and external signals. Cofilin-1's ability to bind and sever actin filaments forms an important part of its functional mechanism altering the cytoskeletal architecture for various cellular processes.
Pathways	Cofilin-1 participates principally in the actin dynamics component of the cytoskeleton remodeling pathway. Within this framework cofilin-1 acts alongside other proteins like actin-depolymerizing factor (ADF) and tropomyosin influencing assembly and disassembly cycles of actin filaments. The Rho family of GTPases regulates cofilin-1 activity through signaling pathways such as the Rho/ROCK/LIMK pathway. This signaling regulation affects actin filament organization impacting cellular processes like axonal guidance and directed cell movement.
Associated diseases and disorders	There is a strong connection between cofilin-1 dysregulation and neurodegenerative diseases such as Alzheimer's disease and certain types of cancer. In Alzheimer's disease cofilin-1 activity influenced by abnormal cofilin phosphorylation forms part of the pathological mechanisms that lead to synaptic dysfunction and neurofibrillary tangles. In the context of cancer cofilin-1 collaborates with proteins such as RhoA and LIM kinase contributing to cancer cell invasion and metastasis by reorganizing the actin cytoskeleton to facilitate cell motility. Understanding these interactions helps in targeting therapeutic strategies against these conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

11 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cofilin (phospho S3) antibody

Immunohistochemical analysis of paraffin-embedded Human stomach tissue labelling Cofilin (phospho S3) with ab283500 at 1/5000 (0.121 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human stomach without alkaline phosphatase treatment (image A). No signal was detected when tissues were treated with alkaline phosphatase (image B). The section was incubated with ab283500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Immunocytochemistry/ Immunofluorescence - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Cofilin (phospho S3) antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HeLa cells labelling Cofilin (phospho S3) with ab283500 at 1/1000 (0.604 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing nuclear and cytoplasmic staining in HeLa cells, no staining in HeLa treated with Alkaline Phosphatase (37°C) for 1 hour is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5 ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cofilin (phospho S3) antibody

Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue labelling Cofilin (phospho S3) with ab283500 at 1/5000 (0.121 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human breast carcinoma without alkaline phosphatase treatment (image A). (PMID: 27799793) No signal was detected when tissues were treated with alkaline phosphatase (image B). The section was incubated with ab283500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins


---



Flow Cytometry (Intracellular) - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Flow Cytometry (Intracellular) staining using rabbit Anti-Cofilin (phospho S3) antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HeLa (Human cervix adenocarcinoma epithelial cell) cells labelling Cofilin (phospho S3) with ab283500 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cofilin (phospho S3) antibody

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labelling Cofilin (phospho S3) with ab283500 at 1/5000 (0.121 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse colon without alkaline phosphatase treatment (image A). (PMID: 15649475) No signal was detected when tissues were treated with alkaline phosphatase (image B). The section was incubated with ab283500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins

---




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Cofilin (phospho S3) antibody

Immunohistochemical analysis of paraffin-embedded Rat colon tissue labelling Cofilin (phospho S3) with ab283500 at 1/5000 (0.121 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat colon without alkaline phosphatase treatment (image A). No signal was detected when tissues were treated with alkaline phosphatase (image B). The section was incubated with ab283500 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Western blot staining using rabbit Anti-Cofilin (phospho S3) antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 3 seconds

All lanes:
Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500) at 1/1000 dilution

Lane 1:
293T (Phosphatase treated membrane) at 10 µg


Lane 2:
NIH/3T3 (Phosphatase treated membrane) at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 18 kDa

Observed band size: 18 kDa



Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Western blot staining using rabbit Anti-Cofilin (phospho S3) antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 37 seconds

All lanes:
Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500) at 1/1000 dilution

Lane 1:
HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate (Untreated membrane) at 10 µg

Lane 2:


HeLa(human cervix adenocarcinoma epithelial cell), whole cell lysate (Phosphatase treated membrane) at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 18 kDa

Observed band size: 18 kDa



Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Cofilin (phospho S3) Western blot staining using rabbit Anti-Cofilin (phospho S3) antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 10 seconds

All lanes:
Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500) at 1/1000 dilution

Lane 1:
C6 (rat glial tumor glial cell), whole cell lysate (Untreated membrane) at 10 µg


Lane 2:
C6 (Phosphatase treated membrane) at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 18 kDa

Observed band size: 18 kDa



Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500) at 1/1000 dilution

Lane 1:
NIH/3T3 (mouse embryonic fibroblast), whole cell lysate (Untreated membrane) at 10 µg

Lane 2:
NIH/3T3 (Phosphatase treated membrane) at 10 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 18 kDa

Observed band size: 18 kDa

Exposure time: 3s



Dot Blot - Anti-Cofilin (phospho S3) antibody [EPR24753-28] (ab283500)

Dot blot analysis of Cofilin (phospho S3) using ab283500 at 1/1000 (0.604 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100,000 dilution.
Lane 1: Cofilin-1 (phospho S3) peptide

Lane 2: Cofilin-1 non-phospho peptide

Exposure time: 3 minutes

Blocking and diluting buffer and concentration: 5% NFDM/TBST

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.