abcam

ab283568

Anti-CGRP-I+CGRP-II antibody [EPR23804-95]

Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568) is a rabbit monoclonal antibody detecting CGRP-I+CGRP-II in **Western Blot, Flow Cytometry (Intra), IHC-P, IHC-Fr, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23804-95
Purification technique	Affinity purification Protein A
Concentration	0.55 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/5000

Notes	-

Not recommended

Species	Mouse, Rat, Recombinant full length protein - Mouse
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	-
Notes	-

Species	Rat
Dilution info	-
Notes	-

Species	Human
Dilution info	1/100
Notes	-

Not recommended

Species	Recombinant full length protein - Mouse
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat, Recombinant full length protein - Mouse
Dilution info	-

Notes	-

IHC-P

Tested

Species	Rat
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Not recommended

Species	Human
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Species	Recombinant full length protein - Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Recombinant full length protein - Mouse
Dilution info	1/1000

Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Species	Rat
Dilution info	1/100
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Not recommended

Species	Human
Dilution info	-
Notes	Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).

Species	Recombinant full length protein - Mouse
Dilution info	-
Notes	-

Target data

See full target information Calca

Function	CGRP1/CALCA is a peptide hormone that induces vasodilation mediated by the CALCRL-RAMP1 receptor complex. Dilates a variety of vessels including the coronary, cerebral and systemic vasculature. Its abundance in the CNS also points toward a neurotransmitter or neuromodulator role. It also elevates platelet cAMP. CGRP1 can also bind and activate CALCR-RAMP1 (AMYR1) receptor complex.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P), Immunohistochemistry (IHC-Fr), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

Other related products
We have a range of other formats of antibody clone [EPR23804-95] also available for your convenience: Biotin - ab322855, Alexa Fluor® 647 - ab302831

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The target designated as CGRP-I and CGRP-II refers to calcitonin gene-related peptide isoforms which play key roles in various physiological functions. CGRP around 37 amino acids long has alternative names such as alpha-CGRP and beta-CGRP occurring mainly in the central nervous system and peripheral nervous tissues. These peptides are potent vasodilators integral to the modulation of blood vessel tone and immune responses. Both isoforms though small in mass allow for significant physiological impacts due to their receptor interactions.
Biological function summary	CGRP-I and CGRP-II contribute to vasodilation and pain transmission in the nervous system. Each isoform can activate specific CGRP receptors mainly located on smooth muscle cells and neurons without forming stable complexes. Their activity involves increasing intracellular cAMP levels facilitating vascular relaxation and modulating nociceptive pathways. Consequently CGRP peptides are essential for conveying pain signals and supporting neurogenic inflammation mechanisms.
Pathways	CGRP-I and CGRP-II serve critical roles in the cardiovascular and nervous system pathways. The cAMP-dependent pathway one of the essential routes of action elevates vasodilation responses where CGRP interacts with the RAMP1 protein a receptor accessory protein. Another pathway involves the modulation of pain where CGRP interacts with other neurotransmitters and proteins such as substance P and neurokinin.
Associated diseases and disorders	CGRP-I and CGRP-II show significant associations with migraines and hypertension. These peptides through their vasodilatory capabilities and nociceptive signaling connect to migraines by interacting with proteins like RAMP1 in trigeminal neurons exacerbating migraine pathology. Furthermore CGRP's role in promoting vasodilation links to hypertension management where altered CGRP activity affects blood pressure regulation. These connections make CGRP-related pathways viable targets for therapeutic intervention.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

17 product images




Immunocytochemistry/ Immunofluorescence - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized TT cells labelling CGRP-I+CGRP-II with ab283568 at 1/100 (6.39 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).
Confocal image showing cytoplasmic staining in TT cell line, and no staining in 293T cell line is observed.


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Flow Cytometry (Intracellular) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Flow Cytometry (Intracellular) staining using rabbit Anti-CGRP-I+CGRP-II antibody


Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized 293T (Human embryonic kidney epithelial cell) (Left) / TT (Human thyroid carcinoma epithelial cell) (Right) cells labelling CGRP-I+CGRP-II with ab283568 at 1/5000 dilution (0.01ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat F(ab')2 Anti-Rabbit IgG(DyLight® 488, ab98507) at 1/500 dilution was used as the secondary antibody.
Negative control: 293T.



Immunocytochemistry/ Immunofluorescence - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunocytochemistry/ Immunofluorescence staining of Mouse splenocytes using rabbit Anti-CGRP-I+CGRP-II antibody


Immunofluorescent analysis of 4% Paraformaldehyde fixed 0.1% Triton X-100 permeabilized Mouse splenocytes labelling CGRP-I+CGRP-II with ab283568 at 1/200 (2.9 µg/ml) followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody as secondary at 1/1000 (2 µg/ml) dilution. **Negative control:** confocal image showing no staining in mouse splenocytes (shown in green). The counterstain was observed in magenta. Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunocytochemistry/ Immunofluorescence - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunocytochemistry/ Immunofluorescence staining of Rat DRG primary neurons using rabbit Anti-CGRP-I+CGRP-II antibody

Immunofluorescent analysis of 4% Paraformaldehyde fixed rat DRG primary neurons permeabilized using 0.1% Triton X-100 labelling CGRP-I+CGRP-II with ab283568 (green) followed by ab150081 AlexaFluor®488 preadsorbed Goat anti-Rabbit secondary. ab78078 anti-beta III Tubulin (mouse mAb) was used as counterstain (magenta) with ab150120 AlexaFluor®594 Goat anti-Mouse as secondary. Nuclear DNA was labelled with DAPI (shown in blue). Image showing cytoplasmic staining in rat DRG primary neurons was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of paraffin-embedded Mouse spinal cord tissue labelling CGRP-I+CGRP-II with ab283568 at 1/1000 (0.639 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse central termination sensory neurons (PMID: 29634489). The section was incubated with ab283568 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins




Immunohistochemistry (Frozen sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Frozen sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat spinal cord tissue labeling CGRP-I+CGRP-II with ab283568 at 1/100 (6.39 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution (Green). Positive staining on the dorsal horn of rat spinal cord is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of paraffin-embedded Rat skin tissue labelling CGRP-I+CGRP-II with ab283568 at 1/1000 (0.639 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). The section was incubated with ab283568 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Negative control: no staining on rat skin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins



Immunohistochemistry (Frozen sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Frozen sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse skin tissue labeling CGRP-I+CGRP-II with ab283568 at 1/100 (6.39 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution (Green). The nuclear counterstain was DAPI (Blue).
Negative control: No staining on mouse skin is observed.

Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of paraffin-embedded Rat spinal cord tissue labelling CGRP-I+CGRP-II with ab283568 at 1/1000 (0.639 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on rat central termination sensory neurons. The section was incubated with ab283568 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins




Immunohistochemistry (Frozen sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Frozen sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat skin tissue labeling CGRP-I+CGRP-II with ab283568 at 1/100 (6.39 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution (Green). The nuclear counterstain was DAPI (Blue).
Negative control: No staining on rat skin is observed.

Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labelling CGRP-I+CGRP-II with ab283568 at 1/1000 (0.639 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on the muscularis nerve fibres of mouse colon. The section was incubated with ab283568 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of paraffin-embedded Mouse skin tissue labelling CGRP-I+CGRP-II with ab283568 at 1/1000 (0.639 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). The section was incubated with ab283568 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Negative control: no staining on mouse skin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins



Immunocytochemistry/ Immunofluorescence - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunocytochemistry/ Immunofluorescence staining of Mouse DRG primary neurons using rabbit Anti-CGRP-I+CGRP-II antibody


Immunofluorescent analysis of 4% Paraformaldehyde fixed mouse DRG primary neurons permeabilized using 0.1% Triton X-100 labelling CGRP-I+CGRP-II with ab283568 (green) followed by ab150081 AlexaFluor®488 preadsorbed Goat anti-Rabbit secondary. ab78078 anti-beta III Tubulin (mouse mAb) was used as counterstain (magenta) with ab150120 AlexaFluor®594 Goat anti-Mouse as secondary. Nuclear DNA was labelled with DAPI (shown in blue). Image showing cytoplasmic staining in mouse DRG primary neurons was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunocytochemistry/ Immunofluorescence - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunocytochemistry/ Immunofluorescence staining of Rat splenocytes using rabbit Anti-CGRP-I+CGRP-II antibody

Immunofluorescent analysis of 4% Paraformaldehyde fixed 0.1% Triton X-100 permeabilized Rat splenocytes labelling CGRP-I+CGRP-II with ab283568 at 1/200 (2.9 µg/ml) followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody as secondary at 1/1000 (2 µg/ml) dilution. **Negative control:** confocal image showing no staining in rat splenocytes (shown in green). The counterstain was observed in magenta. Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunohistochemistry (Frozen sections) - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Immunohistochemistry (Frozen sections) staining using rabbit Anti-CGRP-I+CGRP-II antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse spinal cord tissue labeling CGRP-I+CGRP-II with ab283568 at 1/100 (6.39 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution (Green). Positive staining on the dorsal horn of mouse spinal cord is observed. The nuclear counterstain was DAPI (Blue).
Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Western blot - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

CGRP-I+CGRP-II Western blot staining using rabbit Anti-CGRP-I+CGRP-II antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBSTFresh lysate of TT cell line was used in this blot.
Negative control: 293T, Hela, Daudi (HPA database)

Exposure time: 26 seconds

All lanes:
Western blot - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568) at 1/1000 dilution

Lane 1:
TT (human thyroid carcinoma epithelial cell), whole cell lysate at 20 µg

Lane 2:
293T (human embryonic kidney epithelial cell), whole cell lysate at 20 µg


Lane 3:
HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate at 20 µg

Lane 4:
Daudi (human Burkitt's lymphoma lymphoblast), whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 10 kDa



Western blot - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
This antibody has cross-reaction with mouse CALCB.

The rec proteins CALCB and CLACA were made in-house.

CALCB recombinant protein was made in-house and expressed from the E.coli expression system.

CLACA recombinant protein was expressed from a mammalian - HEK-293 expression system.

Exposure time: 1 second

All lanes:
Western blot - Anti-CGRP-I+CGRP-II antibody [EPR23804-95] (ab283568) at 1/1000 dilution

Lane 1:
His-tagged mouse CALCB recombinant protein at 0.01 µg

Lane 2:
Mouse CALCA recombinant protein at 0.01 µg

Secondary

Lanes 1 - 2:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Lanes 1 - 2:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 14 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.