abcam

ab283660

Anti-Id1 antibody [EPR24304-18A] - BSA and Azide free

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR24304-18A
Purification technique	Affinity purification Protein A
Specificity	Rat only suitable for WB, but not ICC or FC application
Concentration	1.009 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human

Dilution info	-
Notes	-

Not recommended

Species	Rat, Mouse
Dilution info	-
Notes	-

IP

Not recommended

Species	Rat, Human, Mouse
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Rat, Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Rat, Human
Dilution info	-

Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

Target data

See full target information ID1

Function	Transcriptional regulator (lacking a basic DNA binding domain) which negatively regulates the basic helix-loop-helix (bHLH) transcription factors by forming heterodimers and inhibiting their DNA binding and transcriptional activity. Implicated in regulating a variety of cellular processes, including cellular growth, senescence, differentiation, apoptosis, angiogenesis, and neoplastic transformation. Inhibits skeletal muscle and cardiac myocyte differentiation. Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-BMAL1 heterodimer (By similarity).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

ab283660 is the carrier-free version of ab283650.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The Id1 protein also known as Inhibitor of Differentiation 1 functions as a helix-loop-helix (HLH) transcription factor. It belongs to the family of inhibitor of differentiation proteins and weighs approximately 16 kDa. It lacks a basic DNA-binding domain which causes it to prevent transcription when it forms heterodimers with other HLH proteins. Id1 shows expression in various tissues but has noticeably high levels in embryonic and adult tissues involved in developmental processes.
Biological function summary	The Id1 protein plays significant roles in cellular differentiation proliferation and senescence. It does not function within a larger complex but acts as a suppressor of basic HLH transcription factors influencing cell differentiation pathways. Particularly it inhibits functions of factors that promote differentiation allowing cells to maintain a proliferative state. This inhibition capability marks Id1 as an important regulator in development and adult tissue maintenance.
Pathways	The Id1 protein influences key processes by interacting with the TGF-β (transforming growth factor-beta) pathway. It helps regulate cellular responses to growth factors that control cell cycle and apoptosis. In addition the protein has connections to the Notch signaling pathway which is important in cell fate determination. Through these pathways Id1 interacts with proteins such as Smad

and Notch receptors highlighting its influence on developmental and immune responses.

Associated diseases and disorders

Id1 has been linked to cancer and neurodegenerative diseases. Its dysregulation often leads to abnormal cell proliferation and resistance to differentiation processes that are hallmarks in cancer progression. Related proteins like c-Myc have interactions with Id1 in cancer endorsing cell cycle advancement. Additionally elevated Id1 expression connects to neurodegenerative conditions as it may disrupt normal neuronal differentiation linking to proteins such as NF-kB involved in cell survival and inflammation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

5 product images




Immunocytochemistry/ Immunofluorescence - Anti-Id1 antibody [EPR24304-18A] - BSA and Azide free (ab283660)

Id1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Id1 antibody

This data was developed using ab283650, the same antibody clone in a different buffer formulation.
Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HCT 116 cells labelling Id1 with ab283650 at 1/50 (9.64 ug/ml) dilution, followed by ab150081 Goat

Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing nuclear and weakly cytoplasmic staining in HCT 116 cell line. is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5 ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Flow Cytometry - Anti-Id1 antibody [EPR24304-18A] - BSA and Azide free (ab283660)

Id1 Flow Cytometry staining using rabbit Anti-Id1 antibody

This data was developed using ab283650, the same antibody clone in a different buffer formulation.
Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HCT 116 (Human colorectal carcinoma epithelial cell) cells labelling Id1 with ab283650 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat F(ab')2 Anti-Rabbit IgG (DyLight® 488, ab98507) at 1/500 dilution was used as the secondary antibody.



Western blot - Anti-Id1 antibody [EPR24304-18A] - BSA and Azide free (ab283660)

Id1 Western blot staining using rabbit Anti-Id1 antibody

This data was developed using ab283650, the same antibody clone in a different buffer formulation.
Blocking and diluting buffer and concentration: 5% NFDM/TBST

The molecular weight observed is consistent with what has been described in the literature (PMID: 11034201, PMID: 1644289).

Exposure time: 3 minutes

All lanes:
Western blot - Anti-Id1 antibody [EPR24304-18A] (ab283650) at 1/1000 dilution

Lane 1:
MDA-MB-231 (human breast adenocarcinoma epithelial cell) whole cell lysate at 40 µg

Lane 2:
PC-3 (human prostate adenocarcinoma epithelial cell) whole cell lysate at 40 µg


Lane 3:
HCT 116 (human colorectal carcinoma epithelial cell) whole cell lysate at 40 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 16 kDa

Observed band size: 14-16 kDa



Western blot - Anti-Id1 antibody [EPR24304-18A] - BSA and Azide free (ab283660)

Id1 Western blot staining using rabbit Anti-Id1 antibody

This data was developed using ab283650, the same antibody clone in a different buffer formulation.
Blocking and diluting buffer and concentration: 5% NFDM/TBST

This blot was developed using a higher sensitivity ECL substrate.

In materials from rats, ab283650 is suitable only for WB application.

Exposure time: 92 seconds

All lanes:
Western blot - Anti-Id1 antibody [EPR24304-18A] (ab283650) at 1/1000 dilution

Lane 1:
Untreated PC-12 (rat adrenal gland pheochromocytoma) whole cell lysate at 20 µg


Lane 2:
PC-12 treated with 50 ng/ml EGF (ab9697) for 48 hours, whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 16 kDa

Observed band size: 14-16 kDa



Western blot - Anti-Id1 antibody [EPR24304-18A] - BSA and Azide free (ab283660)

This data was developed using ab283650, the same antibody clone in a different buffer formulation.
Blocking and diluting buffer and concentration: 5% NFDM/TBST

Exposure time: 8 seconds

All lanes:
Western blot - Anti-Id1 antibody [EPR24304-18A] (ab283650) at 1/1000 dilution

Lane 1:
HEK-293T (human embryonic kidney) transfected with an empty vector (vector control), containing a myc-His-tag®, whole cell lysate at 10 µg

Lane 2:
HEK-293T transfected with human Id1 (WT) expression vector containing a myc-His-tag®, whole cell lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 16 kDa

Observed band size: 14-16 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.