abcam

ab283679

Anti-DAP12 antibody [EPR24244-76]

Anti-DAP12 antibody [EPR24244-76] (ab283679) is a rabbit monoclonal antibody detecting DAP12 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24244-76
Purification technique	Affinity purification Protein A
Concentration	0.48 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Mouse
Dilution info	1/30

Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse

Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

IHC-Fr

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Human
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information Tyrobp

Function

Adapter protein which non-covalently associates with activating receptors found on the surface of a variety of immune cells to mediate signaling and cell activation following ligand binding by the receptors (PubMed:15471863, PubMed:9647200). TYROBP is tyrosine-phosphorylated in the ITAM domain following ligand binding by the associated receptors which leads to activation of additional tyrosine kinases and subsequent cell activation (PubMed:15728241, PubMed:16715077). Also has an inhibitory role in some cells (PubMed:21727189). Non-covalently associates with activating receptors of the CD300 family to mediate cell activation (By similarity). Also mediates cell activation through association with activating receptors of the CD200R family (PubMed:15471863). Required for neutrophil activation mediated by

integrin (PubMed:17086186). Required for the activation of myeloid cells mediated by the CLEC5A/MDL1 receptor (By similarity). Associates with natural killer (NK) cell receptors such as the KLRD1/KLRC2 heterodimer to mediate NK cell activation (By similarity). Also associates non-covalently with the NK cell receptors KLRA4/LY49D and KLRA8/LY49H which leads to NK cell activation (PubMed:9647200). Associates with TREM1 to mediate activation of neutrophils and monocytes (By similarity). Associates with TREM2 on monocyte-derived dendritic cells to mediate up-regulation of chemokine receptor CCR7 and dendritic cell maturation and survival (By similarity). Association with TREM2 mediates cytokine-induced formation of multinucleated giant cells which are formed by the fusion of macrophages (PubMed:18957693). Stabilizes the TREM2 C-terminal fragment (TREM2-CTF) which is produced by TREM2 ectodomain shedding (By similarity). In microglia, required with TREM2 for phagocytosis of apoptotic neurons (PubMed:15728241). Required with ITGAM/CD11B in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development (PubMed:18685038). Promotes pro-inflammatory responses in microglia following nerve injury which accelerates degeneration of injured neurons (PubMed:25690660). Positively regulates the expression of the IRAK3/IRAK-M kinase and IL10 production by liver dendritic cells and inhibits their T cell allostimulatory ability (PubMed:21257958). Negatively regulates B cell proliferation (PubMed:21727189). Required for CSF1-mediated osteoclast cytoskeletal organization (PubMed:18691974). Positively regulates multinucleation during osteoclast development (PubMed:12569157, PubMed:14969392).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-DAP12 antibody [EPR24244-76] (ab283679) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Mouse, Rat samples.

What is the molecular weight of DAP12?

Anti-DAP12 [EPR24244-76] (ab283679) specifically detects a band for DAP12 (UniProt: O54885) at a molecular weight of 12kDa.

Other related products
We have a range of other formats of antibody clone [EPR24244-76] also available for your convenience: ab283679, Carrier free - ab283703, Alexa Fluor® 647 - ab315262, Alexa Fluor® 488 - ab315294

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:


10 product images



Flow Cytometry (Intracellular) - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Flow Cytometry (Intracellular) staining using rabbit Anti-DAP12 antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized Mouse blood cells cells labelling DAP12 with ab283679 at 1/50 dilution (1ug) (Right) compared with a Rabbit monoclonal IgG (ab172730) (Left) isotype control. A Goat F(ab')2 Anti-Rabbit IgG(DyLight® 488, ab98507) at 1/500 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DAP12 antibody

Immunohistochemical analysis of paraffin-embedded Rat liver tissue labelling DAP12 with ab283679 at 1/2000 (0.232 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on kupffer cells of rat liver. The section was incubated with ab283679 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DAP12 antibody

Immunohistochemical analysis of paraffin-embedded Rat lung tissue labelling DAP12 with ab283679 at 1/2000 (0.232 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat lung. The section was incubated with ab283679 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DAP12 antibody

Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labelling DAP12 with ab283679 at 1/2000 (0.232 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse spleen. The section was incubated with ab283679 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-DAP12 antibody

Immunohistochemical analysis of paraffin-embedded Mouse liver tissue labelling DAP12 with ab283679 at 1/2000 (0.232 ug/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on kupffer cells of mouse liver. The section was incubated with ab283679 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-DAP12 antibody

Immunofluorescent analysis of 100% methanol-fixed, 0.1% TritonX-100 permeabilized RAW264.7 cells labelling DAP12 with ab283679 at 1/50 (9.28 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing cytoplasmic and membranous staining in RAW 264.7 cell line. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5 ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).
Negative control: M1 (PMID: 11034384).


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Flow Cytometry (Intracellular) - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Flow Cytometry (Intracellular) staining using rabbit Anti-DAP12 antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized M1 (mouse myeloid leukemia myeloblast) (Left), RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) (Right) cells labelling DAP12 with ab283679 at 1/500 dilution (0.1ug)(Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat F(ab')2 Anti-Rabbit IgG(DyLight® 488, ab98507) at 1/500 dilution was used as the secondary antibody.
Negative control: M1 (PMID: 11034384).



Immunoprecipitation - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 was immunoprecipitated from 0.35 mg RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate 10 ug with ab283679 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab283679 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: RAW264.7 whole cell lysate 10 ug

Lane 2: ab283679 IP in RAW264.7 whole cell lysate


Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab283679 in RAW264.7 whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 5.5 seconds

All lanes:
Immunoprecipitation - Anti-DAP12 antibody [EPR24244-76] (ab283679)

Predicted band size: 12 kDa



Western blot - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Western blot staining using rabbit Anti-DAP12 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Negative control: M1 (PMID: 11034384).

Exposure time: 114 seconds

All lanes:
Western blot - Anti-DAP12 antibody [EPR24244-76] (ab283679) at 1/1000 dilution

Lane 1:
RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage), whole cell lysate at 20 µg


Lane 2:
N9 (mouse microglia), whole cell lysate at 20 µg

Lane 3:
M1 (mouse myeloid leukemia myeloblast), whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 12 kDa



Western blot - Anti-DAP12 antibody [EPR24244-76] (ab283679)

DAP12 Western blot staining of Rat spleen tissue lysate using rabbit Anti-DAP12 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 3 minutes

All lanes:
Western blot - Anti-DAP12 antibody [EPR24244-76] (ab283679) at 1/1000 dilution

All lanes:
Rat spleen tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 12 kDa

Observed band size: 12 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.