abcam

ab284761

Anti-YB1 antibody [EP2706Y] - BSA and Azide free

RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EP2706Y
Purification technique	Affinity purification Protein A

Reactivity data

ICC/IF

Not recommended

Species	Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Target data

See full target information YBX1

Function

DNA- and RNA-binding protein involved in various processes, such as translational repression, RNA stabilization, mRNA splicing, DNA repair and transcription regulation (PubMed:10817758, PubMed:11698476, PubMed:14718551, PubMed:18809583, PubMed:31358969, PubMed:8188694). Predominantly acts as a RNA-binding protein: binds preferentially to the 5'-[CU]CUGCG-3' RNA motif and specifically recognizes mRNA transcripts modified by C5-methylcytosine (m5C) (PubMed:19561594, PubMed:31358969). Promotes mRNA stabilization: acts by binding to m5C-containing mRNAs and recruiting the mRNA stability maintainer ELAVL1, thereby preventing mRNA decay (PubMed:10817758, PubMed:11698476, PubMed:31358969). Component of the CRD-mediated complex that promotes MYC mRNA stability (PubMed:19029303). Contributes to the regulation of translation by modulating the interaction between the mRNA and eukaryotic initiation factors (By similarity). Plays a key role in RNA composition of extracellular exosomes by defining the sorting of small non-coding RNAs, such as tRNAs, Y RNAs, Vault RNAs and miRNAs (PubMed:27559612, PubMed:29073095). Probably sorts RNAs in

exosomes by recognizing and binding C5-methylcytosine (m5C)-containing RNAs (PubMed:28341602, PubMed:29073095). Acts as a key effector of epidermal progenitors by preventing epidermal progenitor senescence: acts by regulating the translation of a senescence-associated subset of cytokine mRNAs, possibly by binding to m5C-containing mRNAs (PubMed:29712925). Also involved in pre-mRNA alternative splicing regulation: binds to splice sites in pre-mRNA and regulates splice site selection (PubMed:12604611). Binds to TSC22D1 transcripts, thereby inhibiting their translation and negatively regulating TGF-beta-mediated transcription of COL1A2 (By similarity). Also able to bind DNA: regulates transcription of the multidrug resistance gene MDR1 is enhanced in presence of the APEX1 acetylated form at 'Lys-6' and 'Lys-7' (PubMed:18809583). Binds to promoters that contain a Y-box (5'-CTGATTGGCCAA-3'), such as MDR1 and HLA class II genes (PubMed:18809583, PubMed:8188694). Promotes separation of DNA strands that contain mismatches or are modified by cisplatin (PubMed:14718551). Has endonucleolytic activity and can introduce nicks or breaks into double-stranded DNA, suggesting a role in DNA repair (PubMed:14718551). The secreted form acts as an extracellular mitogen and stimulates cell migration and proliferation (PubMed:19483673).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

ab284761 is the carrier-free version of ab76540

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Y-box Binding Protein 1 (YB1) also known as YBX1 is a multifunctional protein with a mass of approximately 36 kDa. It belongs to the cold shock protein family that contains a highly conserved cold shock domain. YB1 is broadly expressed across many tissues predominantly found in the cytoplasm and nucleus where it functions as a transcription factor and RNA-binding protein. Through its transcriptional regulatory activity YB1 can bind to Y-box sequences in the promoter regions of genes and modulate their expression.
Biological function summary	YB1 plays vital roles in regulating cell proliferation stress response and differentiation. It does not operate alone; it interacts with various complexes notably ribonucleoprotein complexes. YB1 regulates gene expression at both transcriptional and translational levels impacting cell cycle progression and apoptosis pathways. Its ability to bind RNA and DNA makes it integral in controlling the expression of genes related to stress responses and developmental processes.
Pathways	YB1 is essential in many signaling pathways that govern cell growth and survival including the PI3K/AKT pathway and MAPK pathway. In these pathways it interacts with proteins like AKT1 and MAPK3 influencing cellular responses to external and internal stimuli. These interactions reveal the adaptability of YB1 in various cellular contexts allowing it to mediate pathway-specific responses to environmental cues thereby promoting the maintenance of cellular homeostasis and adaptation.
Associated diseases and disorders	The dysregulation of YB1 has been linked to cancer and neurodegenerative diseases. Overexpression or altered localization of YB1 often in connection with other proteins such as p53 contributes to tumorigenesis by enhancing cell proliferation and inhibiting programmed cell death. In neurodegenerative diseases impaired YB1 function can negatively affect neuronal survival potentially mediated through its interactions with proteins like tau which are implicated in conditions such as Alzheimer's disease. Understanding of these associations highlights the potential of YB1 as a therapeutic target in disease management.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


2 product images



Flow Cytometry (Intracellular) - Anti-YB1 antibody [EP2706Y] - BSA and Azide free (ab284761)

YB1 Flow Cytometry (Intracellular) staining using rabbit Anti-YB1 antibody

This data was developed using ab76540, the same antibody clone in a different buffer formulation.Intracellular flow cytometric analysis of permeabilized HeLa cells using ab76540 at 1/60 dilution (red) or a rabbit IgG (negative) (green)



Western blot - Anti-YB1 antibody [EP2706Y] - BSA and Azide free (ab284761)

This data was developed using ab76540, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-YB1 antibody [EP2706Y] (ab76540) at 1/20000 dilution

Lane 1:
HeLa cell lysate at 10 µg

Lane 2:
293T cell lysate at 10 µg

Lane 3:
Jurkat cell lysate at 10 µg

Lane 4:
MCF7 cell lysate at 10 µg

Secondary

All lanes:
goat anti-rabbit HRP at 1/1000 dilution

Predicted band size: 36 kDa

Observed band size: 47 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.