abcam

ab287961

Anti-Dectin-1 antibody [EPR25360-64]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR25360-64
Purification technique	Affinity purification Protein A
Concentration	0.502 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

WB

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

Target data

See full target information Clec7a

Function	Lectin that functions as a pattern recognizing receptor (PRR) specific for beta-1,3-linked and beta-1,6-linked glucans, which constitute cell wall constituents from pathogenic bacteria and fungi (PubMed:11544516, PubMed:15213161, PubMed:17159984). Necessary for the TLR2-mediated inflammatory response and activation of NF-kappa-B: upon beta-glucan binding, recruits SYK via its ITAM motif and promotes a signaling cascade that activates some CARD domain-BCL10-MALT1 (CBM) signalosomes, leading to the activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:12719479, PubMed:15731053, PubMed:16825490, PubMed:32358020). Enhances cytokine production in macrophages and dendritic cells (PubMed:15845454). Mediates production of reactive oxygen species in the cell (PubMed:12719479, PubMed:15731053, PubMed:16825490). Mediates phagocytosis of C.albicans conidia (PubMed:15729357, PubMed:16825490). Binds T-cells in a way that does not involve their surface glycans and plays a role in T-cell activation (PubMed:10779524). Stimulates T-cell proliferation (PubMed:10779524). Induces phosphorylation of SCIMP after binding beta-glucans (PubMed:27288407).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

3 product images




Flow Cytometry - Anti-Dectin-1 antibody [EPR25360-64] (ab287961)

Dectin-1 Flow Cytometry staining using rabbit Anti-Dectin-1 antibody

Flow cytometric analysis of Mouse blood cells cells labelling Dectin-1 with ab287961 at 1/500 dilution (0.1ug)(Right), compared with a Rabbit monoclonal IgG (ab172730) (Left) isotype control.


Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody. Gated on viable cells.



Flow Cytometry - Anti-Dectin-1 antibody [EPR25360-64] (ab287961)

Dectin-1 Flow Cytometry staining using rabbit Anti-Dectin-1 antibody

Flow cytometric analysis of Mouse blood cells cells labelling Dectin-1 with ab287961 at 1/500 dilution (0.1ug)(Right) compared with a Rabbit monoclonal IgG (ab172730) (Left) isotype control. Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody. Cells were stained with rabbit IgG or ab287961. Then stained with anti-Ly6g conjugated to Pacific blue.Gated on viable cells.



Immunocytochemistry/ Immunofluorescence - Anti-Dectin-1 antibody [EPR25360-64] (ab287961)

Dectin-1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Dectin-1 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse PBMC cells labelling Dectin-1 with ab287961 at 1/500 (1.004 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing positive staining in partly mouse PBMC is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.