abcam

ab288538

Anti-IL-13 antibody [B-B13] - BSA and Azide free

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	pH: 7.3 - 7.5 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Recombinant Full Length Protein corresponding to Human IL13. Database link P35225
Carrier free	Yes
Clone number	B-B13
Purification technique	Affinity purification Protein A
Light chain type	kappa
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from cell culture supernatant

Reactivity data

Flow Cyt

Tested

Species	Human

Dilution info	1/2 - 1/256
Notes	-

Target data

See full target information IL13

Function	Cytokine that plays important roles in allergic inflammation and immune response to parasite infection (PubMed:8096327, PubMed:8097324). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). Stimulates B-cell proliferation, and activation of eosinophils, basophils, and mast cells (PubMed:7903680, PubMed:8759755). Plays an important role in controlling IL33 activity by modulating the production of transmembrane and soluble forms of interleukin-1 receptor-like 1/IL1RL1 (By similarity). Displays the capacity to antagonize Th1-driven proinflammatory immune response and downregulates synthesis of many proinflammatory cytokines including IL1, IL6, IL10, IL12 and TNF through a mechanism that partially involves suppression of NF-kappa-B (By similarity). Also functions on nonhematopoietic cells, including endothelial cells where it induces vascular cell adhesion protein 1/VCAM1, which is important in the recruitment of eosinophils (PubMed:8639787). Exerts its biological effects through its receptors which comprises the IL4R chain and the IL13RA1 chain, to activate JAK1 and TYK2, leading to the activation of STAT6 (PubMed:9013879). Aside from IL13RA1, another receptor IL13RA2 acts as a high affinity decoy for IL13 and mediates internalization and depletion of extracellular IL13 (PubMed:21622864).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

Hybridoma: Myeloma X63/AG.8653 x Balb/c spleen cells.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry - Anti-IL-13 antibody [B-B13] - BSA and Azide free (ab288538)


IL-13 Flow Cytometry staining using mouse Anti-IL-13 antibody

TF1 cells proliferation measurement with resazurin after 4 days with IL-13 (6 ng/mL) and different concentrations of antibody.



Flow Cytometry - Anti-IL-13 antibody [B-B13] - BSA and Azide free (ab288538)

CD4+ activated by PMA and ionomycine then marked with B-B13 on intracytoplasmic staining.



Flow Cytometry - Anti-IL-13 antibody [B-B13] - BSA and Azide free (ab288538)

B-B13 with non-activated CD4+.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.