abcam

ab288715

Anti-S100A8 + S100A9 antibody [RM1038]

Anti-S100A8 + S100A9 antibody [RM1038] (ab288715) is a rabbit recombinant multiclonal antibody detecting S100A8 + S100A9 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, IHC-Fr, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Multiclonal
Immunogens	The exact immunogen used to generate this antibody is proprietary information.
Clone number	RM1038
Purification technique	Affinity purification Protein A
Concentration	0.505 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-Fr

Tested

Species	Mouse

Dilution info	1/100
Notes	-

Expected

Species	Rat, Human
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Rat
Dilution info	1/5000
Notes	-

Species	Human
Dilution info	1/5000
Notes	-

Species	Mouse
Dilution info	1/5000
Notes	-

IP

Tested

Species	Mouse
Dilution info	-
Notes	-

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Species	Mouse
Dilution info	1/500
Notes	-

Expected

Species	Rat

Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Species	Human
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information S100A9

Function	S100A9 is a calcium- and zinc-binding protein which plays a prominent role in the regulation of inflammatory processes and immune response (PubMed:12626582, PubMed:15331440, PubMed:16258195, PubMed:19122197, PubMed:20103766, PubMed:21325622, PubMed:8423249). It can induce neutrophil chemotaxis, adhesion, can increase the bactericidal activity of neutrophils by promoting phagocytosis via activation of SYK, PI3K/AKT, and ERK1/2 and can induce degranulation of neutrophils by a MAPK-dependent mechanism (PubMed:12626582, PubMed:15331440, PubMed:20103766). Predominantly found as calprotectin (S100A8/A9) which has a wide plethora of intra- and extracellular functions (PubMed:16258195, PubMed:19122197, PubMed:8423249). The intracellular functions include: facilitating leukocyte arachidonic acid trafficking and metabolism, modulation of the tubulin-dependent cytoskeleton during migration of phagocytes and activation of the neutrophilic NADPH-oxidase (PubMed:15331440, PubMed:21325622). Also participates in regulatory T-cell differentiation together with CD69 (PubMed:26296369). Activates NADPH-oxidase by facilitating the enzyme complex assembly at the cell membrane, transferring arachidonic acid, an essential cofactor, to the enzyme complex and S100A8

contributes to the enzyme assembly by directly binding to NCF2/P67PHOX (PubMed:15642721, PubMed:22808130). The extracellular functions involve pro-inflammatory, antimicrobial, oxidant-scavenging and apoptosis-inducing activities (PubMed:19534726, PubMed:8423249). Its pro-inflammatory activity includes recruitment of leukocytes, promotion of cytokine and chemokine production, and regulation of leukocyte adhesion and migration (PubMed:15598812, PubMed:21487906). Acts as an alarmin or a danger associated molecular pattern (DAMP) molecule and stimulates innate immune cells via binding to pattern recognition receptors such as Toll-like receptor 4 (TLR4) and receptor for advanced glycation endproducts (AGER) (PubMed:19402754). Binding to TLR4 and AGER activates the MAP-kinase and NF-kappa-B signaling pathways resulting in the amplification of the pro-inflammatory cascade (PubMed:19402754, PubMed:22804476). Has antimicrobial activity towards bacteria and fungi and exerts its antimicrobial activity probably via chelation of Zn(2+) which is essential for microbial growth (PubMed:19087201). Can induce cell death via autophagy and apoptosis and this occurs through the cross-talk of mitochondria and lysosomes via reactive oxygen species (ROS) and the process involves BNIP3 (PubMed:19935772). Can regulate neutrophil number and apoptosis by an anti-apoptotic effect; regulates cell survival via ITGAM/ITGB and TLR4 and a signaling mechanism involving MEK-ERK (PubMed:22363402). Its role as an oxidant scavenger has a protective role in preventing exaggerated tissue damage by scavenging oxidants (PubMed:21912088, PubMed:22489132). Can act as a potent amplifier of inflammation in autoimmunity as well as in cancer development and tumor spread (PubMed:16258195). Has transnitrosylase activity; in oxidatively-modified low-densitity lipoprotein (LDL(ox))-induced S-nitrosylation of GAPDH on 'Cys-247' proposed to transfer the NO moiety from NOS2/iNOS to GAPDH via its own S-nitrosylated Cys-3 (PubMed:25417112). The iNOS-S100A8/A9 transnitrosylase complex is proposed to also direct selective inflammatory stimulus-dependent S-nitrosylation of multiple targets such as ANXA5, EZR, MSN and VIM by recognizing a [IL]-x-C-x-x-[DE] motif (PubMed:25417112).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-S100A8 + S100A9 antibody [RM1038] (ab288715) is a rabbit recombinant recombinant multiclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunohistochemistry (IHC-Fr), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of S100A8 + S100A9?
Anti-S100A8 + S100A9 [RM1038] (ab288715) specifically detects a band for S100A8 + S100A9 (UniProt: P06702) at a molecular weight of 13kDa.

What are recombinant multiclonals?
Recombinant multiclonals are a mixture of recombinant antibodies co-expressed from a library of heavy and light chains. They offer several advantages including:

- The sensitivity of polyclonal antibodies by recognising multiple epitopes
- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

View our range of recombinant multiclonal antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	S100A8 and S100A9 often known together as calprotectin are proteins that form a heterodimer and function as a complex. Each protein has a mass of approximately 10.8 kDa for S100A8 and 13.2 kDa for S100A9. These proteins express mainly in neutrophils and monocytes acting as calcium-binding proteins. Their expression can modulate various cellular functions like migration and inflammation responses.
Biological function summary	Calprotectin serves as an important player in the immune response. This complex possesses antimicrobial properties and contributes to the body's defense by modulating inflammation. It binds to zinc and manganese to deprive pathogens of essential nutrients an action called nutrient bioavailability restriction. In immune cells calprotectin's presence signals the activation and recruitment of further immune components.
Pathways	S100A8 and S100A9 influence the NF-kB signaling pathway which governs immune responses and inflammation. Their function affects other pathways such as MAPK by mediating pro-inflammatory cytokines' production and release. These proteins

	often engage with Toll-like receptors impacting the innate immune system and contributing to the body's defensive mechanisms against pathogens.
Associated diseases and disorders	High levels of calprotectin are associated with inflammatory diseases like rheumatoid arthritis and inflammatory bowel disease. Its involvement in these conditions makes it a valuable biomarker. Calprotectin often interacts with other inflammatory molecules like TNF-alpha an important factor in propagating inflammation. The calprotectin ELISA test kit plasma calprotectin immunoassay and similar tools can aid in diagnosing and monitoring these conditions providing insight into disease progression and response to treatment.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

14 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-S100A8 + S100A9 antibody

Immunohistochemical analysis of paraffin-embedded Human spleen tissue labeling S100A8+S100A9 with ab288715 at 1/5000 (0.096 ug/ml) followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection). Positive staining on the human spleen. The section was

incubated with ab288715 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND™ RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection).


Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins



Immunocytochemistry/ Immunofluorescence - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HL-60 cells labelling S100A8+S100A9 with ab288715 at 1/500 (0.96 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing membranous staining in HL-60 cell line is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-S100A8 + S100A9 antibody


Immunohistochemical analysis of paraffin-embedded Human stomach carcin tissue labeling S100A8+S100A9 with ab288715 at 1/5000 (0.096 ug/ml) followed by a ready to use LeicaDS9800 (Bond® Polymer Refine Detection). Positive staining on the stroma inflammatory cells in human stomach carcinoma. The section was incubated with ab288715 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND™ RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond® Polymer Refine Detection).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins



Flow Cytometry (Intracellular) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

Flow cytometric analysis of 4% paraformaldehyde fixed 0.1% Tween-20 permeabilized HL-60 (Human acute promyelocytic leukemia promyeloblast) cells labelling S100A8+S100A9 with ab288715 at 1/500 dilution (0.1ug)(Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8+S100A9 was immunoprecipitated from 0.35 mg HL-60 (Human acute promyelocytic leukemia promyeloblast) whole cell lysate 10ug with ab288715 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab288715 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HL-60 (Human acute promyelocytic leukemia promyeloblast) whole cell lysate 10ug

Lane 2: ab288715 IP in HL-60 whole cell lysate

Lane 3:Rabbit monoclonal IgG (ab172730) instead of ab288715 in HL-60 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 30 seconds

All lanes:
Immunoprecipitation - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

Predicted band size: 13 kDa

Observed band size: 11 kDa, 14 kDa




Immunocytochemistry/ Immunofluorescence - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunocytochemistry/ Immunofluorescence staining of Mouse PBMC cells using rabbit Anti-S100A8 + S100A9 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized Mouse PBMC cells labelling S100A8+S100A9 with ab288715 at 1/500 (0.96 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing membranous staining in subsets of mouse PBMCs is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.



Flow Cytometry (Intracellular) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Flow Cytometry (Intracellular) staining using rabbit Anti-S100A8 + S100A9 antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 0.1% Tween-20 permeabilized Mouse blood cells cells labelling S100A8+S100A9 with ab288715 at 1/500 dilution (0.1 µg)/ Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control. A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-S100A8 + S100A9 antibody

Immunohistochemical analysis of paraffin-embedded Rat spleen tissue labeling S100A8+S100A9 with ab288715 at 1/5000 (0.096 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on the rat spleen. The section was incubated with ab288715 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-S100A8 + S100A9 antibody

Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling S100A8+S100A9 with ab288715 at 1/5000 (0.096 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on the mouse spleen. The section was incubated with ab288715 for 10 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins



Immunohistochemistry (Frozen sections) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunohistochemistry (Frozen sections) staining using rabbit Anti-S100A8 + S100A9 antibody


Positive staining on the red pulp of mouse spleen.
Fresh mouse spleen was fixed with 4% PFA and parmeabilised with 0.2 % Triton X100. ab288715 was used as a primary antibody at 1/100 dilution. ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) was preadsorbed and used as a sacondary antiobdy at 1/1000 dilution. DAPI was used as a nuclear counter stain.



Immunohistochemistry (Frozen sections) - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Immunohistochemistry (Frozen sections) staining using rabbit Anti-S100A8 + S100A9 antibody

Positive staining on the red pulp of rat spleen.
Fresh rat spleen was fixed with 4% PFA and parmeabilised with 0.2 % Triton X100. ab288715 was used as a primary antibody at 1/100 dilution. ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) was preadsorbed and used as a sacondary antiobdy at 1/1000 dilution. DAPI was used as a nuclear counter stain.



Immunoprecipitation - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8+S100A9 was immunoprecipitated from 0.35 mg Mouse spleen lysate 10ug with ab288715 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab288715 at 1/1000 dilution. VeriBlot for IP secondary antibody (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: Mouse spleen lysate 10ug

Lane 2: ab288715 IP in Mouse spleen lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab288715 in Mouse spleen lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds

All lanes:
Immunoprecipitation - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

Predicted band size: 13 kDa

Observed band size: 11 kDa, 14 kDa



Western blot - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)

S100A8 + S100A9 Western blot staining using rabbit Anti-S100A8 + S100A9 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 3 min

All lanes:
Western blot - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715) at 1/1000 dilution

Lane 1:
Mouse spleen lysate

Lane 2:
Rat spleen lysate

Lane 3:
SK-BR-3 (Human breast adenocarcinoma epithelial cell) whole cell lysate

Lane 4:
HL-60 (Human acute promyelocytic leukemia promyeloblast) whole cell lysate


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 13 kDa

Observed band size: 11 kDa, 14 kDa

Western blot - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715)



S100A8 + S100A9 Western blot staining using rabbit Anti-S100A8 + S100A9 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 3 min

All lanes:
Western blot - Anti-S100A8 + S100A9 antibody [RM1038] (ab288715) at 1/1000 dilution

Lane 1:
Mouse PBMC lysate

Lane 2:
Mouse spleen lysate

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 13 kDa

Observed band size: 11 kDa, 14 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.