abcam

ab289968

Anti-FGFR2 antibody [EPR24075-418]

Anti-FGFR2 antibody [EPR24075-418] (ab289968) is a rabbit monoclonal antibody detecting FGFR2 in **Western Blot, Flow Cytometry, IP, IHC-P, ICC/IF**. Suitable for **Human**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR24075-418
Purification technique	Affinity purification Protein A
Concentration	0.632 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/50
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-

Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Mouse
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information FGFR2

Function	Tyrosine-protein kinase that acts as a cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation, migration and apoptosis, and in the regulation of embryonic development. Required for normal embryonic patterning, trophoblast function, limb bud development, lung morphogenesis, osteogenesis and skin development. Plays an essential role in the regulation of osteoblast differentiation, proliferation and apoptosis, and is required for normal skeleton development. Promotes cell proliferation in keratinocytes and immature osteoblasts, but promotes apoptosis in differentiated osteoblasts. Phosphorylates PLCG1, FRS2 and PAK4. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. FGFR2 signaling is down-regulated by ubiquitination, internalization and degradation. Mutations that lead to constitutive kinase activation or impair normal FGFR2 maturation, internalization and degradation lead to aberrant signaling. Over-expressed FGFR2 promotes activation of STAT1.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-FGFR2 antibody [EPR24075-418] (ab289968) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Flow Cyt), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human samples.

What is the molecular weight of FGFR2?
Anti-FGFR2 [EPR24075-418] (ab289968) specifically detects a band for FGFR2 (UniProt: P21802) at a molecular weight

of 92kDa.

Specificity confirmed
The specificity of Anti-FGFR2 antibody [EPR24075-418] (ab289968) has been confirmed by Western blot testing in FGFR2 Knockout HAP1 cells.

Other related products
We have a range of other formats of antibody clone [EPR24075-418] also available for your convenience: ab289968, Carrier free - ab289993, Alexa Fluor® 488 - ab309606, PE - ab322671

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The fibroblast growth factor receptor 2 (FGFR2) also known as CD332 is a protein that functions as a transmembrane receptor tyrosine kinase. FGFR2 is important in signal transduction where it binds to fibroblast growth factors triggering downstream intracellular pathways. FGFR2 has two main isoforms FGFR2b and FGFR2c created through alternative splicing. The molecular weight of FGFR2 varies by isoform typically around 100-110 kDa. It gets expressed in various tissues including epithelial and mesenchymal tissues.
Biological function summary	FGFR2 plays a role in diverse cellular processes like proliferation differentiation and migration. As part of a receptor-ligand complex FGFR2 partners with fibroblast growth factors to initiate signaling cascades. It contributes to embryonic development tissue repair and angiogenesis. FGFR2b expression predominantly appears in epithelial cells while FGFR2c is found in mesenchymal cells reflecting its distinct role in tissue-specific signaling.
Pathways	FGFR2 integrates into key signaling networks especially the MAPK and PI3K/AKT pathways. It interacts with proteins like FRS2 and GRB2 which facilitate the activation of downstream signals promoting cell survival and growth. FGFR2's engagement in these pathways is critical for maintaining normal cellular functions and contributes to its involvement in regulating complex cellular responses.

Associated diseases and disorders	FGFR2 mutations have connections with certain cancers and craniosynostosis syndromes. Alterations in FGFR2 signaling pathways can lead to aberrant cellular growth and have been found in various cancers including breast and gastric cancer. In craniosynostosis syndromes mutations in FGFR2 affect bone development linking it with disorders like Apert syndrome. These conditions demonstrate the pivotal role of FGFR2 in maintaining cellular homeostasis and the consequences of its dysfunction.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

8 product images




Immunocytochemistry/ Immunofluorescence - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-FGFR2 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized KATO III cells labeling FGFR2 with ab289968 at 1/50 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green). Confocal image showing membrane staining in the KATO III cell line, while no staining was observed in the SW620 cell line. **Negative control:** SW620 (PMID: 24968263). ab195889 Anti-alpha Tubulin mouse monoclonal

antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-FGFR2 antibody

Immunohistochemical analysis of paraffin-embedded Human breast carcinoma tissue labelling FGFR2 with ab289968 at 1/100 dilution, followed by LeicaDS9800 (Bond™, Polymer Refine Detection). The section was incubated with ab289968 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is LeicaDS9800 (Bond™, Polymer Refine Detection).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Flow Cytometry - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 Flow Cytometry staining using rabbit Anti-FGFR2 antibody

Flow cytometric analysis of SW620 (human colorectal adenocarcinoma epithelial cell, Left) / KATO III (human gastric carcinoma, Right) cells labelling FGFR2 with ab289968 at 1/500 dilution (0.1µg) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody. **Negative control:** SW620 (PMID: 24968263). Gated on viable cells.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-FGFR2 antibody

Immunohistochemical analysis of paraffin-embedded Human gastric adenocarcinoma tissue labelling FGFR2 with ab289968 at 1/100 dilution, followed LeicaDS9800 (Bond™, Polymer Refine Detection). Membraneous staining on human gastric adenocarcinoma (PMID: 24518603). The section was incubated with ab289968 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is LeicaDS9800 (Bond™, Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-FGFR2 antibody

Immunohistochemical analysis of paraffin-embedded (A) KATO III cells labeling FGFR2 with ab289968 at 1/100 dilution followed by LeicaDS9800 (Bond™, Polymer Refine Detection). Positive staining on (A) KATO III cells. **Negative control:** no staining on (B) SW620 cells (PMID: 24968263). The section was incubated with ab289968 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is LeicaDS9800 (Bond™, Polymer Refine Detection).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Immunoprecipitation - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 was immunoprecipitated from KATO III (human gastric carcinoma) whole cell lysate with ab289968 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab289968 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: KATO III (human gastric carcinoma) whole cell lysate 10 µg

Lane 2: ab289968 IP in KATO III whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab289968 in KATO III whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 5.5 seconds

All lanes:
Immunoprecipitation - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

Predicted band size: 92 kDa

Observed band size: 145 kDa, 85 kDa



Western blot - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) at 1/1000000 dilution.

This blot was developed using a high sensitivity ECL substrate. The high-sensitivity ECL substrate used allows for the detection of proteins in the mid-femtogram range.

All lanes:
Western blot - Anti-FGFR2 antibody [EPR24075-418] (ab289968) at 1/1000 dilution

Lane 1:
Wild-type HAP1 (human chronic myelogenous leukemia near-haploid cell) whole cell lysate at 20 µg

Lane 2:
FGFR2 KO HAP1 whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Developed using the ECL technique.

Observed band size: 85 kDa, 145 kDa

Exposure time: 80s



Western blot - Anti-FGFR2 antibody [EPR24075-418] (ab289968)

FGFR2 Western blot staining using rabbit Anti-FGFR2 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
The antibody recognizes endogenous levels of total FGFR2 protein and a C-terminal truncated form K-sam (PMID:17505008).

Negative control: SW620 (PMID: 24968263)

Exposure time: 10 seconds

All lanes:
Western blot - Anti-FGFR2 antibody [EPR24075-418] (ab289968) at 1/1000 dilution

Lane 1:
KATO III (human gastric carcinoma) whole cell lysate at 20 µg

Lane 2:
SW620 (human colorectal adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100000 dilution

Predicted band size: 92 kDa

Observed band size: 145 kDa, 85 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.