abcam

ab300486

Anti-MICB antibody [EPR26362-65] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR26362-65
Purification technique	Affinity purification Protein A
Concentration	1.032 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	-

Notes	-

Not recommended

Species	Recombinant full length protein - Human, Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Human
Dilution info	1/500
Notes	-

Not recommended

Species	Recombinant full length protein - Human, Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Not recommended

Species	Recombinant full length protein - Human, Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human, Recombinant full length protein - Human
Dilution info	-
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Mouse, Rat, Recombinant full length protein - Human
Dilution info	-
Notes	-

Target data

See full target information MICB

Function Widely expressed membrane-bound protein which acts as a ligand to stimulate an activating receptor KLRK1/NKG2D, expressed on the surface of essentially all human natural killer (NK), gammadelta T and CD8+ alphabeta T-cells (PubMed:11491531, PubMed:11777960). Up-regulated in stressed conditions, such as viral and bacterial infections or DNA damage response, serves as signal of cellular stress, and engagement of KLRK1/NKG2D by MICA triggers NK-cells resulting in a range of immune effector functions, such as cytotoxicity and cytokine production.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	+4°C
Aliquoting information	Upon delivery aliquot
Storage information	Do Not Freeze

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	MICB also known as MHC class I polypeptide-related sequence B is a protein with a molecular weight of around 42 kDa. It anchors on cell surfaces and is widely expressed in epithelial cells fibroblasts and monocytes. MICB surfaces on stressed cells but remains minimal in normal conditions indicating its involvement in cell stress response. It is an activating ligand for NKG2D receptor on natural killer (NK) cells and certain T cells playing a major role in immune surveillance.
Biological function summary	MICB contributes to immune system functioning by presenting stress-induced antigens to immune cells. It does not belong to a complex but engages in interactions critical for activating cytotoxic responses. By binding to the NKG2D receptor it helps activate NK cells and certain T cell subsets which results in the

	elimination of damaged or transformed cells. This immune activation is important for the surveillance against cancer and infected cells maintaining tissue homeostasis.
Pathways	MICB plays a role in the stress-induced NKG2D signaling pathway and is involved in the immune response pathway. It cooperates with proteins such as NKG2D and MICA another ligand of the NKG2D receptor to initiate immune cell-mediated cytotoxicity. These pathways activate NK cells and CD8+ T cells important for removing stressed or altered cells thereby influencing the body's defense mechanisms.
Associated diseases and disorders	MICB has connections to cancer and autoimmune disorders. Aberrant expression of MICB appears in tumors affecting tumor immune evasion and associates with autoimmune diseases influencing immune tolerance. Dysregulation of MICB activity can affect interactions with proteins like NKG2D altering immune responses and contributing to disease progression. Therefore understanding MICB's role is critical for developing therapeutic interventions targeting immune-related diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


6 product images



Immunocytochemistry/ Immunofluorescence - Anti-MICB antibody [EPR26362-65] - BSA and Azide free (ab300486)

MICB Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MICB antibody


This data was developed using ab300485, the same antibody clone in a different buffer formulation.Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized K-562 (human chronic myelogenous leukemia lymphoblast) cells labelling MICB with ab300485 at 1/50 (9.62ug/mL) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green). Confocal image showing membranous staining in K-562 cell lineNegative control: Daudi (PMID: 19404979) is observed. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Flow Cytometry - Anti-MICB antibody [EPR26362-65] - BSA and Azide free (ab300486)

MICB Flow Cytometry staining using rabbit Anti-MICB antibody

This data was developed using ab300485, the same antibody clone in a different buffer formulation.Flow cytometric analysis of 293T (human embryonic kidney epithelial cell, Left) / Daudi (human Burkitt's lymphoma lymphoblast, Middle) / K562 (human chronic myelogenous leukemia lymphoblast, Right) cells labelling MICB with ab300485 at 1/500 dilution (0.1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody. Gated on viable cells.Negative control: 293T, Daudi (PMID: 28154561).



Immunoprecipitation - Anti-MICB antibody [EPR26362-65] - BSA and Azide free (ab300486)

This data was developed using ab300486, the same antibody clone in a different buffer formulation. MICB was immunoprecipitated from 0.35 mg K562 (human chronic myelogenous leukemia lymphoblast), whole cell lysate 10 ug with ab300485 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab300485 at 1/1000 dilution. VeriBlot for IP secondary antibody (HRP)(ab131366) was used at 1/5000 dilution. Lane 1: K562 (human chronic myelogenous leukemia lymphoblast), whole cell lysate 10 ug Lane 2: ab300485 IP in K562 whole cell lysate Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab300485 in K562 whole cell lysate. Blocking and dilution buffer and concentration: 5% NFDM/TBST. Exposure time: 180 seconds
Lane 1:
Immunoprecipitation - Anti-MICB antibody [EPR26362-65] (ab300485) at 1/30 dilution


Lane 2:
Immunoprecipitation - Anti-MICB antibody [EPR26362-65] (ab300485) at 1/1000 dilution

All lanes:
K562 whole cell lysate at 10 µg

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 180s



Western blot - Anti-MICB antibody [EPR26362-65] - BSA and Azide free (ab300486)

MICB Western blot staining using rabbit Anti-MICB antibody

This data was developed using ab300485, the same antibody clone in a different buffer formulation.

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

This antibody does not cross-react with human MICA.

All lanes:
Western blot - Anti-MICB antibody [EPR26362-65] (ab300485) at 1/1000 dilution

Lane 1:
Purified his-tagged human MICB recombinant protein at 10 ng

Lane 2:
Purified his-tagged human MICA recombinant protein at 10 EU/ng


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 43 kDa

Observed band size: 37-50 kDa

Exposure time: 7.75s



Western blot - Anti-MICB antibody [EPR26362-65] - BSA and Azide free (ab300486)

This data was developed using ab300485, the same antibody clone in a different buffer formulation.
Blocking and diluting buffer and concentration: 5% NFDM/TBST

MICB is a glycoprotein of approximately 61 kDa and detected as two 37-42 kDa bands after treated with Protein Deglycosylation MIX II.

The bands beneath the target band (37 kDa) are reported as deglycosylated cleaved soluble MICB.

Molecular weight observed is consistent with described in the literature (PMID:19147769).

This blot was developed using a high sensitivity ECL substrate. The high-sensitivity ECL substrate used allows for the detection of proteins in the mid-femtogram range.

All lanes:
Western blot - Anti-MICB antibody [EPR26362-65] (ab300485) at 1/1000 dilution

Lane 1:
Untreated K562 (human chronic myelogenous leukemia lymphoblast), whole cell lysate at 15 µg

Lane 2:

K562 whole cell lysate treated with Protein Deglycosylation Mix II at 15 µg

Secondary


All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Developed using the ECL technique.

Predicted band size: 43 kDa

Observed band size: 37-61 kDa

Exposure time: 103s



Western blot - Anti-MICB antibody [EPR26362-65] - BSA and Azide free (ab300486)

This data was developed using ab300485, the same antibody clone in a different buffer formulation.
Blocking and diluting buffer and concentration: 5% NFDM/TBST

The expression profile/molecular weight observed is consistent with what has been described in the literature (PMID:10359807).

Negative control: Daudi, 293T (PMID: 28154561).

This blot was developed using a high sensitivity ECL substrate. The high-sensitivity ECL substrate used allows for the detection of proteins in the mid-femtogram range.

All lanes:
Western blot - Anti-MICB antibody [EPR26362-65] (ab300485) at 1/1000 dilution

Lane 1:
K-562 (human chronic myelogenous leukemia lymphoblast) whole cell lysate at 40 µg

Lane 2:
U266B1 (human multiple myeloma B lymphocyte) whole cell lysate at 40 µg

Lane 3:
Daudi (human Burkitt's lymphoma lymphoblast) whole cell lysate at 40 µg

Lane 4:
293T (human embryonic kidney epithelial cell) whole cell lysate at 40 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051)

Developed using the ECL technique.

Predicted band size: 43 kDa

Observed band size: 37-61 kDa

Exposure time: 48s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.