abcam

ab303459

Alexa Fluor® 647 Anti-CD81 antibody [EPR21916]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 647
Excitation/Emission	Ex: 650nm, Em: 665nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR21916
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human

Dilution info	1/100
Notes	-

ICC/IF

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information CD81

Function

Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling. Essential for trafficking and compartmentalization of CD19 receptor on the surface of activated B cells (PubMed:16449649, PubMed:20237408, PubMed:27881302). Upon initial encounter with microbial pathogens, enables the assembly of CD19-CR2/CD21 and B cell receptor (BCR) complexes at signaling TERMs, lowering the threshold dose of antigen required to trigger B cell clonal expansion and antibody production (PubMed:15161911, PubMed:20237408). In T cells, facilitates the localization of CD247/CD3 zeta at antigen-induced synapses with B cells, providing for costimulation and polarization toward T helper type 2 phenotype (PubMed:22307619, PubMed:23858057, PubMed:8766544). Present in MHC class II compartments, may also play a role in antigen presentation (PubMed:8409388, PubMed:8766544). Can act both as positive and negative regulator of homotypic or heterotypic cell-cell fusion processes. Positively regulates sperm-egg fusion and may be involved in acrosome reaction (By similarity). In myoblasts, associates with CD9 and PTGFRN and inhibits myotube fusion during muscle regeneration (By similarity). In macrophages, associates with CD9 and beta-1 and beta-2 integrins, and prevents macrophage fusion into multinucleated giant cells specialized in ingesting complement-opsonized large particles (PubMed:12796480). Also prevents the fusion of mononuclear cell progenitors into osteoclasts in charge of bone

resorption (By similarity). May regulate the compartmentalization of enzymatic activities. In T cells, defines the subcellular localization of dNTPase SAMHD1 and permits its degradation by the proteasome, thereby controlling intracellular dNTP levels (PubMed:28871089). Also involved in cell adhesion and motility. Positively regulates integrin-mediated adhesion of macrophages, particularly relevant for the inflammatory response in the lung (By similarity). (Microbial infection) Acts as a receptor for hepatitis C virus (HCV) in hepatocytes. Association with CLDN1 and the CLDN1-CD81 receptor complex is essential for HCV entry into host cell. (Microbial infection) Involved in SAMHD1-dependent restriction of HIV-1 replication. May support early replication of both R5- and X4-tropic HIV-1 viruses in T cells, likely via proteasome-dependent degradation of SAMHD1. (Microbial infection) Specifically required for Plasmodium falciparum infectivity of hepatocytes, controlling sporozoite entry into hepatocytes via the parasitophorous vacuole and subsequent parasite differentiation to exoerythrocytic forms.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies

Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD81 also known as TAPA-1 or 26 kDa protein is a member of the tetraspanin family featuring four transmembrane domains. The molecular weight of CD81 is approximately 26-28 kDa. This protein exhibits expression in numerous cell types such as leukocytes endothelial cells and epithelial tissues. CD81 plays a mechanical role by facilitating membrane protein interactions and contributing to cellular processes like adhesion and morphogenesis. Researchers have identified specific isoforms like M38 and labels like 1D6 CHAN in studies involving this target often analyzing CD81 through techniques like Western blot to determine expression levels and molecular weight.
Biological function summary	CD81 interacts with other tetraspanins and forms complexes within the membrane to regulate cellular signaling trafficking and adhesion. It participates in the assembly of larger tetraspanin-enriched microdomains which are important for efficient signaling and functional variety. These complexes modulate cell morphology proliferation and development influencing immune responses and pathogen entry to cells. The presence of CD81 in diverse tissues implies its involvement in a wide range of cellular processes forming essential complexes with proteins like integrins that further engage in tissue repair and immunological defense.
Pathways	CD81 plays significant roles in the immune system and viral entry pathways. It interacts with other proteins like CD9 and CD19 within the immune response pathways regulating lymphocyte activation and differentiation. CD81 is notably a coreceptor in the hepatitis C virus (HCV) entry pathway facilitating viral attachment and fusion into host cells. These interactions illustrate CD81's involvement in modulation of immune cell responses and influence on pathogen infection processes integrating into the complex cellular pathways important for maintaining homeostasis and response to external stimuli.
Associated diseases and disorders	CD81 has connections to hepatitis C virus (HCV) infections and immunological disorders such as systemic lupus erythematosus (SLE). During HCV infection CD81 serves as an important entry point for the virus interacting with proteins like claudin-1 and scavenger receptor class B type I (SR-BI) enabling viral entry and replication within liver cells. In SLE alterations in CD81 expression can impact autoantibody production and lymphocyte behavior contributing to the disease's pathology. Understanding CD81's role in these diseases provides insight into

therapeutic targets and potential interventions for managing infections and autoimmune responses.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


2 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-CD81 antibody [EPR21916] (ab303459)

CD81 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD81 antibody

Immunohistochemical analysis of paraffin-embedded human liver tissue stained for CD81 using ab303459 at a 1/100 dilution (5.0 µg/ml) (Red). Positive staining on human liver. The section was incubated with ab303459 for 60 mins at room temperature (shown in red). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®. The immunostaining was performed on a Leica Biosystems BOND RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 40 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-CD81 antibody [EPR21916] (ab303459)

CD81 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD81 antibody

Immunohistochemical analysis of paraffin-embedded human tonsil tissue stained for CD81 using ab303459 at a 1/100 dilution (5.0 µg/ml) (Red). Positive staining on human tonsil. The section was incubated with ab303459 for 60 mins at room temperature (shown in red). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®. The immunostaining was performed on a Leica Biosystems BOND RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 40 mins.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.