abcam

ab303568

Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free

Recombinant | RabMAb | Advanced Validation

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Multiclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	RM1046
Purification technique	Affinity purification Protein A

Reactivity data

Dot

Tested

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

Expected

Species	Human, Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Not recommended

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	-
Notes	Unsuitable for human samples

Species	Rat
Dilution info	-
Notes	Unsuitable for human samples

Not recommended

Species	Human
Dilution info	-
Notes	Unsuitable for human samples

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse, Rat
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human, Mouse
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.

Notes	-

Not recommended

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human, Mouse
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

PepArr

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide - Human
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat, Synthetic peptide - Human
Dilution info	-
Notes	-

ChIP

Tested

Species	Human
Dilution info	-
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide - Human

Dilution info	-
Notes	-

Target data

See full target information H3C1 phospho S10 + S28

Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

What are recombinant multiclonals?
Recombinant multiclonals are a mixture of recombinant antibodies co-expressed from a library of heavy and light chains. They offer several advantages including:

- The sensitivity of polyclonal antibodies by recognising multiple epitopes
- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

View our range of recombinant multiclonal antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


11 product images



Immunocytochemistry/ Immunofluorescence - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)


This data was developed using ab303567, the same antibody clone in a different buffer formulation. Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HeLa (human cervical adenocarcinoma epithelial cell) cells labelling Histone H3 (phospho S10 + S28) with ab303567 at 1/4000 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green). Confocal image showing

nuclear staining in HeLa in M phase. The image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution.




Flow Cytometry (Intracellular) - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)


This data was developed using ab303567, the same antibody clone in a different buffer formulation. Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HeLa (human cervix adenocarcinoma epithelial cell) cells labelling Histone H3 (phospho S10 + S28) with ab303567 at 1/500 dilution (0.1ug)/ Right (Red) compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)




Immunohistochemistry (Frozen sections) - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)




Immunocytochemistry/ Immunofluorescence - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)

This data was developed using ab303567, the same antibody clone in a different buffer formulation. Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling Histone H3 (phospho S10 + S28) with ab303567 at 1/4000 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green). Confocal image showing nuclear staining in NIH/3T3 in M phase. The image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)



Flow Cytometry (Intracellular) - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)

This data was developed using ab303567, the same antibody clone in a different buffer formulation. Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NIH/3T3 (mouse embryonic fibroblast) cells labelling Histone H3 (phospho S10 + S28) with ab303567 at 1/500 dilution (0.1ug)/ Right (Red) compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody.



Western blot - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)

This data was developed using ab303567, the same antibody clone in a different buffer formulation. Blocking and diluting buffer and concentration:5% NFDM/TBST Exposure time: 48 seconds
All lanes:
Western blot - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] (ab303567) at 1/1000 dilution

Lane 1:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg


Lane 2:
HeLa treated with /ml Nocodazole for 24h whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 17 kDa


Exposure time: 48s



Peptide Array - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)

This data was developed using ab303567, the same antibody clone in a different buffer formulation. Peptide array analysis of ab303567 at 0.1ug/ml (0.1ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Fluo 647nm conjugated at 1:50000 dilution. All batches of ab303567 are tested in Peptide Array against 501 different modified and unmodified histone peptides; each peptide is printed on the array at six concentrations (each in triplicate). Circle area represents affinity between the antibody and a peptide: all antigen-containing peptides are displayed as red circles,

all other peptides as blue circles. The affinity is calculated as the area under the curve when antibody binding values are plotted against the corresponding peptide concentration. Each circle area is normalized to the peptide with the strongest affinity.
The complete dataset, including full list of all peptides and information on the position of each peptide in the diagram, is available under the Product Protocol section.



Western blot - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)

This data was developed using ab303567, the same antibody clone in a different buffer formulation. Blocking and diluting buffer and concentration:5% NFDM/TBST Exposure time: 15 seconds
All lanes:
Western blot - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] (ab303567) at 1/1000 dilution

Lane 1:
NIH/3T3 (mouse embryonic fibroblast) whole cell lysate at 20 µg

Lane 2:
NIH/3T3 treated with 100nM Calycin A for 30min whole cell lysate at 20 µg

Lane 3:
C6 (rat glial tumor glial cell) whole cell lysate at 20 µg


Lane 4:
C6 treated with /ml Calyculin A for 60min whole cell lysate 20ug at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 17 kDa

Exposure time: 15s



Dot Blot - Anti-Histone H3 (phospho S10 + S28) antibody [RM1046] - BSA and Azide free (ab303568)

This data was developed using ab303567, the same antibody clone in a different buffer formulation. Dot blot analysis of Histone H3 (phospho S10 + S28) using ab303567 at 1: 1000 dilution (0.496 µg/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1:10,000 dilution. Lane 1: Histone H3 (phospho S10) peptide Lane 2: Histone H3 (phospho S28) peptide Lane 3: Histone H3 unmodified peptide Exposure time: 10 seconds Blocking and diluting buffer and concentration: 5% NFDM/TBST

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.