abcam

ab305111

Alexa Fluor® 488 Anti-Eph receptor A2 antibody [EPR17660-120]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR17660-120
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Mouse
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information Epha2

Function	Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin-A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells (PubMed:29749928). Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 signaling pathway. May also participate in UV radiation-induced apoptosis and have a ligand-independent stimulatory effect on chemotactic cell migration. During development, may function in distinctive aspects of pattern formation and subsequently in development of several fetal tissues. Involved for

instance in angiogenesis, in early hindbrain development and epithelial proliferation and branching morphogenesis during mammary gland development. Engaged by the ligand ephrin-A5/EFNA5 may regulate lens fiber cells shape and interactions and be important for lens transparency development and maintenance. With ephrin-A2/EFNA2 may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are

sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Eph receptor A2 also known as EphA2 or A2 receptor is a type of protein that functions as a receptor tyrosine kinase. It has an approximate mass of 130 kDa. Researchers find this receptor widely expressed in various tissues including epithelial cells and some neuronal populations. EphA2 is involved in cell positioning movement and organization by interacting with ephrin ligands notably ephrin A1. The receptor-ligand interaction triggers bidirectional signaling that affects cell structure and function.
Biological function summary	Eph receptor A2 participates in important cellular processes such as angiogenesis cell proliferation and migration. It often functions as part of larger signaling networks or complexes that include other Eph receptors and ephrin ligands. These interactions help regulate cell adhesion and detachment which are important for embryonic development and tissue organization. EphA2 studies often use fluorescent labeling strategies such as Alexa 588 to track its expression and activity in cells.
Pathways	Eph receptor A2 integrates into the PI3K/AKT and MAPK signaling pathways which are important for cell survival growth and differentiation. It interacts with proteins like Ras and ERK within these pathways to modulate their functions. EphA2 signaling influences cytoskeletal dynamics and morphogenesis making it an important component in maintaining tissue architecture. Research also indicates a connection between EphA2 and the APC protein highlighting its role in the Wnt signaling pathway.
Associated diseases and disorders	The Eph receptor A2 shows significant involvement in cancer and inflammatory conditions. Overexpression or dysregulation of EphA2 occurs frequently in various cancers including breast and prostate cancer correlating with increased tumor aggressiveness and metastasis. The receptor also interacts with other proteins such as APC which links to tumorigenesis and cancer progression. In immunology EphA2 plays a part in chronic inflammatory diseases by modulating immune cell migration and vascular inflammation. Understanding these connections provides insights for potential therapeutic targets using molecular interventions like A2 antibodies and APC-based strategies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - Alexa Fluor® 488 Anti-Eph receptor A2 antibody [EPR17660-120] (ab305111)

Flow cytometric analysis of L929 (mouse connective tissue fibroblast) cells labelling Eph receptor A2 with ab305111 at 1/50 dilution (1ug) (Red) compared with a Rabbit IgG monoclonal [EPR25A] - Isotype Control (Alexa Fluor® 488) (ab199091) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Gated on viable cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.