abcam

ab307601

Anti-xCT antibody [EPR27115-64]

Anti-xCT antibody [EPR27115-64] (ab307601) is a rabbit monoclonal antibody detecting xCT in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR27115-64
Purification technique	Affinity purification Protein A
Specificity	We recommend not to boil your lysates before loading to gel when preparing membrane used for western blot. Boiling leads to the formation of aggregates of membrane proteins, leading to unsuccessful migration in gel. Please do not boil your lysate when testing membrane protein. FURTHER INFORMATION ON SPECIFICITY (Chinese Version) available under the product protocols section. This file includes key technical notes of experience when using this product.
Concentration	0.544 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Mouse
Dilution info	1/500 - 1/1000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000

Notes	-

Target data

See full target information Cystine/glutamate transporter

Function	Heterodimer with SLC3A2, that functions as an antiporter by mediating the exchange of extracellular anionic L-cystine and intracellular L-glutamate across the cellular plasma membrane (PubMed:11133847, PubMed:11417227, PubMed:14722095, PubMed:15151999, PubMed:34880232, PubMed:35245456, PubMed:35352032). Provides L-cystine for the maintenance of the redox balance between extracellular L-cystine and L-cysteine and for the maintenance of the intracellular levels of glutathione that is essential for cells protection from oxidative stress (By similarity). The transport is sodium-independent, electroneutral with a stoichiometry of 1:1, and is drove by the high intracellular concentration of L-glutamate and the intracellular reduction of L-cystine (PubMed:11133847, PubMed:11417227). Acts as an inhibitor of ferroptosis by mediating the import of L-kynurenine leading to anti-ferroptotic signaling propagation required to maintain L-cystine and glutathione homeostasis (PubMed:35245456, PubMed:40246981). Also inhibits ferroptosis by acting as an atypical proton transporter that mediates a slow proton efflux from lysosomes via cystine and glutamate flux (PubMed:40280132). Glutamate and cystine contain side-chain groups that are protonatable in the physiological range of lysosomal pH and cytosolic pH, respectively, enabling a slow lysosomal proton leak through a substrate-as-proton mechanism (PubMed:40280132). Moreover, mediates N-acetyl-L-cysteine uptake into the placenta leading to subsequently down-regulation of pathways associated with oxidative stress, inflammation and apoptosis (PubMed:34120018). In vitro can also transport L-aspartate (PubMed:11417227). May participate in astrocyte and meningeal cell proliferation during development and can provide neuroprotection by promoting glutathione synthesis and delivery from non-neuronal cells such as astrocytes and meningeal cells to immature neurons (By similarity). Controls the production of pheomelanin pigment directly (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot

Storage information	Avoid freeze / thaw cycle

Notes

Anti-xCT antibody [EPR27115-64] (ab307601) was developed by Abcam using patented rabbit monoclonal antibody technology and is validated for use in Flow Cyt (Intra), ICC/IF, IHC-P, IP and WB.

Abcam's high quality manufacturing and validation processes ensure Anti-xCT antibody [EPR27115-64] (ab307601) has high sensitivity and specificity alongside high lot-to-lot consistency and reproducibility.

Anti-xCT antibody [EPR27115-64] (ab307601) specifically detects xCT (UniProt ID: Q9UPY5; Molecular weight: 56kDa) and is sold in a convenient trial size to enable initial testing (20 µL) and larger sizes for subsequent scaling up experiments (100 µL and 1 mL).

Conjugation-ready, carrier free format available for antibody clone EPR27115-64 - ab307602.

A highly validated xCT (SLC7A11) antibody, essential for studying the cystine/glutamate antiporter SLC7A11, ferroptosis and redox balance. This antibody is crucial in cancer research, particularly in understanding xCT's role in tumor progression and antioxidant defense and is widely used in studies of reactive oxygen species (ROS) and xCT inhibitors for therapeutic applications. SLC7A11 is frequently overexpressed in human cancers. SLC7A11 promotes cystine uptake and glutathione biosynthesis, resulting in protection from oxidative stress and ferroptotic cell death. Recent studies have unexpectedly revealed that SLC7A11 also plays critical roles in glutamine metabolism and regulates the glucose and glutamine dependency of cancer cells. Expression of SLC7A11/xCT correlates with advanced lung cancer stage and predicted a worse 5-year survival.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

18 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)

Immunohistochemical analysis of paraffin-embedded human thyroid carcinoma tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human thyroid carcinoma.


The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection. Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-xCT antibody [EPR27115-64] (ab307601)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized A549 (human lung carcinoma epithelial cell) cells labeling xCT with ab307601 at 1/500 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green). Confocal image showing membranous and cytoplasmic staining in A549 cell line. Low expression: COLO 205 (PMID: 31000598). Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (2.5 µg/ml) (Red). Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)


Immunohistochemical analysis of paraffin-embedded human cerebrum tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human cerebrum. The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)


XCT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human skeletal muscle using rabbit Anti-xCT antibody

Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on human skeletal muscle. The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Flow Cytometry (Intracellular) - Anti-xCT antibody [EPR27115-64] (ab307601)

Flow cytometric analysis of 4% paraformaldehyde-fixed 90% methanol-permeabilized COLO 205 (human colon adenocarcinoma tumor ascitic epithelial cell) (Left panel) / A549 (human lung carcinoma epithelial cell) (Right panel) cells labeling xCT with ab307601 at 1/50 dilution (1µg) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as the secondary antibody. Low expression: COLO 205 (PMID: 31000598).



Immunoprecipitation - Anti-xCT antibody [EPR27115-64] (ab307601)

xCT was immunoprecipitated from 0.35 mg A549 (human lung carcinoma epithelial cell) whole cell lysate 10 µg with ab307601 at 1/30 dilution (2 µg in 0.3 5mg lysates). Western blot was performed on the immunoprecipitate using ab307601 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution. Lane 1: A549 whole cell lysate 10 µg Lane 2: ab307601 IP in A549 whole cell lysate Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab307601 in A549 whole cell lysate Blocking and dilution buffer and concentration: 5% NFDM/TBST. Exposure time: 41 seconds.
All lanes:
Immunoprecipitation - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/30 dilution


All lanes:
A549 (human lung carcinoma epithelial cell)

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution


Observed band size: 37 kDa

Exposure time: 41s



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)

Immunohistochemical analysis of paraffin-embedded rat cerebrum tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat cerebrum. The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)


XCT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-xCT antibody

Immunohistochemical analysis of paraffin-embedded (A) Cerebrum tissue from wild-type C57BL/6J mice and (B) Cerebrum tissue from xCT knockout mice tissue labeling xCT with ab307601 at 1/1000 dilution.
Positive staining on (A) Cerebrum tissue from wild-type C57BL/6J mice, no staining on (B) Cerebrum tissue from xCT knockout mice.

The primary antibody was incubated for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.


The tissue samples were kindly provided by GemPharmatech. C57BL/6JGpt wildtype mice and Slc7a11-KO homozygous mice (Strain ID: T015610).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)

XCT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Rat skeletal muscle using rabbit Anti-xCT antibody


Immunohistochemical analysis of paraffin-embedded rat skeletal muscle tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on rat skeletal muscle (PMID:17401668). The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)


XCT Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse skeletal muscle using rabbit Anti-xCT antibody

Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on mouse skeletal muscle. The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-xCT antibody [EPR27115-64] (ab307601)

Immunohistochemical analysis of paraffin-embedded mouse cerebrum tissue labeling xCT with ab307601 at 1/500 dilution (1.024 µg/ml) followed by ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse cerebrum (PMID:29350434). The section was incubated with ab307601 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

Western blot: Anti-SLC7A11 antibody [EPR27115-64] (ab307601) staining at 1/1000 dilution, shown in black; Mouse anti-CANX [CANX/1543] (ab238078) loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 35 kDa in wild-type HepG2 cell lysates with a reduction in signal observed at this size in SLC7A11 heterozygous knockout cell line. To generate this image, wild-type and SLC7A11 heterozygous knockout HepG2 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times before development with a high-sensitivity ECL substrate kit and imaged with 16 minutes exposure time. Secondary antibodies used were Goat anti-Rabbit HRP (H+L) and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution

Lane 1:
Wild-type HepG2 Vehicle Control Diethyl Maleate (0mM, 24h) cell lysate at 20 µg

Lane 2:
Wild-type HepG2 Treated Diethyl Maleate (0.1mM, 24h) cell lysate at 20 µg

Lane 3:
SLC7A11 heterozygous knockout HepG2 Vehicle Control Diethyl Maleate (0mM, 24h) cell lysate at 20 µg


Lane 4:
SLC7A11 heterozygous knockout HepG2 Treated Diethyl Maleate (0.1mM, 24h) cell lysate at 20 µg

Secondary

All lanes:
Goat anti-Rabbit HRP (H+L) and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 35 kDa



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

Blocking buffer and concentration: 5% NFDM/TBST. **Diluting buffer and concentration:** 5% NFDM/TBST.
Boiling leads to the formation of aggregates of membrane proteins, leading to unsuccessful migration in gel. Please do not boil your lysate when testing membrane protein.

No xCT expression or very low levels were found in lung, heart, liver, and kidney. (PMID: 22667998).

The molecular weight observed is consistent with what has been described in the literature (PMID: 26494316, 18806690, 25384799, 30425240, 29693305).

ab181602 was used as a loading control.

All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution

Lanes 1 and 6:
HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Lanes 2 and 7:
Mouse cerebellum tissue lysate at 20 µg

Lanes 3 and 8:
Mouse colon tissue lysate at 20 µg

Lanes 4 and 9:
Mouse kidney tissue lysate at 20 µg

Lanes 5 and 10:
Mouse heart tissue lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 55 kDa

Observed band size: 37 kDa

Exposure time: 140s



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

Blocking and dilution buffer: 5% NFDM/TBST.

xCT protein is easy to form aggregates generating strong signal at the top.

All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution

Lanes 1 and 4:
293T cells transfected with an empty vector containing a His tag whole cell lysate at 20 µg

Lanes 2 and 5:
293T cells transfected with a Human SLC7A5 expression vector containing a His-tag whole cell lysate at 20 µg

Lanes 3 and 6:
293T cells transfected with a Human xCT expression vector containing a His-tag whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 55 kDa

Observed band size: 37 kDa

Exposure time: 7s



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

Blocking and diluting buffer and concentration: 5% NFDM/TBST. Low expression cell line: COLO 205 (PMID: 31000598). The molecular weight observed is consistent with what has been described in the literature (PMID: 26494316, 18806690, 25384799, 30425240, 29693305).
All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution

Lane 1:
Human cerebellum tissue lysate at 20 µg

Lane 2:
A549 (human lung carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
COLO 205 (human colon adenocarcinoma tumor ascitic epithelial cell) whole cell lysate at 20 µg


Lane 4:
HepG2 (human hepatocellular carcinoma epithelial cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 37 kDa

Exposure time: 70s



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

Western blot: Anti-SLC7A11 antibody [EPR27115-64] (ab307601) staining at 1/1000 dilution, shown in black; Mouse anti-CANX [CANX/1543] (ab238078) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab307601 was shown to bind specifically to SLC7A11. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times before development with a high-sensitivity ECL substrate kit and imaged with 30 seconds exposure time. Secondary antibodies used were Goat anti-Rabbit HRP (H+L) and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution


All lanes:
Recombinant Human xCT protein cell lysate at 0.01 µg

Secondary

All lanes:
Goat anti-Rabbit HRP (H+L) and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Observed band size: 35 kDa



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

Negative control: skeletal muscle (PMID:17401668).

The molecular weight observed is consistent with what has been described in the literature (PMID: 26494316, 18806690, 25384799, 30425240, 29693305).

All lysates were not boiled. For getting better result, we recommend not boiling the sample
All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution

Lane 1:
Mouse cerebellum tissue lysate at 20 µg

Lane 2:
Mouse skeletal muscle tissue lysate at 20 µg


Lane 3:
Rat cerebellum tissue lysate at 20 µg

Lane 4:
Rat skeletal muscle tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Exposure time: 180s



Western blot - Anti-xCT antibody [EPR27115-64] (ab307601)

XCT Western blot staining using rabbit Anti-xCT antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
In Western blot, Anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution.

All lysates were not boiled. For getting better result we recommend not boiling the sample. The tissue samples were kindly provided by GemPharmatech. C57BL/6JGpt wildtype mice and Slc7a11-KO homozygous mice (Strain ID: T015610).

All lanes:
Western blot - Anti-xCT antibody [EPR27115-64] (ab307601) at 1/1000 dilution

Lane 1:
Wild-type mouse brain tissue lysate (male) at 40 µg

Lane 2:
Slc7a11 knockout mouse brain tissue lysate (male case1) at 40 µg

Lane 3:
Slc7a11 knockout mouse brain tissue lysate (male case2) at 40 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 35 kDa

Exposure time: 26s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.