abcam

ab312324

Anti-BASP1 antibody [EPR28066-51]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR28066-51
Purification technique	Affinity purification Protein A
Concentration	0.502 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	0.502 ug/ml

Species	Rat

Dilution info	1/1000
Notes	0.502 ug/ml

Not recommended

Species	Human
Dilution info	-
Notes	0.502 ug/ml

IHC-Fr

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Species	Rat
Dilution info	1/50
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/8000

Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human
Dilution info	-
Notes	-

Target data

See full target information Basp1

Storage

Shipped at conditions	Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


13 product images



Immunocytochemistry/ Immunofluorescence - Anti-BASP1 antibody [EPR28066-51] (ab312324)

BASP1 Immunocytochemistry/ Immunofluorescence staining of mouse primary neuron using rabbit Anti-BASP1 antibody


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Tween-20 permeabilized mouse primary neuron cells labelling BASP1 with ab312324 at 1/1000 (0.502 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green). The confocal image showing cytoplasmic staining in mouse primary neuron.Confocal scanning Z step was set as 0.3 µm followed by image processing with maximum Z projection. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. ab11267 Anti-MAP2 mouse monoclonal antibody was used to counterstain tubulin at 1/500 4ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

BASP1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse cardiac muscle using rabbit Anti-BASP1 antibody

Immunohistochemical analysis of paraffin-embedded Mouse cardiac muscle tissue labeling BASP1 with ab312324 at 1/8000 (0.063 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on mouse cardiac muscle.The section was incubated with ab312324 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrumentIncubate slides with 3% Hydrogen Peroxide for 10 mins at room temperature after secondary antibody incubation to reduce the background Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Frozen sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

BASP1 Immunohistochemistry (Frozen sections) staining of Rat skeletal muscle (fresh) using rabbit Anti-BASP1 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat skeletal muscle (fresh) tissue labeling BASP1 with ab312324 at 1/50 (10.04 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Negative control: confocal image showing no staining on rat skeletal muscle


(PMID: 2148567). The nuclear counterstain was DAPI (Blue). The section was incubated with ab312324 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.



Immunohistochemistry (Frozen sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

BASP1 Immunohistochemistry (Frozen sections) staining of Rat liver (fresh) using rabbit Anti-BASP1 antibody


Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat liver (fresh) tissue labeling BASP1 with ab312324 at 1/50 (10.04 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Negative control: confocal image showing no staining on rat liver (PMID: 14701728). The nuclear counterstain was DAPI (Blue). The section was incubated with ab312324 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.



Immunohistochemistry (Frozen sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

BASP1 Immunohistochemistry (Frozen sections) staining of Mouse liver (fresh) using rabbit Anti-BASP1 antibody

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse liver (fresh) tissue labeling BASP1 with ab312324 at 1/50 (10.04 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Negative control: confocal image showing no staining on mouse liver (PMID: 14701728). The nuclear counterstain was DAPI (Blue). The section was incubated with ab312324 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.




Immunohistochemistry (Frozen sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

BASP1 Immunohistochemistry (Frozen sections) staining of Mouse skeletal muscle (fresh) using rabbit Anti-BASP1 antibody


Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse skeletal muscle (fresh) tissue labeling BASP1 with ab312324 at 1/50 (10.04 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Negative control: confocal image showing no staining on mouse skeletal muscle (PMID: 2148567). The nuclear counterstain was DAPI (Blue). The section was incubated with ab312324 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

Immunohistochemical analysis of paraffin-embedded Mouse cerebrum tissue labeling BASP1 with ab312324 at 1/8000 (0.063 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse cerebrum.The section was incubated with ab312324 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrumentIncubate slides with 3% Hydrogen Peroxide for 10 mins at room temperature after secondary antibody incubation to reduce the background Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Frozen sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat cerebrum (fresh) tissue labeling BASP1 with ab312324 at 1/50 (10.04 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Confocal image showing positive staining on rat cerebrum. The nuclear counterstain was DAPI (Blue). The section was incubated with ab312324 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.




Immunohistochemistry (Frozen sections) - Anti-BASP1 antibody [EPR28066-51] (ab312324)

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Mouse cerebrum (fresh) tissue labeling BASP1 with ab312324 at 1/50 (10.04 ug/ml) dilution followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/mL dilution (Green). Confocal image showing positive staining on mouse cerebrum. The nuclear counterstain was DAPI (Blue). The section was incubated with ab312324 for 60 mins at room temperature. The section was then mounted using Fluoromount®.The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). is observed. The nuclear counterstain was DAPI (Blue). Secondary antibody control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbedat 1/1000 2 ug/mL dilution.




Flow Cytometry - Anti-BASP1 antibody [EPR28066-51] (ab312324)

Flow cytometric analysis of Mouse spleen cell cells labelling BASP1 with ab312324 at 1/50 dilution (1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Gated on viable cells.



Flow Cytometry - Anti-BASP1 antibody [EPR28066-51] (ab312324)

Flow cytometric analysis of Mouse primary neuron cells labelling BASP1 with ab312324 at 1/50 dilution (1 ug)/Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Gated on viable cells.



Immunoprecipitation - Anti-BASP1 antibody [EPR28066-51] (ab312324)


BASP1 was immunoprecipitated from 0.35 mg Mouse brain tissue lysate with ab312324 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab312324 at 1/1000 dilution. VeriBlot for IP secondary antibody(HRP)(ab131366) was used at 1/5000 dilution. Lane 1: Mouse brain tissue lysate Lane 2: ab312324 IP in Mouse brain tissue lysate Lane 3:Rabbit monoclonal IgG (ab172730) instead of ab312324 in mouse brain tissue lysate Blocking and dilution buffer and concentration: 5% NFDM/TBST. Exposure time: 180 seconds.
All lanes:
Immunoprecipitation - Anti-BASP1 antibody [EPR28066-51] (ab312324) at 1/30 dilution

All lanes:
Mouse brain tissue lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Exposure time: 180s



Western blot - Anti-BASP1 antibody [EPR28066-51] (ab312324)

Blocking and diluting buffer and concentration: 5% NFDM/TBST Negative control: kidney, lung, heart, pancreas, liver (PMID: 9749536, 14701728, 8390468). In Western blot, anti-GAPDH antibody (ab181602) loading control staining at 1/200000 dilution. Exposure time: 180 seconds
All lanes:
Western blot - Anti-BASP1 antibody [EPR28066-51] (ab312324) at 1/1000 dilution

Lane 1:
Mouse brain tissue lysate at 20 µg

Lane 2:
Mouse liver tissue lysate at 20 µg

Lane 3:

Mouse kidney tissue lysate at 20 µg

Lane 4:
Mouse lung tissue lysate at 20 µg

Lane 5:
Mouse heart tissue lysate at 20 µg

Lane 6:
Mouse pancreas tissue lysate at 20 µg

Lane 7:
Rat brain tissue lysate at 20 µg

Lane 8:
Rat liver tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 55 kDa

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.