abcam

ab312840

Anti-Sialoadhesin/CD169 antibody [EPR27102-11]

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR27102-11
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/250
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human, Transfected cell line - Mouse
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Not recommended

Species	Human, Transfected cell line - Mouse
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Human, Transfected cell line - Mouse
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/100
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Transfected cell line - Mouse
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Human
Dilution info	-
Notes	-

mIHC

Tested

Species	Mouse
Dilution info	1/100
Notes	-

Species	Rat
Dilution info	1/100
Notes	-

Not recommended

Species	Human, Transfected cell line - Mouse
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Rat, Human, Transfected cell line - Mouse
Dilution info	-
Notes	-

Target data

See full target information Siglec1

Function

Macrophage-restricted adhesion molecule that mediates sialic-acid dependent binding to lymphocytes, including granulocytes, monocytes, natural killer cells, B-cells and CD8 T-cells. Plays a crucial role in limiting bacterial dissemination by engaging sialylated bacteria to promote effective phagocytosis and antigen presentation for the adaptive immune response (PubMed:24788876). Mediates the uptake of various enveloped viruses via sialic acid recognition and subsequently induces the formation of intracellular compartments filled with virions (VCCs) (PubMed:26370074). In turn, enhances macrophage-to-T-cell transmission of several viruses including murine leukemia virus (PubMed:26370074). Acts as an endocytic receptor mediating clathrin dependent endocytosis. Preferentially binds to alpha-2,3-linked sialic acid (By similarity). Binds to SPN/CD43 on T-cells (By

similarity). May play a role in hemopoiesis. Plays a role in the inhibition of antiviral innate immune by promoting TBK1 degradation via TYROBP and TRIM27-mediated ubiquitination (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


14 product images



Multiplex immunohistochemistry - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Multiplex immunohistochemistry staining of Mouse spleen using rabbit Anti-Sialoadhesin/CD169 antibody


Multiplex immunohistochemistry analysis of formalin/PFA-fixed paraffin-embedded Mouse spleen tissue labeling Sialoadhesin/CD169, CD3 epsilon and CD20 with ab312840 at 1/100 dilution, ab16669 at 1:150 dilution and ab236434 at 1:5000 dilution. Panel A: merged staining of anti-Sialoadhesin/CD169 (red; Opal™690), anti-CD3 epsilon (green; Opal™520) and anti-CD20 (gray; Opal™570) on mouse spleen. Panel B: anti-Sialoadhesin/CD169 stained on macrophages. Panel C: anti-CD3 epsilon stained on T cells. Panel D: anti-CD20 stained on B cells. The section was incubated in three rounds of staining: in the order of ab312840, ab16669, and ab236434 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system. The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Rat kidney using rabbit Anti-Sialoadhesin/CD169 antibody


Immunohistochemical analysis of paraffin-embedded Rat kidney tissue labeling Sialoadhesin/CD169 with ab312840 at 1/100 (5.16 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on rat kidney. The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse kidney using rabbit Anti-Sialoadhesin/CD169 antibody


Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labeling Sialoadhesin/CD169 with ab312840 at 1/100 (5.16 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on mouse kidney. The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)


Sialoadhesin/CD169 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Mouse skeletal muscle using rabbit Anti-Sialoadhesin/CD169 antibody

Immunohistochemical analysis of paraffin-embedded Mouse skeletal muscl tissue labeling Sialoadhesin/CD169 with ab312840 at 1/100 (5.16 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on mouse skeletal muscle. The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Immunohistochemical analysis of paraffin-embedded (A) HEK-293T (human tissue labeling Sialoadhesin/CD169 with ab312840 at 1/2000 (0.258 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on (A) HEK-293T transfected with a mouse SIGLEC1 expression vector containing a his tag. No staining on (B) HEK-293T transfected with empty vector containing a his tag. The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling Sialoadhesin/CD169 with ab312840 at 1/100 (5.16 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse spleen interfollicular zone (PMID: 15107454). The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Immunohistochemical analysis of paraffin-embedded Rat spleen tissue labeling Sialoadhesin/CD169 with ab312840 at 1/100 (5.16 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat spleen interfollicular zone. The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.




Immunocytochemistry/ Immunofluorescence - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized RAW 264.7 (mouse abelson murine leukemia virus-induced tumor macrophage) cells labelling Sialoadhesin/CD169 with ab312840 at 1/250 (2.064 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 (2 ug/ml) dilution (Green). Confocal image showing positive staining in RAW 264.7 cells treated with lipopolysaccharide (100ng/ml) for 16 hours.Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 (2.5ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 ug/ml) dilution.




Flow Cytometry - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Flow cytometric analysis of Mouse bone marrow cells labelling Sialoadhesin/CD169 with ab312840 at 1/50 dilution (1ug)/ Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Cells co-stained with CD3 conjugated to A647. Gated on viable cells.



Flow Cytometry - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)


Flow cytometric analysis of Mouse bone marrow cells labelling Sialoadhesin/CD169 with ab312840 at 1/50 dilution (1ug)/ Right compared with a Rabbit monoclonal IgG (ab172730) / Left isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Cells co-stained with F4/80 conjugated to PE. Gated on viable cells.



Multiplex immunohistochemistry - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Multiplex immunohistochemistry staining of Rat spleen using rabbit Anti-Sialoadhesin/CD169 antibody


Multiplex immunohistochemistry analysis of formalin/PFA-fixed paraffin-embedded Rat spleen tissue labeling Sialoadhesin/CD169,CD3 epsilon and CD20 with ab312840 at 1/100 dilution, ab16669 at 1:150 dilution and ab236434 at 1:5000 dilution. Panel A: merged staining of anti-Sialoadhesin/CD169 (red; Opal™690), anti-CD3 epsilon (green; Opal™520) and anti-CD20 (gray; Opal™570) on rat spleen. Panel B: anti-Sialoadhesin/CD169 stained on macrophages. Panel C: anti-CD3 epsilon stained on T cells. Panel D: anti-CD20 stained on B cells. The section was incubated in three rounds of staining: in the order of ab312840, ab16669, and ab236434 for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system. The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Rat skeletal muscle using rabbit Anti-Sialoadhesin/CD169 antibody

Immunohistochemical analysis of paraffin-embedded Rat skeletal muscle tissue labeling Sialoadhesin/CD169 with ab312840 at 1/100 (5.16 ug/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Negative control: no staining on rat skeletal muscle. The section was incubated with ab312840 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Western blot - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Western blot staining using rabbit Anti-Sialoadhesin/CD169 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST The expression of Sialoadhesin/CD169 is upregulated in response to LPS treatment (PMID: 27129263). In Western blot, anti- Vinculin antibody (ab129002) loading control staining at 1/10000 dilution. Exposure time: 180 seconds
All lanes:
Western blot - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840) at 1/1000 dilution

Lane 1:
Untreated RAW 264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage) whole cell lysate at 20 µg


Lane 2:
RAW 264.7 treated with /ml LPS for 16h whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 250 kDa

Exposure time: 180s



Western blot - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840)

Sialoadhesin/CD169 Western blot staining using rabbit Anti-Sialoadhesin/CD169 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST Low expression: kidney (PMID 33408860). This blot of Lane 6 was developed using a high sensitivity ECL substrate. In Western blot, anti- Vinculin antibody (ab129002) loading control staining at 1/10000 dilution. Exposure time: Lane 1-5: 180 seconds; Lane 6: 114 seconds.
All lanes:
Western blot - Anti-Sialoadhesin/CD169 antibody [EPR27102-11] (ab312840) at 1/1000 dilution

Lane 1:
Mouse spleen tissue lysate at 20 µg

Lane 2:
Mouse lymph node tissue lysate at 20 µg

Lane 3:
Mouse kidney tissue lysate at 20 µg

Lane 4:
Rat lymph node tissue lysate at 20 µg

Lane 5:
Rat kidney tissue lysate at 20 µg

Lane 6:
Rat spleen tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 250 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.