abcam

ab313392

Anti-EZHIP antibody [EPR28396-50]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR28396-50
Purification technique	Affinity purification Protein A
Specificity	Unsuitable for mouse IHC-P.
Concentration	0.52 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/200
Notes	

Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Mouse, Rat
Dilution info	-

Notes	-

Target data

See full target information EZHIP

Function	Inhibits PRC2/EED-EZH1 and PRC2/EED-EZH2 complex function by inhibiting EZH1/EZH2 methyltransferase activity, thereby causing down-regulation of histone H3 trimethylation on 'Lys-27' (H3K27me3) (PubMed:29909548, PubMed:30923826, PubMed:31086175, PubMed:31451685). Probably inhibits methyltransferase activity by limiting the stimulatory effect of cofactors such as AEBP2 and JARID2 (PubMed:30923826). Inhibits H3K27me3 deposition during spermatogenesis and oogenesis (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The EZHIP (EZH Inhibitory Protein) also known as CATACOMB is a protein that inhibits the activity of EZH2 an important enzyme in the PRC2 (Polycomb Repressive Complex 2). EZHIP is a protein with a molecular mass of approximately 90 kDa. It expresses in various tissues with notable presence in the male and female reproductive tracts as well as in several cancer cell lines. The protein's role involves modulating chromatin structure by interfering with methyltransferase activity affecting gene silencing.
Biological function summary	EZHIP interacts with the Polycomb Repressive Complex 2 (PRC2) altering its methylation activity on histone H3 at lysine 27 (H3K27me3) an important mark for epigenetic gene silencing. By engaging the complex EZHIP hinders the catalytic efficiency of EZH2 reducing the repression of target genes. This interaction significantly impacts epigenetic regulation as alterations to methylation patterns influence cellular functions and differentiation processes.
Pathways	EZHIP's function affects pathways related to chromatin modification and epigenetic regulation. It is central to the regulation of the JAK-STAT signaling pathway and interacts closely with EZH2 and possibly other members of the Polycomb group proteins. EZHIP shapes the chromatin landscape influencing the transcriptional output by collaborating or competing with other transcription factors and modifiers within these pathways.
Associated diseases and disorders	EZHIP's aberrant expression is linked to cancers including gliomas and germ cell tumors. The relationship between EZHIP and EZH2 becomes important as alterations in EZH2 activity lead to abnormal gene silencing associated with oncogenesis. Additionally research indicates that dysregulation involving EZHIP might contribute to developmental disorders through improper epigenetic marks. Further study about its interactions could provide insights into therapeutic approaches for conditions linked to epigenetic misregulation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


8 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EZHIP antibody [EPR28396-50] (ab313392)

EZHIP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human skeletal muscl using rabbit Anti-EZHIP antibody


Immunohistochemical analysis of paraffin-embedded Human skeletal muscl tissue labeling CXorf67 with ab313392 at 1/200 (2.555 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Negative control: No staining on human skeletal muscle.The section was incubated with ab313392 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EZHIP antibody [EPR28396-50] (ab313392)

EZHIP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human cardiac muscle using rabbit Anti-EZHIP antibody


Immunohistochemical analysis of paraffin-embedded Human cardiac muscle tissue labeling CXorf67 with ab313392 at 1/200 (2.555 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Negative control: No staining on human cardiac muscle.The section was incubated with ab313392 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EZHIP antibody [EPR28396-50] (ab313392)


Immunohistochemical analysis of paraffin-embedded Human testis tissue labeling CXorf67 with ab313392 at 1/200 (2.555 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human testis (PMID: 31451685).The section was incubated with ab313392 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody

only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EZHIP antibody [EPR28396-50] (ab313392)

Immunohistochemical analysis of paraffin-embedded Human placenta tissue labeling CXorf67 with ab313392 at 1/200 (2.555 ug/ml) followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on human placenta.The section was incubated with ab313392 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin. Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunocytochemistry/ Immunofluorescence - Anti-EZHIP antibody [EPR28396-50] (ab313392)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized U-2 OS (human bone osteosarcoma epithelial cell) cells labelling CXorf67 with ab313392 at 1/500 (1.022 ug/ml) dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 2 ug/ml dilution (Green). Confocal image showing nuclear staining in U-2 OS cell line.Low expression: 293T (PMID: 34880314).Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8). ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5ug/ml dilution (Red). The Nuclear counterstain was DAPI (Blue). Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 2 ug/ml dilution.



Flow Cytometry (Intracellular) - Anti-EZHIP antibody [EPR28396-50] (ab313392)

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized 293T (human embryonic kidney epithelial cell, Left) / U-2 OS (human bone osteosarcoma epithelial cell, Right) cells labelling EZHIP with ab313392 at 1/500 dilution (0.1 ug)/Red (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) at 1/5000 dilution was used as the secondary antibody. Low expression: 293T (PMID: 34880314).



Western blot - Anti-EZHIP antibody [EPR28396-50] (ab313392)

EZHIP Western blot staining using rabbit Anti-EZHIP antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Low expression: 293T (PMID: 34880314).

The molecular weight observed is consistent with what has been described in the literature (PMID: 34880314).

Lysates were freshly made and used for Western blotting immediately to minimize protein degradation.

In Western blot, anti-GAPDH antibody (ab181602) loading control staining at 1/200000 dilution.
All lanes:
Western blot - Anti-EZHIP antibody [EPR28396-50] (ab313392) at 1/1000 dilution

Lane 1:
U-2OS (human bone osteosarcoma epithelial cell) whole cell lysate at 50 µg

Lane 2:
293T (human embryonic kidney epithelial cell) whole cell lysate at 50 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Performed under non-reducing conditions.

Observed band size: 50 kDa, 75 kDa

Exposure time: 92s



Western blot - Anti-EZHIP antibody [EPR28396-50] (ab313392)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
This blot was developed using a high sensitivity ECL substrate. The high-sensitivity ECL substrate used allows for the detection of proteins in the mid-femtogram range.

In Western blot, anti- H3 antibody (ab176842) loading control staining at 1/100000 dilution.

In Western blot, anti-GAPDH antibody (ab181602) loading control staining at 1/200000 dilution.

All lanes:
Western blot - Anti-EZHIP antibody [EPR28396-50] (ab313392) at 1/1000 dilution

Lane 1:
Mouse testis cytoplasmic fraction at 50 µg

Lane 2:
Mouse testis nuclear fraction at 50 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Developed using the ECL technique.

Observed band size: 75 kDa

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.