abcam

ab313647

Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR28380-83
Purification technique	Affinity purification Protein A
Concentration	1.027 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Transfected cell lysate - Human
Dilution info	-

Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Transfected cell line - Human, Transfected cell line
Dilution info	-
Notes	-

IP

Tested

Species	Transfected cell lysate - Human
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Transfected cell line - Human, Transfected cell line
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Transfected cell line - Human
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Transfected cell lysate - Human, Transfected cell line
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human, Transfected cell line
Dilution info	-
Notes	-

Not recommended

Species	Transfected cell lysate - Human, Transfected cell line - Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Transfected cell line - Human

Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Transfected cell lysate - Human, Transfected cell line
Dilution info	-
Notes	-

Target data

See full target information EGFR

Function

Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:10805725, PubMed:27153536, PubMed:2790960, PubMed:35538033). Known ligands include EGF, TGFA/TGF-alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF (PubMed:12297049, PubMed:15611079, PubMed:17909029, PubMed:20837704, PubMed:27153536, PubMed:2790960, PubMed:7679104, PubMed:8144591, PubMed:9419975). Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity). Isoform 2 may act as an antagonist of EGF action. (Microbial infection) Acts as a receptor for hepatitis C virus (HCV) in hepatocytes and facilitates its cell entry. Mediates HCV entry by promoting the formation of the

CD81-CLDN1 receptor complexes that are essential for HCV entry and by enhancing membrane fusion of cells expressing HCV envelope glycoproteins.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	EGFR VIII also known as EGFRvIII EGFR of 83 or 83 EGFR is a mutant form of the epidermal growth factor receptor (EGFR) with a molecular weight of approximately 145 kDa. This variant results from a deletion of exons 2 to 7 in the EGFR gene

	which produces a truncated protein that lacks a portion of the extracellular ligand-binding domain. The EGFRvIII does not bind ligands and is constitutively active. It is often expressed in cancer cells particularly in glioblastomas and some other solid tumors but not in normal tissues.
Biological function summary	EGFRvIII modifies signaling pathways that promote cell survival and proliferation. It is a part of cell membrane-based complexes interacting with proteins like PI3K and Src which helps in activating downstream signaling cascades. This constitutive signaling leads to increased oncogenic potential allowing the cells to thrive in uncontrolled manner contributing to tumor progression and resistance to conventional therapies.
Pathways	EGFRvIII participates in key signaling pathways such as the PI3K/AKT pathway and the RAS/RAF/MEK/ERK pathway. These pathways are important in transmitting signals from receptor tyrosine kinases to the cell nucleus. EGFRvIII links with proteins like AKT and ERK to drive processes that control growth and survival. The dysregulation of these pathways due to EGFRvIII activity can result in enhanced proliferative and survival signals in cancer cells.
Associated diseases and disorders	EGFRvIII is strongly associated with glioblastoma and non-small cell lung cancer. In these conditions EGFRvIII expression correlates with poor prognosis and an aggressive tumor phenotype. The presence of EGFRvIII often accompanies other oncogenic proteins such as PTEN whose loss of function can amplify the oncogenic signaling of EGFRvIII. As a result targeting EGFRvIII represents a potential therapeutic intervention in such aggressive cancer types.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

9 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)


EGFRvIII Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human glioblastoma and placenta tissues using rabbit Anti-EGFRvIII antibody

Immunohistochemical analysis of formalin fixed paraffin-embedded human glioblastoma and placenta tissues labelling EGFRvIII with ab313646 and EGFR with ab52894 at a concentration of 2.5 µg/ml and 0.5 µg/ml.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with a Bond™ Polymer Refine Detection kit. Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution 2) for 20 mins. Primary antibodies were incubated for 30 mins at room temperature. Sections were counterstained with Hematoxylin.

Tissue obtained from the Human Research Tissue Bank, supported by the NHR Cambridge Biomedical Research Centre.

This data was developed using ab313646, the same antibody clone in a different buffer formulation.




Immunocytochemistry/ Immunofluorescence - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

This data was developed using ab313646, the same antibody clone in a different buffer formulation.
Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized 293T (human embryonic kidney epithelial cell) labelling EGFRvIII with ab313646 at 1/500 dilution, followed by ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed antibody at 1/1000 dilution (Green). Cells were counterstained with anti-Myc tag Mouse mab [9E10] (Alexa Fluor® 594) at 1/100 dilution (Red). The Nuclear counterstain is DAPI (Blue).

Confocal image showing cytoplasmic and membrane staining in 293T cells overexpressing human EGFRvIII. No staining in 293T cells overexpressing human wild-type EGFR. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Secondary antibody only control: Secondary antibody is ab150081 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 (2 µg/mL) dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

EGFRvIII Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human glioblastoma tissue using rabbit Anti-EGFRvIII antibody


Chromogenic duplex (2-plex) immunohistochemical staining of formalin fixed paraffin-embedded human glioblastoma tissue performed on a Roche Ventana Discovery Ultra instrument.

Heat mediated antigen retrieval was performed with CC1 solution for 64 min at 95°C. Two rounds of primary antibody incubation were performed each for 16 min at 37°C. anti-EGFRvIII (ab313646; 1st primary antibody; 3 µg/ml) and anti-EGFR (ab52894; 2nd primary antibody; 0.5 µg/ml). Antibody denaturation was conducted after the first round of staining using Ultra CC2 solution for 8 min at 100°C to avoid cross reactivity.

Signal was developed with anti-rabbit HQ followed by anti-HQ HRP coupled with purple (Discovery purple kit (RUO) #760-229) and teal (Discovery teal kit (RUO) #760-247) chromogens and counterstained with haematoxylin II.

Tissue obtained from the Human Research Tissue Bank, supported by the NHR Cambridge Biomedical Research Centre.

This data was developed using ab313646, the same antibody clone in a different buffer formulation.




Flow Cytometry - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

EGFRvIII Flow Cytometry staining using rabbit Anti-EGFRvIII antibody

This data was developed using ab313646, the same antibody clone in a different buffer formulation.
Flow cytometric analysis of 293T (human embryonic kidney epithelial cell) transfected with a human EGFRvIII expression vector containing a His-Myc tag (top left, bottom left); human wild-type EGFR expression vector containing a His-Myc tag (top right); empty vector containing a His-Myc tag (bottom right). Cells were fixed with 2% paraformaldehyde and permeabilised with 0.1% Tween-20. Cell labelling of EGFRvIII with ab313646 at 1/500 dilution. A Goat Anti-Rabbit IgG (Alexa Fluor® 488, ab150081) was used at 1/5000 dilution. Isotype control - Rabbit monoclonal IgG (ab172730). No cross-reactivity with human wild-type EGFR.

Cells were surface stained with primary antibody. Fixed with 2% PFA for 10 min followed by intracellularly staining with anti-Myc tag conjugated to Alexa Fluor® 647.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)


This data was developed using ab313646, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin-embedded 293T (human embryonic kidney epithelial cell) transfected with (A) human EGFRvIII expression vector containing a Myc-His tag; (B) human wild-type EGFR expression vector containing a Myc-His tag; (C) empty vector containing a His tag.

Whole cell pellet labelling of EGFRvIII with ab313646 at 1/100 (5.09 µg/ml) followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Positive staining on 293T overexpressing human EGFRvIII (image A). No staining on 293T overexpressing human wild-type EGFR (image B) or the empty vector control (image C). The section was incubated with ab313646 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

This data was developed using ab313646, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin-embedded (A, D) human placenta; (B, E) human glioblastoma; (C, F) human lung adenocarcinoma tissue.

A-C: Sections were incubated with anti-EGFR antibody [SP84] - C-terminal (ab227642; 1:100) for 30 mins at room temperature. Positive staining on human placenta (image A), glioblastoma (image B), and lung adenocarcinoma (image C). D-F: Sections were incubated with anti-EGFRvIII antibody [EPR28380-83] (ab313646; 1:100 [5.09 µg/mL]) for 30 mins at room temperature. Positive staining on human glioblastoma (image E), no staining on placenta (image D) or lung adenocarcinoma (image F). The immunostaining was performed on a Leica Biosystems BOND® RX instrument counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunoprecipitation - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

This data was developed using ab313646, the same antibody clone in a different buffer formulation.
EGFRvIII was immunoprecipitated from 0.35 mg of 293T (human embryonic kidney epithelial cell) transfected with a human EGFRvIII expression vector containing a His-tag using ab313646 at 1/30 dilution (2µg in 0.35mg whole cell lysate). Western blot was performed on the immunoprecipitate using ab313646 at 1/1000 dilution. VeriBlot for IP secondary antibody (HRP)(ab131366) was used at 1/5000 dilution. Isotype control - Rabbit monoclonal IgG (ab172730).

Lanes 1-3: 293T (human embryonic kidney epithelial cell) transfected with a human EGFRvIII expression vector containing a His-tag

Lane 1: whole cell lysate at 0.3µg

Lane 2: IP with ab313646

Lane 3: IP with Isotype control (ab172730)


Blocking and dilution buffer and concentration: 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-EGFRvIII antibody [EPR28380-83] (ab313646) at 1/30 dilution

All lanes:
293T (human embryonic kidney epithelial cell) transfected with a human EGFRvIII expression vector containing a His-tag whole cell lysate

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/1000 dilution

Exposure time: 5s



Western blot - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

EGFRvIII Western blot staining using rabbit Anti-EGFRvIII antibody

This data was developed using ab313646, the same antibody clone in a different buffer formulation. This antibody recognizes human EGFRvIII but not human wild-type EGFR. In western blot, anti-GAPDH antibody [EPR16891] - Loading Control (ab181602) staining at 1/200000 dilution, anti-His antibody (ab213204) staining at 1/5000 dilution, and anti-EGFR antibody staining at 1/1000 dilution.
All lanes:
Western blot - Anti-EGFRvIII antibody [EPR28380-83] (ab313646) at 1/1000 dilution

Lane 1:
293T (human embryonic kidney epithelial cell) transfected with an empty vector containing a His tag, whole cell lysate at 10 µg

Lane 2:
293T transfected with a human EGFR expression vector containing a His tag, whole cell lysate at 10 µg

Lane 3:
293T transfected with a human EGFRvIII expression vector containing a His tag, whole cell lysate at 10 µg

Lane 4:
293T transfected with a human EGFRvIII expression vector containing a His tag, whole cell lysate at 2.5 µg

Lane 5:
293T transfected with a human EGFRvIII expression vector containing a His tag, whole cell lysate at 0.62 µg

Lane 6:
293T transfected with a human EGFRvIII expression vector containing a His tag, whole cell lysate at 0.31 µg


Lane 7:
293T transfected with a human EGFRvIII expression vector containing a His tag, whole cell lysate at 0.16 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Observed band size: 134 kDa

Exposure time: 5.5s



Western blot - Anti-EGFRvIII antibody [EPR28380-83] - BSA and Azide free (ab313647)

This data was developed using ab313646, the same antibody clone in a different buffer formulation. Blocking and diluting buffer and concentration: 5% NFDM/TBST This antibody does not recognize human wild-type EGFR. (A): anti-EGFRvIII antibody (ab313646) staining at 1/1000 dilution, (B): anti-EGFR antibody staining at 1/1000 dilution, (C): anti-GAPDH antibody [EPR16891] – Loading Control (ab181602) staining at 1/200000 dilution.
All lanes:
Western blot - Anti-EGFRvIII antibody [EPR28380-83] (ab313646) at 1/1000 dilution

Lane 1:
A431 (human epidermoid carcinoma epithelial cell) whole cell lysate at 20 µg

Lane 2:
Wild-type HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate at 20 µg

Lane 3:
EGFR knockout HeLa whole cell lysate at 20 µg

Lane 4:
Human placenta tissue lysate at 20 µg

Secondary

Lanes 1 - 3:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Lane 4:
Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Observed band size: 134 kDa

Exposure time: 180s

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.